

This is the peer reviewed version of the following article: Inauen, J. , Bolger, N. , Shrout, P. E., Stadler, G. , Amrein, M. , Rackow, P. and Scholz, U. (2017), Using Smartphone-Based Support Groups to Promote Healthy Eating in Daily Life: A Randomised Trial. *Applied Psychology: Health and Well-Being*, 9: 303-323, which has been published in final form at <https://doi.org/10.1111/aphw.12093>. This article may be used for non-commercial purposes in accordance With Wiley Terms and Conditions for self-archiving.

Running Head: SMARTPHONE SUPPORT GROUPS

**Using Smartphone-Based Support Groups to Promote Healthy Eating in Daily Life:
A Randomized Trial**

Jennifer Inauen, Ph.D., Corresponding author

Department of Environmental Social Sciences, Eawag: Swiss Federal Institute of Aquatic
Science & Technology, Duebendorf, Switzerland and

Department of Psychology, Columbia University, New York, USA

Niall Bolger, Ph.D.

Department of Psychology, Columbia University, New York, USA

Patrick E. Shrout, Ph.D.

Department of Psychology, New York University, New York, USA

Gertraud Stadler, Ph.D.

The Institute of Applied Health Sciences, University of Aberdeen, Aberdeen, Scotland

Melanie Amrein, MSc

Department of Psychology, University of Zurich, Zurich, Switzerland

Pamela Rackow, Ph.D.

The Institute of Applied Health Sciences, University of Aberdeen, Aberdeen, Scotland

Urte Scholz, Ph.D.

Department of Psychology & University Research Priority Program “Dynamics of Healthy
Aging”, University of Zurich, Zurich, Switzerland

Acknowledgement

The first author was supported by a fellowship of the Swiss National Science
Foundation (Fellowship P2ZHP1_155103).

Conflict of interest

The authors declare that they have no conflict of interest.

Authors

Jennifer Inauen, Eawag: Swiss Federal Institute of Aquatic Science & Technology,
Department of Environmental Social Sciences, Environmental and Health Psychology,
Ueberlandstrasse 133, 8600 Duebendorf, Switzerland, phone: +41 58 765 5780, email:
jennifer.inauen@eawag.ch

Niall Bolger, Columbia University, Department of Psychology, 402A Schermerhorn Hall,
1190 Amsterdam Avenue, New York, NY 10027, USA, phone: +1 212 854 9034, email:
bolger@psych.columbia.edu

Patrick E. Shrout, New York University, Department of Psychology, 6 Washington Place,
New York, NY 10003, USA, phone: +1 212 998 7895, email: pat.shrout@nyu.edu

Gertraud Stadler, University of Aberdeen, The Institute of Applied Health Sciences, 2nd
Floor Health Sciences Building, Foresterhill Campus, Ashgrove Road, Aberdeen AB25 2ZD,
United Kingdom, phone: +44 1224 438407, email: gertraud.stadler@abdn.ac.uk

Melanie Amrein, University of Zurich, Department of Psychology, Applied Social and Health
Psychology, Binzmuehlestrasse 14, Box 14, 8050 Zurich, Switzerland, phone: +41 44 635 72
60, email: melanie.amrein@uzh.ch

Pamela Rackow, University of Aberdeen, The Institute of Applied Health Sciences, 2nd Floor
Health Sciences Building, Foresterhill Campus, Ashgrove Road, Aberdeen AB25 2ZD,
United Kingdom, phone: +44 1224 438108, email: pamela.rackow@abdn.ac.uk

Urte Scholz, University of Zurich, Department of Psychology, Applied Social and Health
Psychology, Binzmuehlestrasse 14, Box 14, 8050 Zurich, Switzerland, phone: +41 44 635 72
50, email: urte.scholz@psychologie.uzh.ch

Using Smartphone-Based Support Groups to Promote Healthy Eating in Daily Life: A Randomized Trial

Abstract

Background: Although many people intend to eat healthily, they often fail to do so. We report the first randomized trial testing if smartphone-based support groups can enhance healthy eating. **Methods:** Adults ($N=203$) were randomized to the support or control condition (information), and to one of two eating goals (increasing fruit and vegetable / decreasing unhealthy snack consumption). After baseline, participants received information on their assigned eating goal, and completed a 13-day electronic diary. During Days 4-10, support participants were asked to support each other in achieving their eating goal in smartphone-based groups. The primary outcome was daily servings of fruits/vegetables or unhealthy snacks. Maintenance of intervention effects was assessed on Days 11-13, and at 1-month and 2-month follow-ups. **Results:** Support participants showed a gradual increase in healthy eating over time, and ate 1.4 fruits and vegetables more, 95% CI [0.3, 2.6], or 0.8 unhealthy snacks less, 95% CI [-1.4, -0.2] than controls on Day 10. Most effects were not maintained at follow-ups. **Conclusions:** Smartphone-based groups can promote fruit and vegetable consumption and decrease unhealthy snack intake. This study extends previous findings of the benefits of support groups, and sheds light on the temporal dynamics of behavior change.

Keywords: Eating behavior; Fruits and vegetables; Unhealthy snacks; Social support groups; Intensive longitudinal methods

What we eat influences our health. To prevent chronic disease, current dietary guidelines include increasing fruit and vegetable consumption, and reducing fat and sugar intake (U.S. Department of Health and Human Services and U.S. Department of Agriculture, 2015). Higher fruit and vegetable consumption has been found to reduce risks of cancer (Turati, Rossi, Pelucchi, Levi, & La Vecchia, 2015) and coronary artery disease (Brunzell, 2007), and is associated with delayed mortality (Wang et al., 2014). Unhealthy snack consumption, which is the consumption of energy dense foods between main meals, has been linked to increased energy intake (Graaf, 2006), and increased risk of being overweight and obese (Murakami & Livingstone, 2016). Although people are often motivated to eat healthily, they frequently fail to do so in their daily lives. This so-called intention-behavior gap may be due to self-regulation failures during certain times of day when temptation is high or in circumstances when persons can rationalize setting aside their healthy eating goals (Inauen, Shrout, Bolger, Stadler, & Scholz, 2016).

One important influence on self-regulation success is social support. Social support has been defined as “a set of processes through which another person helps individuals engage in effective self-regulation” (Fitzsimons & Finkel, 2010, p. 103) to accomplish a particular goal. The association of social support with physical health is well documented (Uchino, Bowen, Carlisle, & Birmingham, 2012), and an increasing body of research indicates that social support may promote health behavior change. In correlational studies, social support is associated with healthy eating (McKinley, 2009), such as eating a low-fat diet (Scholz, Ochsner, Hornung, & Knoll, 2013). Yet, intervention studies of social support have not shown strong effects. One partner-based intervention found small and marginally significant effects (Prestwich et al., 2014), and others found no effects on eating behavior (Anderson Steeves, Jones-Smith, Hopkins, & Gittelsohn, 2016; Kwan & Gordon, 2016) or eating-related behavioral intentions (McKinley & Wright, 2014).

One reason that some support interventions are not shown to be effective is that the quality of support provided by a partner, family member or peer is not uniformly high (Kiernan et al., 2012). A promising alternative to spontaneous support is the formation of specially constructed support groups. Helgeson and Gottlieb (2000) define support groups to be a group of peers with a common goal or experience that includes a moderator or coordinator, exist for a fixed period and do not engage in advocacy. These features distinguish support groups from self-help groups. Support groups with moderators have been shown to be more effective than unmoderated groups (Mohr, Burns, Schueller, Clarke, & Klinkman, 2013).

Because group members share a common goal, they often promote empathy and acceptance. They can also reinforce self-efficacy and provide new information, such as ways to achieve the common goal. Such support groups are increasingly being administered online and on smartphones, harnessing the popularity of social media, such as WhatsApp, Twitter, and Facebook (Rains, Peterson, & Wright, 2015). They are low cost and promising for eating behavior change, which happens in response to many environmental cues in daily life (Papies & Hamstra, 2010) where offline interventions may not be readily accessible. Particular times of day are especially prone to self-regulation failure (Inauen et al., 2016). Therefore, support through a smartphone-based support group (i.e., ecological momentary interventions; Heron & Smyth, 2010) seems especially promising to help regulate eating.

Ecological momentary interventions for weight regulation are increasingly being used and scientifically investigated (Bardus, Smith, Samaha, & Abraham, 2015; Williams, Hamm, Shulhan, Vandermeer, & Hartling, 2014). They have successfully promoted healthy eating using text messaging (Joo & Kim, 2007) or self-monitoring applications, such as diaries where participants keep track of their food consumption (Atienza, King, Oliveira, Ahn, & Gardner, 2008). Commercial dieting programs have also started harnessing the availability of social media on mobile phones and the possibility to deliver group support to regulate eating

in people's daily lives (e.g., Weight Watchers®, Ballantine & Stephenson, 2011). Yet, the experimental evidence on their ability to promote health behavior is scarce. To the best of our knowledge, the effects of smartphone-based support groups on healthy eating have not been experimentally tested.

Smartphone-Based Support Groups to Promote Healthy Eating

In this article, we report results of a randomized trial to test a smartphone-based support group intervention to promote healthy eating. Participants who were randomly allocated to the intervention condition were assigned to a small group that included a trained member who provided frequent support. Group members communicated with each other via WhatsApp, a popular smartphone application that allows for the private exchange of multimedia content via the phone's internet connection, and further internet-connected devices. Such support groups should effectively promote healthy eating because, first, social support provided by a group of persons sharing the same eating goals should provide beneficial effects by meeting participants' needs in terms of timing and content. Second, it can promote self-efficacy by allowing group members to provide support in addition to receiving it. Third, the messages exchanged within the group should serve as reminders of the shared eating goals. Fourth, the group can provide informational support by sharing healthy recipes and information about purveyors of healthy food and ingredients.

The moderator in each group was not identified as a leader, but was rather introduced as another peer participant. We used this design to create a safe space where members with similar behavioral goals could share their experiences and interact as peers. The trained support moderator provided timely supportive responses to every post in the group, and served as a role model of support provision for other group members. We hypothesized that participants in the social support group compared to a control group would eat a healthier diet (with more servings of fruit/vegetables or less unhealthy snacking) by the end of the intervention period.

Besides the overall effectiveness of this approach to promote healthy eating, we are interested in gaining understanding of the temporal development of behavior change in daily life. For this purpose, this study uses an intensive longitudinal design that provides a rare opportunity to investigate how the effects of the social support condition on healthy eating may develop over time. To the best of our knowledge, no temporal theory of behavior change exists to date. The opportunity to look at temporal dynamics of intervention effects is also often missed, even when temporal data is being collected (Brookie, Mainvil, Carr, Vissers, & Conner, 2017). To make a first contribution to understanding the temporal dynamics of the effects of social support on eating behavior change, we will explore two temporal effects. First, it is possible that groups of strangers first have to get to know each other before gradually increasing their support behavior. Second, the invitation to join a smartphone-based support group might have an immediate impact on eating.

Finally, we seek to determine whether this form of social support can promote persistent health behavior change or whether the effects fade after the group support ends. The previous literature on social support suggests such maintenance effects (Kwasnicka, Dombrowski, White, & Sniehotta, 2016; White & Dorman, 2001). However, this has mostly been shown for romantic couples, where support is usually ongoing. The present study, in contrast, provides the rare opportunity to investigate what happens when support ends.

Methods

We conducted a randomized factorial trial of a social support vs. a control condition by two eating goals: eating more fruits and vegetables or eating fewer unhealthy snacks. Participants were randomly allocated to the support or control condition. Furthermore, they were randomly assigned to one of two eating goals: Eating more fruits and vegetables or eating fewer unhealthy snacks. Data was collected using an intensive longitudinal assessment: 3 days prior to the intervention (Days 1-3), 7 days during (Days 4-10), and 3 days after the intervention (Days 11-13). Three panel surveys additionally served to assess participants'

baseline characteristics (T1), and to determine the maintenance of behavior change at 1-month (T2) and 2-month follow-ups (T3). This trial was approved by the Internal Review Board of the University of Zurich. Also, the data analyses of the trial was registered on the Open Science Framework available here: osf.io/9v73s).

Participants and Procedures

Adult staff and student members of the University of Zurich were recruited in September 2014 via flyers, email notifications, and social networks. The recruitment materials aimed at attracting persons with an intention-behavior gap regarding eating, using the heading “Do you intend to eat healthily but find that difficult sometimes?”. Participants were included if they were at least 18 years old, currently not dieting nor participating in a weight loss program, had a Body Mass Index (BMI) of at least 18, owned a smartphone with internet access, and were fluent in German.

The sample size was a priori determined to detect a small to medium effect ($d=0.35$) of social support on healthy eating using an independent samples t-test with 80% power. Using G*Power (Faul, Erdfelder, Lang, & Buchner, 2007) we determined that a total sample size of 204 participants was needed, assuming a two-tailed Type I error probability of .05. Because our pilot study suggested up to 15% dropout, we aimed at recruiting a total of 236 participants to obtain an effective sample of 204 persons.

Prior to participants’ individual appointments at the University, a research assistant randomized them to the conditions by entering their names in the sequence that they signed up for the study into a list of block-randomized cells (with block size 8) that was created by random number generation. As shown in Figure 1, 232 participants were randomly allocated to the conditions.

>>>>> ADD FIGURE 1 HERE <<<<<<

Participants and the interventionists were blind to condition until the participant’s appointment at the lab, where written informed consent was first obtained from all individual

participants included in the study. Using scale and meter, members of the research team subsequently measured participants' weight and height before they answered the baseline survey. Then, all participants received the informational intervention. Support condition participants additionally received instructions on the social support intervention (details see below). Participants were further instructed to keep a 13-day diary, starting the next day. One month and two months after the diary phase, in December 2014 and January 2015 respectively, participants were asked to fill in the two online follow-up surveys. After the last follow-up survey, participants were fully debriefed (including information on the experimental confederate, see below), and entered a lottery to win a \$1000.- prize (helicopter flight or wellness weekend), or one of eight shopping vouchers. Students were offered the choice of receiving study credit for their participation instead of entering the lottery.

Intervention

Information on healthy eating and the social support intervention were delivered individually by trained psychology master's students. Participants in the control condition only received the information on healthy eating, whereas participants in the support condition additionally partook in the social support intervention. All intervention materials and protocols can be found in the Electronic Supplementary Material.

Information on healthy eating. Information on healthy eating was presented at participants' individual lab appointments, after baseline assessment. It included behavior change techniques (BCTs) 5.1 "Information about health consequences", and 1.1 "Goal setting" (Michie et al., 2013). The information about healthy eating that was tailored to the participant's randomly assigned eating goal (increasing fruit and vegetable consumption or decreasing unhealthy snack consumption). After the presentation, participants received a fact sheet with identical information as presented by the interventionist: a definition of [fruit and vegetable consumption / unhealthy snacks], health effects of [fruit and vegetable consumption / unhealthy snack consumption], and current recommendations for consumption. Also, the

interventionist reinforced participants' assigned eating goal by saying: "Therefore, it is very important that you [eat more fruits and vegetables / avoid unhealthy snacks]". This was also printed on the factsheet.

Social support intervention. The social support intervention included BCT 3.1 "Social support (unspecified)" (Michie et al., 2013). After receiving the informational intervention, social support participants were informed that they would be invited by the group administrator (a male Master's student of the research team) to join a WhatsApp chat group starting Day 4 of the diary for seven days. Participants who did not have WhatsApp were assisted in installing the app. Participants were asked to support each other to reach their assigned eating goal. They were assured that their identity and exchanged content would be kept confidential.

We randomly assigned social support participants to small chat groups, plus one female confederate moderator ($n=32$ groups with $Md=3$ participants; $Min=2$, and $Max=5$ participants). The purpose of the moderator was to ensure that a minimum of social support was delivered to the chat group. She ostensibly participated in the group like a regular participant, but was trained to provide a standardized support message on each of the seven intervention days (e.g. "Hey guys, how's it going for you on the second day? Wishing you lots of success with consuming fewer unhealthy snacks! And just reach out if things get rough. I will do the same :)"). Furthermore, moderators were instructed to respond with a supportive message to any message posted. If suitable, moderators followed a list of supportive responses to common support demands; otherwise, they wrote unstandardized supportive responses.

On the evening of the seventh day of the intervention, participants were informed that the chat group would be deleted by the administrator the next morning. At 8 AM on the eighth day, the administrator downloaded the chat content and then deleted the chat.

Measures

Primary outcome: Healthy eating.

Self-reported healthy eating was assessed using an electronic daily evening diary. Participants were asked, “How many servings of fruits and vegetables did you eat today?”, and “How many unhealthy snacks did you eat today?”. Following the conceptualization by Kelly, Smith, King, Flood, & Bauman (2007), an unhealthy snack is any food consumed between main meals that belongs to the non-core categories (e.g. cake, candy, fast food). The servings of fruits and vegetables represented the outcome for the fruit and vegetable goal group, whereas the servings of unhealthy snacks served as the outcome for the unhealthy snack goal group. Because the reports were based on a single question each day, reliability on each day could not be estimated, but we assessed the consistency of responses of each item over the 10 days. Cronbach’s alpha for fruit/vegetable consumption was .92 and for unhealthy snacks it was .78, indicating systematic responses. Because we made identical predictions for each outcome in its respective goal group, we constructed a single harmonized index as the pre-registered outcome variable. First, for the fruit and vegetable goal group, the group’s mean fruit and vegetable consumption was subtracted from each participant’s fruit and vegetable servings on a particular day, and divided by the group’s standard deviation of fruit and vegetable consumption. Second, for the snack goal group, the group’s mean snack consumption was subtracted from each participant’s snacks on a particular day, divided by the group’s standard deviation of snack consumption, and reverse coded. In both goal groups, the relevant eating outcome therefore had a mean of zero and a standard deviation of one, which harmonized them for a combined analysis while eliminating any differences between goal groups. Across both goal groups, positive values indicate healthier eating than the typical person, and negative values indicate unhealthier eating than the typical person.

To investigate the maintenance of behavior change, self-reported daily healthy eating was also assessed at 1-month and 2-month follow-up. The same items were used as in the

diary, the only difference being that they referred to “yesterday” rather than “today” in order to assure a full daily consumption was reported.

Covariates.

At study registration, several covariates were pre-specified. Besides gender and age, being vegetarians, vegans, or having diabetes, BMI was calculated from participants’ objectively measured weight and height, and active participation in the chat groups was coded (0=no message sent, 1=at least one message sent). Further assessed were restrained eating (Dutch Eating Behavior Questionnaire; Grunert, 1989; van Strien, Frijters, Jan E. R., Bergers, Gerard P. A., & Defares, 1986, $\alpha=0.88$), stress (Kuhl & Fuhrmann, 1998, $\alpha=0.87$), and social desirability (Balanced Inventory of Desirable Responding; Paulhus, 1991; Winkler, Kroh, & Spiess, 2006, $\alpha=0.61$).

Data Analysis

Analysis of intervention effects on healthy eating. As specified in the study registration, we conducted independent samples t-tests to test for differences in fruit and vegetable consumption, unhealthy snack consumption, and healthy eating between the social support and the control condition during the last three days of the intervention. We computed the average of each outcome over Day 8 to 10 for each participant: average scores for fruit and vegetable consumption (for the fruits and vegetable goal group), unhealthy snack consumption (for the snack goal group), and healthy eating (for the total sample).

Analysis of temporal dynamics of intervention effects. To investigate the day-to-day intervention effects, we constructed a segmented linear mixed model (i.e., a spline model, Fitzmaurice, Laird, & Ware, 2012). This allowed for separate time slopes for the three days prior to the intervention and for the intervention period (see also Figure S-2 in the Electronic Supplementary Material):

$$Y_{igt} = b_0 + b_1I_g + b_2T_t + b_3I_gT_t + b_4I_gpreT_t + u_g + u_{i(g)} + w_{i(g)}T_t + e_{igt} \quad \text{Eq. 1}$$

Y_{igt} is the health behavioral outcome of person i in chat group g at day t . The interpretation of the regression coefficients (b) are determined from the scoring of the independent variables, I , T , and $preT$. I is a dummy code for intervention condition (0: control; 1: social support); T_t is day in the study, centered on Day 10 (last intervention day); $preT_t$ is a time variable that is coded -2, -1, and 0 for the first three days and zero for the remaining seven days. With this coding and the model represented in Eq. 1, b_0 reflects average healthy eating for the control condition at Day 10. b_1 represents the intervention effect at Day 10, i.e., the difference in healthy eating between the control condition and the social support condition. b_2 is the time trend (slope) over the diary period in the control condition. Furthermore, b_3 represents the difference in the slope for healthy eating between the support condition and control condition from one day to the next during the intervention period. b_4 represents the possibility of a difference between the support and the control condition during the pre-intervention time.

The model accounted with random effects for the fact that the daily observations were nested in individuals and chat groups, and allowed for the possibility that the intervention effect might vary across the 32 chat groups (u_g), and across individuals (at Day 10: $u_{i(g)}$; over time: $w_{i(g)}T_t$). We also estimated the residual variance (e_{igt}), and fit an AR(1) covariance structure of the repeated measure residuals. The models were fitted using the PROC MIXED procedure in SAS. The code is provided in the Electronic Supplementary Material.

We further conducted several additional analyses. First, we investigated the possibility of a discrete change in healthy eating at the onset of the intervention. For this purpose, we added an interaction term of a dummy coded variable for intervention phase (-1: pre intervention; 0: intervention phase) with the intervention condition. Furthermore, we conducted sensitivity analyses. Models were computed again adding the pre-registered

covariates¹. Also, in line with recommendations (White, Horton, Carpenter, & Pocock, 2011), we conducted a series of missing analyses, to investigate whether the intervention effects depended on dropout. Our main model used all available data, and treated missing values as missing at random. A second model replaced missing values of participants who dropped out during the diary with their first diary day (baseline value carried forward). This can be considered the most conservative model as null intervention effects are assumed for dropouts. Similarly, a third model used the third diary day to replace missing values, as first diary day was missing for some persons. Finally, a fourth model replaced missing values of dropouts with the last observed value (last observation carried forward).

Analysis of maintenance of intervention effects. To investigate the maintenance of intervention effects on eating outcomes, we conducted independent samples t-tests for the immediate follow-up (averaged Day 11 to 13 per person) as well as the 1-month and 2-month follow-ups.

Results

Preliminary Analyses

Of the 232 participants, 203 participants (87.5%) had at least one diary entry and were included in the main analyses. See Figure 1 for sample sizes of the four experimental conditions (support vs. control condition by assigned eating goal). The analyzed participants had filled in on average 10.0 out of 13 diary days ($SD=4.3$). They were on average 27.5 years old ($SD=8.6$), and had a mean BMI of 23.5 ($SD=4.0$). Most participants were female (75.5%), 24.5% were male, and enrolled as students (58.7%; 41.3% staff members and other adults). The 29 persons who were randomized but did not fill out a single diary did not significantly differ from compliers in age, gender, and student/work status. There were also no significant baseline differences between the social support condition and the control condition (see Table S-4 in the Electronic Supplementary Material).

¹ Diabetes was not included, because none of the participants reported having diabetes.

In total, 1249 messages were exchanged during the 7-day intervention period. Of these, 666 messages were sent by the participants (53.3%), whereas 478 messages were written by the confederate (38.3%) and 105 were written by the group administrator (8.4 %). Per group, 14 to 95 messages were exchanged ($M= 41.4$, $SD= 4.3$). On average, participants wrote 7.1 messages ($SD = 0.7$, range: 0-29). There were 10 silent participants who attended the group, but never wrote a message, and nine participants left the group after invitation.

Regarding intervention fidelity, 95% of the standardized support messages to be sent by the confederate were actually posted into the chat groups, whereas 5% were not sent. Eighty percent of the standardized messages were sent at the pre-specified time, and 20% of the messages were sent between 13 to 108 minutes later than scheduled. Sixty-one percent of the sent messages were correct in terms of contents, whereas 39% of the sent messages were modified in minor ways, e.g. to fit it more naturally into an ongoing conversation or to adapt it to the audience (e.g., also addressing workers instead of students only).

Immediate Intervention Effects on Healthy Eating

At the last three days of the intervention, social support participants ate almost one and a half fruits and vegetables more ($M_{diff}=1.31$; 95% CI [0.17, 2.45], $t[76]=2.28$, $p=0.025$), or about two thirds of an unhealthy snack less than controls ($M_{diff}=-0.59$; 95% CI [-1.11, -0.06], $t[79]=2.23$, $p=.028$). The combined healthy eating outcome indicated that social support participants ate two fifths of a standard deviation more healthily than controls ($M_{diff}=0.39$; 95% CI [0.15, 0.62], $t[157]=3.27$, $p=.001$).

Temporal Dynamics of Intervention Effects

>>>>> ADD FIGURE 2 HERE <<<<<<

The temporal model displayed in Figure 2 and Table 1 provides a detailed account of the primary results (see also Figure S-2 in the Electronic Supplementary Material). There was a significant gradual increase in healthy eating over the course of the intervention. On Day 10, social support participants in the fruit and vegetable goal subgroup ate more than five fruits

and vegetables – almost one and a half fruits and vegetables more than controls ($B=1.43$, $SE=0.55$, $p=.020$). In the snack goal group, they ate four fifths of an unhealthy snack less than controls ($B=-0.80$, $SE=0.27$, $p=.010$). The analysis for the total sample indicated that social support participants ate about half of a standard deviation more healthily than controls on Day 10 ($B=0.48$, $SE=0.12$, $p<.001$).

Social support participants increased their fruit and vegetable consumption by almost a sixth of a serving on each day of the intervention ($B=0.15$, $SE=0.07$, $p=.031$). In the unhealthy snack goal group, they reduced their unhealthy snack consumption by about a seventh more than controls from day to day ($B=-0.14$, $SE=0.05$, $p=.007$). Correspondingly, the combined healthy eating outcome in the total sample indicated a small day to day increase in healthy eating compared to controls ($B=0.06$, $SE=0.02$, $p<.001$). Control group participants seemed to decrease in terms of healthy eating over the study period, although this was not significant ($B=-0.01$, $SE=0.01$, $p=.162$).

>>>>> ADD TABLE 1 HERE <<<<<

To rule out the possibility that eating already started changing in the social support condition before the beginning of the chat groups, we further compared the intervention and control conditions during the 3-day pre-intervention phase. We found no indication of pre-intervention differences in fruit and vegetable consumption for the fruit and vegetable goal group ($B=-0.27$, $SE=0.24$, $p=.259$), unhealthy snack consumption for the snack goal group ($B=0.24$, $SE=0.15$, $p=.120$), or healthy eating in the total sample respectively ($B=-0.09$, $SE=0.05$, $p=.100$).

The random effect estimate for the chat groups shows that fruit and vegetable consumption on Day 10 varied significantly across chat groups in the intervention condition (see Table 1, lower half). Expressed in standard deviations, the results indicate that fruit and vegetable consumption on Day 10 varied by $\sqrt{0.77} = 0.88$ between the chat groups.

Assuming normal distribution, this means that 95% of social support participants consumed between 3.5 to 7.0 fruits and vegetables on the last day of the intervention.

Additionally, there was variation in the outcomes at Level 2, meaning that participants significantly differed in their fruit and vegetable consumption, and unhealthy snack consumption at the last day of the intervention. Level 1 results indicated the presence of autocorrelation, which attained significance for the healthy eating outcome. Finally, there was a significant proportion of residual variance in healthy eating, fruit and vegetable, and unhealthy snack consumption that was unaccounted for by the models.

The additional analyses indicated no instantaneous effects on healthy eating at the beginning of the intervention. Also, the inclusion of the covariates did not substantively change the model results. Finally, all missing analyses confirmed the intervention effects. See Table S-5 in the Electronic Supplementary Material.

>>>>> ADD FIGURE 3 HERE <<<<<<

Maintenance of Intervention Effects

To determine whether the intervention effects held in the days after the intervention and 1-month and 2-month follow-ups, we compared the effects at each of those time points. The results are displayed in Figure 3. The intervention effect remained significant immediately after the intervention for unhealthy snack consumption ($M_{diff}=-0.68$; 95% CI [-1.32, -0.05], $t=-2.14$, $df=53.160$, $p=.043$, $d=0.59$), and healthy eating in the total sample ($M_{diff}=0.34$; 95% CI [0.07, 0.62], $t=2.46$, $df=150$, $p=.015$, $d=0.40$), but not for fruit and vegetable consumption ($M_{diff}=0.58$; 95% CI [-0.40, 1.55], $t=1.18$, $df=72$, $p=.242$, $d=0.28$). There was no indication of the intervention effect at 1-month follow-up for fruit and vegetable consumption ($M_{diff}=0.32$; 95% CI [-.47, 1.10], $t=-0.81$, $df=71$, $p=.423$, $d=0.19$), unhealthy snack consumption ($M_{diff}=0.08$; 95% CI [-0.73, 0.89], $t=0.21$, $df=68.377$, $p=.838$, $d=0.05$), or for healthy eating in the total sample ($M_{diff}=0.07$; 95% CI [-0.26, 0.40], $t=0.43$, $df=143$, $p=.671$, $d=0.07$). Neither was there an intervention effect at 2-month follow-up for fruit and

vegetable consumption ($M_{\text{diff}}=0.52$; 95% CI [-0.46, 1.49], $t=1.05$, $df=69$, $p=.296$, $d=0.25$), unhealthy snack consumption ($M_{\text{diff}}=-0.23$; 95% CI [-0.88, 0.43], $t=-0.68$, $df=57.856$, $p=.501$, $d=0.18$), or for healthy eating in the total sample ($M_{\text{diff}}=0.21$; 95% CI [-0.12, 0.54], $t=1.24$, $df=136.237$, $p=.0217$, $d=0.21$). Adding gender, age, BMI, restrained eating, stress, and social desirability did not substantively change the results.

Discussion

The results provide first experimental evidence for the hypothesis that smartphone-based support groups can promote healthy eating in daily life. The intervention successfully increased fruit and vegetable consumption, and decreased unhealthy snack consumption, suggesting generalizability to both eating behaviors. By the last day of the intervention, social support participants ate almost one and a half more servings of fruits and vegetables or almost one unhealthy snack less than control participants did. These are sizeable effects considering the short intervention period of seven days.

The intensive longitudinal assessment allowed for interesting new insights into the temporal effects of the intervention. The data supported a gradual increase in healthy eating due to the intervention rather than an immediate discrete increase after the support groups were formed. Likely, participants of chat groups first have to be acquainted and possibly gain trust before social support processes begin to take place.

There was variance in healthy eating between the different chat groups at the end of the intervention, which was significant for the fruit and vegetable goal group. This suggests that in some groups, the social support processes worked better than in others. Future research might investigate factors that contribute to the success of social support groups, such as group climate (Bakali, Wilberg, Klungsøyr, & Lorentzen, 2013), or normative processes (Robinson, Fleming, & Higgs, 2014).

Contrary to expectations, the intervention effects, for the most part, did not last beyond the end of the social support groups. The effects would have persisted if the participants had

created healthy eating habits that became more automatic or had enabled self-efficacy (Rackow, Scholz, & Hornung, 2015), but our 7-day intervention period appeared to have been too short to foster such effects. As such, previous studies have used longer intervention periods of typically four weeks or longer (Prestwich et al., 2014; Wing & Jeffery, 1999). Another possible explanation for the lack of persistent effects is the timing of our intervention. The intervention period took place during the month of November and early December, and the 1-month follow-up survey was conducted around Christmas and New Year's Eve. This is a period when participants may have found it particularly difficult to adhere to their eating goals. Hence, the effects may have been lost. The negative tendencies towards unhealthier eating in the control condition lends some support to this explanation. Future studies that replicate and extend this intervention at a more favorable time for changing eating may therefore expect larger effects and greater maintenance.

Strengths and Limitations

The present study has several strengths. First, the study provides first experimental evidence that smartphone-based peer support groups can promote healthy eating. We successfully created a scenario where persons sharing the same eating goal were able to share their experiences related to achieving their goals. This finding encourages the practical application of the intervention for eating regulation, by training an expert to provide emotional, informational, and instrumental social support to group members. From our results, this supportive effect is expected while the support is ongoing. Further research is needed to investigate how the effects can be sustained after the chat groups end. Alternatively, a possible low-cost intervention could be to encourage chat group participants to continue the peer support after the trained expert exits the chat group. Second, the experimental and intensive longitudinal study design allowed for rare new insights into the temporal development of intervention effects. Even when researchers collect intensive longitudinal data in randomized trials, the opportunity to investigate temporal intervention effects is often

missed, and the data are analyzed on aggregate level only (e.g. Brookie et al., 2017). Third, the intensive longitudinal design allowed for assessing the heterogeneity of the intervention effects at the individual and group level, rather than the average effects of the intervention alone. For example, our analysis revealed that not all individuals equally benefitted from the intervention, which can inspire further research on moderators of the effects of support groups. A fourth strength is the investigation of the intervention for two different eating goals.

This randomized trial also had some limitations that offer opportunities for future research. First, the primary outcome measure of the study was self-reported. Retrospective bias was minimized by daily diary, but social desirability effects may have biased the reports. The sensitivity analysis adjusting for social desirability lends confidence to the conclusion that the differences between the control condition and the social support condition were not due to social desirability. Furthermore, studies using objective biomarkers of eating have indicated good validity of self-reports (Brookie et al., 2017). Second, the effects were shown with a sample of participants who had a healthy baseline diet but who reported having difficulty in maintaining the diet. The intervention might not necessarily work with participants who had weak motivation to eat healthily, but we predict that it would generalize to motivated persons with lower fruit and vegetable consumption or higher snack consumption at baseline. Studies in other populations, including those who are overweight or obese, are needed to assess the generalizability of the findings. Third, our effects were obtained by treating the trained moderator as a confederate peer group member. The presence of an unidentified peer moderator rather than an identified moderator labeled as coach or staff member or no moderator may have altered the effects of this support group intervention. The effects might be larger due to having a trained professional in the group, but they might also have been diminished due to possible reactance of participants suspecting deception. This may be an additional reason for the between-person variability in intervention effects. We included the covert expert in this study, because we wanted to prompt peer support in this

study, and ensure minimal support in each group. Future research is needed to address the question if moderator presence and type in support groups contributes to their effect.

Conclusions

This study extends literature on the benefits of social support for health behavior change to smartphone-based support groups. The results are encouraging as the integration of social support groups via personal smartphones provides a low-cost way to promote healthy eating among persons motivated to change their eating behavior. The maintenance of behavior change remains a challenge that will need to be addressed in the future.

References

- Anderson Steeves, E., Jones-Smith, J., Hopkins, L., & Gittelsohn, J. (2016). Perceived social support from friends and parents for eating behavior and diet quality among low-income, urban, minority youth. *Journal of Nutrition Education and Behavior*, *48*(5), 304-310.e1. <https://doi.org/10.1016/j.jneb.2015.12.014>
- Atienza, A. A., King, A. C., Oliveira, B. M., Ahn, D. K., & Gardner, C. D. (2008). Using hand-held computer technologies to improve dietary intake. *American Journal of Preventive Medicine*, *34*(6), 514–518. <https://doi.org/10.1016/j.amepre.2008.01.034>
- Bakali, J. V., Wilberg, T., Klungøy, O., & Lorentzen, S. (2013). Development of group climate in short- and long-term psychodynamic group psychotherapy. *International Journal of Group Psychotherapy*, *63*(3), 366–393. <https://doi.org/10.1521/ijgp.2013.63.3.366>
- Ballantine, P. W., & Stephenson, R. J. (2011). Help me, I'm fat! Social support in online weight loss networks. *Journal of Consumer Behaviour*, *10*(6), 332–337. <https://doi.org/10.1002/cb.374>

- Bardus, M., Smith, J. R., Samaha, L., & Abraham, C. (2015). Mobile phone and Web 2.0 technologies for weight management: A systematic scoping review. *Journal of Medical Internet Research, 17*(11), e259. <https://doi.org/10.2196/jmir.5129>
- Brookie, K. L., Mainvil, L. A., Carr, A. C., Vissers, M. C., & Conner, T. S. (2017). The development and effectiveness of an ecological momentary intervention to increase daily fruit and vegetable consumption in low-consuming young adults. *Appetite, 108*, 32–41. <https://doi.org/10.1016/j.appet.2016.09.015>
- Brunzell, J. D. (2007). Hypertriglyceridemia. *New England Journal of Medicine, 357*(10), 1009–1017. <https://doi.org/10.1056/NEJMcp070061>
- Faul, F., Erdfelder, E., Lang, A.-G., & Buchner, A. (2007). G*power 3: A flexible statistical power analysis program for the social, behavioral, and biomedical sciences. *Behavior Research Methods, 39*(2), 175–191.
- Fitzmaurice, G. M., Laird, N. M., & Ware, J. H. (2012). *Applied longitudinal analysis* (2nd Edition). Hoboken, NJ: John Wiley & Sons.
- Fitzsimons, G. M., & Finkel, E. J. (2010). Interpersonal influences on self-regulation. *Current Directions in Psychological Science, 19*(2), 101–105. <https://doi.org/10.1177/0963721410364499>
- Graaf, C. de. (2006). Effects of snacks on energy intake: An evolutionary perspective. *Appetite, 47*(1), 18–23. <https://doi.org/10.1016/j.appet.2006.02.007>
- Grunert, S. C. (1989). Ein Inventar zur Erfassung von Selbstaussagen zum Ernährungsverhalten. *Diagnostica, 35*(2), 167–179.
- Helgeson, V. S., & Gottlieb, B. H. (2000). Support groups. In S. Cohen, L. G. Underwood, & B. H. Gottlieb (Eds.), *Social support measurement and intervention: A guide for health and social scientists* (pp. 221–245). New York: Oxford University Press.

- Heron, K. E., & Smyth, J. M. (2010). Ecological momentary interventions: Incorporating mobile technology into psychosocial and health behaviour treatments. *British Journal of Health Psychology, 15*, 1–39. <https://doi.org/10.1348/135910709X466063>
- Inauen, J., ShROUT, P. E., Bolger, N., Stadler, G., & Scholz, U. (2016). Mind the gap? An intensive longitudinal study of between-person and within-person intention-behavior relations. *Annals of Behavioral Medicine, 1*–7. <https://doi.org/10.1007/s12160-016-9776-x>
- Joo, N.-S., & Kim, B.-T. (2007). Mobile phone short message service messaging for behaviour modification in a community-based weight control programme in Korea. *Journal of telemedicine and telecare, 13*(8), 416–420. <https://doi.org/10.1258/135763307783064331>
- Kelly, B., Smith, B., King, L., Flood, V., & Bauman, A. (2007). Television food advertising to children: The extent and nature of exposure. *Public Health Nutrition, 10*(11), 1234–1240. <https://doi.org/10.1017/S1368980007687126>
- Kiernan, M., Moore, S. D., Schoffman, D. E., Lee, K., King, A. C., Taylor, C. B., . . . Perri, M. G. (2012). Social support for healthy behaviors: Scale psychometrics and prediction of weight loss among women in a behavioral program. *Obesity (Silver Spring, Md.), 20*(4), 756–764. <https://doi.org/10.1038/oby.2011.293>
- Kuhl, J., & Fuhrmann, A. (1998). Decomposing self-regulation and self-control: The Volitional Components Inventory. In J. Heckhausen (Ed.), *Motivation and self-regulation across the life span* (pp. 15–49). New York, NY: Cambridge University Press.
- Kwan, M. Y., & Gordon, K. H. (2016). The effects of social support and stress perception on bulimic behaviors and unhealthy food consumption. *Eating Behaviors, 22*, 34–39. <https://doi.org/10.1016/j.eatbeh.2016.03.024>

- Kwasnicka, D., Dombrowski, S. U., White, M., & Sniehotta, F. (2016). Theoretical explanations for maintenance of behaviour change: a systematic review of behaviour theories. *Health psychology review*, 1–20. <https://doi.org/10.1080/17437199.2016.1151372>
- McKinley, C. J. (2009). Investigating the influence of threat appraisals and social support on healthy eating behavior and drive for thinness. *Health Communication*, 24(8), 735–745. <https://doi.org/10.1080/10410230903264303>
- McKinley, C. J., & Wright, P. J. (2014). Informational social support and online health information seeking: Examining the association between factors contributing to healthy eating behavior. *Computers in Human Behavior*, 37, 107–116. <https://doi.org/10.1016/j.chb.2014.04.023>
- Michie, S., Richardson, M., Johnston, M., Abraham, C., Francis, J., Hardeman, W., . . . Wood, C. E. (2013). The behavior change technique taxonomy (v1) of 93 hierarchically clustered techniques: Building an international consensus for the reporting of behavior change interventions. *Annals of Behavioral Medicine*, 46(1), 81–95. <https://doi.org/10.1007/s12160-013-9486-6>
- Mohr, D. C., Burns, M. N., Schueller, S. M., Clarke, G., & Klinkman, M. (2013). Behavioral intervention technologies: Evidence review and recommendations for future research in mental health. *General Hospital Psychiatry*, 35(4), 332–338. <https://doi.org/10.1016/j.genhosppsych.2013.03.008>
- Murakami, K., & Livingstone, M. B. (2016). Associations between meal and snack frequency and overweight and abdominal obesity in US children and adolescents from National Health and Nutrition Examination Survey (NHANES) 2003-2012. *British Journal of Nutrition*, 115(10), 1819–1829. <https://doi.org/10.1017/S0007114516000854>

- Papies, E. K., & Hamstra, P. (2010). Goal priming and eating behavior: Enhancing self-regulation by environmental cues. *Health Psychology, 29*(4), 384–388.
<https://doi.org/10.1037/a0019877>
- Paulhus, D. L. (1991). Measurement and control of response bias. In J. P. Robinson, P. R. Shaver, & L. S. Wrightsman (Eds.), *Measures of personality and social psychological attitudes* (pp. 17–59). San Diego, CA: Academic Press.
- Prestwich, A., Conner, M. T., Lawton, R. J., Ward, J. K., Ayres, K., & McEachan, Rosemary R. C. (2014). Partner- and planning-based interventions to reduce fat consumption: Randomized controlled trial. *British Journal of Health Psychology, 19*(1), 132–148.
<https://doi.org/10.1111/bjhp.12047>
- Rackow, P., Scholz, U., & Hornung, R. (2015). Received social support and exercising: An intervention study to test the enabling hypothesis. *British Journal of Health Psychology, 20*(4), 763–776. <https://doi.org/10.1111/bjhp.12139>
- Rains, S. A., Peterson, E. B., & Wright, K. B. (2015). Communicating social support in computer-mediated contexts: A meta-analytic review of content analyses examining support messages shared online among individuals coping with illness. *Communication Monographs, 82*(4), 403–430. <https://doi.org/10.1080/03637751.2015.1019530>
- Robinson, E., Fleming, A., & Higgs, S. (2014). Prompting healthier eating: testing the use of health and social norm based messages. *Health psychology : official journal of the Division of Health Psychology, American Psychological Association, 33*(9), 1057–1064.
<https://doi.org/10.1037/a0034213>
- Scholz, U., Ochsner, S., Hornung, R., & Knoll, N. (2013). Does social support really help to eat a low-fat diet? Main effects and gender differences of received social support within the Health Action Process Approach. *Applied Psychology: Health and Well-Being, 5*(2), 270–290. <https://doi.org/10.1111/aphw.12010>

- Turati, F., Rossi, M., Pelucchi, C., Levi, F., & La Vecchia, C. (2015). Fruit and vegetables and cancer risk: A review of southern European studies. *British Journal of Nutrition*, *113*(SupplementS2), S102-S110. <https://doi.org/10.1017/S0007114515000148>
- U.S. Department of Health and Human Services and U.S. Department of Agriculture. (2015). 2015–2020 dietary guidelines for Americans (8th Edition). Retrieved from <http://health.gov/dietaryguidelines/2015/guidelines>
- Uchino, B. N., Bowen, K., Carlisle, M., & Birmingham, W. (2012). Psychological pathways linking social support to health outcomes: A visit with the “ghosts” of research past, present, and future. *Social Science & Medicine*, *74*(7), 949–957. <https://doi.org/10.1016/j.socscimed.2011.11.023>
- van Strien, T., Frijters, Jan E. R., Bergers, Gerard P. A., & Defares, P. B. (1986). The Dutch Eating Behavior Questionnaire (DEBQ) for assessment of restrained, emotional, and external eating behavior. *International Journal of Eating Disorders*, *5*(2), 295–315. [https://doi.org/10.1002/1098-108X\(198602\)5:2<295::AID-EAT2260050209>3.0.CO;2-T](https://doi.org/10.1002/1098-108X(198602)5:2<295::AID-EAT2260050209>3.0.CO;2-T)
- Wang, X., Ouyang, Y., Liu, J., Zhu, M., Zhao, G., Bao, W., & Hu, F. B. (2014). Fruit and vegetable consumption and mortality from all causes, cardiovascular disease, and cancer: Systematic review and dose-response meta-analysis of prospective cohort studies. *BMJ*, *349*, g4490. <https://doi.org/10.1136/bmj.g4490>
- White, I. R., Horton, N. J., Carpenter, J., & Pocock, S. J. (2011). Strategy for intention to treat analysis in randomised trials with missing outcome data. *BMJ*, *342*, d40. <https://doi.org/10.1136/bmj.d40>
- White, M., & Dorman, S. M. (2001). Receiving social support online: Implications for health education. *Health Education Research*, *16*(6), 693–707. <https://doi.org/10.1093/her/16.6.693>

- Williams, G., Hamm, M. P., Shulhan, J., Vandermeer, B., & Hartling, L. (2014). Social media interventions for diet and exercise behaviours: A systematic review and meta-analysis of randomised controlled trials. *BMJ Open*, *4*(2), e003926. <https://doi.org/10.1136/bmjopen-2013-003926>
- Wing, R. R., & Jeffery, R. W. (1999). Benefits of recruiting participants with friends and increasing social support for weight loss and maintenance. *Journal of Consulting and Clinical Psychology*, *67*(1), 132–138. <https://doi.org/10.1037/0022-006X.67.1.132>
- Winkler, N., Kroh, M., & Spiess, M. (2006). Entwicklung einer deutschen Kurzskala zur zweidimensionalen Messung von sozialer Erwünschtheit. *Discussion Papers of DIW Berlin*. (579), 1–31. Retrieved from http://www.diw.de/documents/publikationen/73/diw_01.c.44281.de/dp579.pdf

Table 1. *Linear Mixed Model Predicting Intervention Effects on Healthy Eating.*

	Fruits and Vegetables (servings per day) F&V Goal Group (<i>n</i> = 97)		Unhealthy Snacks (servings per day) Snack Goal Group (<i>n</i> = 106)		Healthy Eating (z standardized) Total Sample (<i>N</i> = 203)	
	Estimate (<i>SE</i>)	95% CI	Estimate (<i>SE</i>)	95% CI	Estimate (<i>SE</i>)	95% CI
Fixed effects (intercept, slopes)						
Intercept (mean control condition Day 10)	3.85 (0.34)***	[3.13, 4.57]	1.61 (0.18)***	[1.22, 2.00]	-0.17 (0.08)*	[-0.34, <-0.01]
Intervention effect Day 10	1.43 (0.55)*	[0.26, 2.59]	-0.80 (0.27)**	[-1.38, -0.23]	0.48 (0.12)***	[0.23, 0.73]
Time (slope control condition)	-0.02 (0.04)	[-0.10, 0.06]	<0.01 (0.03)	[-0.05, 0.07]	-0.01 (0.01)	[-0.04, 0.01]
Intervention by time	0.15 (0.07)*	[0.01, 0.29]	-0.14 (0.05)**	[-0.24, -0.04]	0.06 (0.02)***	[0.03, 0.10]
Intervention by pre-intervention time	-0.27 (0.24)	[-0.74, 0.20]	0.24 (0.15)	[-0.06, 0.55]	-0.09 (0.05)†	[-0.20, 0.02]
Random effects ([co-]variances)						
Level 3 (chat groups): Intercept ^a	0.77*	[0.25, 10.39]	0.07	[0.01, 25514]	0.04	[0.01, 1.79]
Level 2 (between-person)						
Intercept ^a	2.79***	[1.76, 5.09]	0.38***	[0.14, 2.87]	0.32***	[0.21, 0.52]
Time	0.11**	[0.02, 0.21]	-0.02	[-0.09, 0.06]	0.01	[-0.01, 0.02]
Intercept-time covariance	^b	^b	0.01†	[<0.01, 0.11]	<0.01	[<0.01, 0.03]
Level 1 (within-person)						
Autocorrelation	0.03	[-0.05, 0.11]	0.01	[-0.09, 0.11]	0.12**	[0.05, 0.19]
Residual	5.00***	[4.50, 5.59]	2.44***	[2.88, 2.76]	0.53***	[0.48, 0.58]

Note. Time: Linear trend from pre-intervention Day 1 (= -10) to the last day of the intervention (= 0), Pre-intervention time = 3 days prior to intervention. Intervention: 1 = social support, 0 = control. F&V = fruit and vegetable.

^a Estimate of variance of healthy eating at the last day of the intervention.


^b Was not estimable.



* $p < .05$, ** $p < .01$, *** $p < .001$, † $p < .10$.

Figure 1. Participant flow through the trial. BMI = Body Mass Index, F&V = Fruit and Vegetable.

Figure 2. Model predicted intervention effects on self-reported healthy eating over time.
F&V: Fruit and Vegetable.

Figure 3. Intervention effects on healthy eating during the intervention (averaged diary days 4-10), immediately post intervention (averaged diary days 11-13), and at 1-month and 2-month follow-up. Solid lines represent the social support condition, dashed lines represent the control condition. Error bars: +/- 2 times the standard error. F&V: Fruit and Vegetable.





Using Smartphone-Based Support Groups to Promote Healthy Eating in Daily Life: A Randomized Trial

Jennifer Inauen, Niall Bolger, Patrick E. Shrout, Gertraud Stadler, Melanie
Amrein, Pamela Rackow, & Urte Scholz

Electronic Supplementary Material

Contents:

1. Intervention Materials (English Translations)
2. Graphic Representation of the Statistical Model
3. SAS Code
4. Results: Supplemental Tables and Figures


1. Intervention Materials (English Translations)

Information on healthy eating: Factsheets

a) Factsheet for the fruit and vegetable goal subgroup

Information about healthy eating

What do we understand by healthy eating?
Healthy eating describes a diversified, balanced diet. It is recommended to eat more plant foods than animal foods, and to eat only small amounts of fat, salt, and sugar.
It is therefore very important that you eat more fruits and vegetables.



Why is healthy eating so important?
Eating is a crucial element to maintain our physical health.

The consumption of fruits and vegetables has positive health effects.


Fruits and vegetables
Fruits and vegetables are the major source for vitamins, minerals, and fibers. They are low in calories and low in fat, and they are easy to digest. A varied diet with lots of fruits and vegetables is therefore beneficial to health and well-being.

Fruits include, for example, apples, bananas, berries, apricots, and pineapple. Vegetables include, for example, tomatoes, spinach, or broccoli. Potatoes, corn and legumes are not considered vegetables. They are considered carbohydrates because of their high starch content.

What are the consequences of fruit and vegetable consumption?
High fruit and vegetable consumption lowers the risk for cardiovascular disease, Type II diabetes, some forms of cancer, rheumatism and osteoporosis.

Did you know that ...?


- only 30% of the Swiss population adhere to the recommended consumption of five fruits and vegetables per day.
- the most popular vegetable in Switzerland is the carrot. The least favorite is Brussels sprouts.
- the degree of self-sufficiency for vegetables is 55 % in Switzerland. The rest is imported from France, Italy, and Spain.



b) Factsheet for the unhealthy snack goal subgroup

Information about healthy eating

What do we understand by healthy eating?
Healthy eating describes a diversified, balanced diet. It is recommended to eat more plant foods than animal foods, and to eat only small amounts of fat, salt, and sugar.
It is therefore very important that you avoid unhealthy snacks.



Why is healthy eating so important?
Eating is a crucial element to maintain our physical health.
The consumption of unhealthy snacks has negative health effects.

Unhealthy snacks
Snacks are any foods consumed in between the three main meals (breakfast, lunch, dinner). Dessert is also included in lunch or dinner. Snacks can be healthy or unhealthy.

Unhealthy snacks are:

- Confectionary
- Fast food restaurant meals
- Cakes, biscuits and muesli bars
- Savory crisps and pastries
- Gravies and sauces
- Sugared drinks and fruit juice
- Ice cream
- High-sugar/high-fat spreads
- fried potato products

What are the consequences of excessive unhealthy snack consumption?
Excessive unhealthy snack consumption can increase the risk of Type II diabetes, cardiovascular disease, high cholesterol, and high blood pressure. It can also lead to weight gain and decreased fat burning.

Did you know that ...?

- the World Health Organization WHO recommends to keep sugar intake at maximum 10% of the total energy intake. In Switzerland in 2005, sugar consumption represented 17% of the total energy intake.
- consumers in Europe, on average, snack 3-4 times daily, and thereby eat 5-6 snacks.
- the consumption of less than 50 grams of sugar leads to significantly less caries. On average, current sugar consumption is at 130 grams per person.



Social Support Intervention

Table S-1. *Standardized Daily Support Messages sent by the Confederate.*

Day	Time	Fruit and Vegetable Goal Subgroup	Unhealthy Snack Goal Subgroup
1	16:35	Hi everyone, I just came across this interesting article: http://eatsmarter.de/ernaehrung/news/aepf-el-gesund [Link to info on healthiness of apples] Pretty impressive!	Hi everyone, I just came across this interesting article: http://www.bauch.de/ernaehrung/die-10-schlimmsten-snacks [Link to list of the 10 most terrible snacks] Pretty impressive!
2	11:30	Hey guys, how's it going for you on the second day? Wishing you lots of success with eating more fruits and vegetables! And just reach out if things get rough. I will do the same :)	Hey guys, how's it going for you on the second day? Wishing you lots of success with consuming fewer snacks! And just reach out if things get rough. I will do the same :)
3	20:18	Hi everybody, my recommendation of the day ;) Order a fruits and vegetable subscription, e.g. here: http://www.oepfelchasper.ch/?gclid=CImQnb7U1sECFQ3LtAodpF8AXw Then one will always have fresh fruits and vegetables at home. My roommates really like this.	Hi everybody, my recommendation of the day ;) Remove all unhealthy snacks from your home and don't buy new ones. Then one doesn't even get tempted. This helped me out a lot today.
4	14:25	Hi everyone, I just wanted to say to you that I think it's really great that we have this group. It's good to know that we are all "sitting in the same boat" :) Have a nice day!	Hi everyone, I just wanted to say to you that I think it's really great that we have this group. It's good to know that we are all "sitting in the same boat" :) Have a nice day!
5	09:00	Hey guys, shall we maybe agree on something so that it'll be easier for us to eat more fruits and vegetables? My suggestion: Today, we eat two servings of fruits and vegetables more than usual! Who's in?	Hey guys, shall we maybe agree on something so that it'll be easier for us to snack less? My suggestion: Today, we avoid unhealthy snacks altogether! Who's in?
6	10:15	Hi all, do you also always have difficulties to eat vegetables at dinner when you have plans directly after school and don't get a chance to shop and cook? The other day, I discovered the "Schnägg" at the train station: http://www.xn--schngg-eua.ch/de/ [Link to takeout restaurant]. They have delicious and affordable vegetable dishes to take out.	Hi all, do you also always have difficulties to refrain from eating crisps and such when watching TV? The other day I went to the theatre instead, where it was very easy to avoid snacks! On Mondays it's half price, by the way: http://www.schauspielhaus.ch/karten-abos/preise [Link to local theater webpage]
7	20:55	Did it also work out well for you with eating more fruits and vegetables these last days? We can be proud of each other, I think :) I wish you much success at keeping to eat yummy fruits and vegetables! Stick to it!	Did it also work out well for you with avoiding unhealthy snacks these last days? We can be proud of each other, I think :) I wish you much success at keeping to avoid unhealthy snacks! Stick to it!

Table S-2. Fruit and Vegetable Goal Subgroup: Possible Support Demands and Exemplary Supportive Responses Provided to Confederates.

Support Demands	Supportive Response
How can I integrate more vegetables in my diet?	Just take an extra side salad with your lunch.
How do you cook healthily?	One possibility is to add a lot of vegetables to the main meal. E.g. a broccoli-pasta gratin.
Does anyone know a delicious recipe for vegetables?	Stir fried vegetables! Or a delicious soup! That's yummy and fast to prepare.
What healthy snacks do you eat at the workplace?	Mandarins, carrots, cherry tomatoes. That's pretty varied!
I have had enough of apples for snacks. What kind of snacks do you eat? Do you have other ideas?	Grapes, pears, pineapples, mandarins, oranges, small tomatoes, carrots... there are so many fruits and vegetables ☺ Go to a large supermarket – there's great variety!
I'd like to eat something healthy when I am on the move. But I never have anything with me.	Plan your snacks! You can pack apples or carrots the night before, so you don't forget them ☺
I don't have much time to eat. Today, I had my lunch at the deli with my girlfriend. I just couldn't resist to have French fries and chicken nuggets.	You'll do better next time! Remember that there's also healthy fast food, such as salad or vegetable wraps.
I am usually running late in the morning, and therefore don't have time to prepare a healthy breakfast.	I'm the same. I prepare my muesli with fruits the night before, and take it with me.
I strongly intend to replace sweets with fruits. Do you also find this particularly difficult at lunch? I can pull myself together and not eat sweets, but I just don't feel like eating an apple instead.	I totally understand! A yummy alternative is to cut fruits into small pieces and mix them with plain yogurt!
What do you do when you have cravings or feel like eating chocolate due to stress?	How about a sweet mandarin instead?
Ideas for meals with fruits or vegetables?	I remember the mango soup I cooked last week. Puree the mango, and then cook with coconut milk and ginger. Or vegetable stew with yogurt-garlic dip? Yum, I am getting hungry ☺
I personally find it easier to replace snacks with fruits and vegetables instead of avoiding them. But after a meal it's difficult to eat fruit instead of chocolate with my coffee.	How about dates? They are deliciously sweet ☺

Table S-3. Unhealthy Snack Goal Subgroup: Possible Support Demands and Exemplary Supportive Responses Provided to Confederates.

Support Demands	Supportive Response
I have eaten several unhealthy snacks today. So, I guess it doesn't matter if I completely forget about my good intentions today...	No! It's still worthwhile to stick with your goals!
Are all snacks unhealthy?	No, there are healthy snacks as well. Carrots, for example. And it's a question of proportion.
I'm hungry if I don't eat any snacks. What should I do?	It's really hard sometimes, right? In my experience, it's oftentimes not hunger but thirst! Therefore, drink a glass of water and distract yourself a bit. Perhaps you won't be hungry after that. And if you still are, eat a plain yogurt or so.
I was invited for coffee and didn't want to say no to cake, because it seemed impolite. What do you do to avoid this even though someone invites you and perhaps won't understand why one doesn't want to have cake?	It's natural to feel a bit bad... But it's fine to decline foods you don't want to eat. The others will habituate to that eventually.
Not everyone is understanding. They often offer snacks that I don't want to eat. I feel like I can't say no. How can I avoid this?	I know the situation! Nasty. It might help if you inform your social environment about your intentions to change your diet, and that you would be happy if they would support you.
What are healthy alternative snacks that go with coffee instead of a chocolate bar?	Grapes or banana?
I'm so hungry, but I don't feel like eating something healthy. Hmm... do you think it'd be bad if I ate a chocolate bar right now?	That's very tempting.... Or you eat a dried fruit instead? Come on, we can make it together!
I am at a birthday party with my girlfriend, and there are so many snacks, and of course cake. Is it bad if I eat a piece?	Of course it would be better to remain strong. But a small piece once in a while is okay. Eat slowly and enjoy it!
I find not snacking particularly difficult in the afternoon break. How can I motivate myself for a healthy snack?	Think about how good you will feel if you eat healthily ☺ There are so many healthy snacks! It helps me when I put the fruit in front of me on the desk. ☺
I would love to eat a chocolate croissant... Any suggestions what I could eat instead?	Fruit buns are a delicious and healthy alternative!
What kind of healthy snacks do you eat in at work?	Mandarins or whole wheat crackers. That's really varied!
Argh, I am badly craving some snacks. I think I will have to grant myself something. I am very hard at work at the moment and don't feel like keeping my eating intentions on top of that.	That's understandable. I usually try to remind myself that I feel more energized when I eat fewer unhealthy snacks!

Administrator: Standardized Chat Messages / Actions**Day 1, 8:00 AM**

- 1) Create chat groups, adding the respective logo depending on eating goal of the group



Figure S-1. Logos for chat groups: Eat more fruits and vegetables (left) or avoid unhealthy snacks (right)

- 2) Invite participants to chat group
- 3) Send initial message:

Dear participants!

Welcome to our 7-day WhatsApp chat to support each other to [eat more fruits and vegetables / avoid unhealthy snacks]. The goal of the group is that you support each other during the study to eat more healthily and to be there for each other in difficult situations. You are welcome to exchange helpful tips, recipes or photos related to healthy eating. Please treat the other members of the group the way you'd like to be treated yourselves, and be respectful to one another. I am the group's administrator and will stay mostly in the background. This chat is meant for you, the participants of this group. Please don't forget that the identity of the participants and the exchanged contents must be kept confidential and must not be forwarded to third parties. I wish you lots of success in eating [more fruits and vegetables / fewer unhealthy snacks]!

- 4) Send request that participants introduce themselves:

Please introduce yourselves in order to get to know the group. It's enough if you say your name and what your favorite food is. If you'd like, you are welcome to write more, too. I wish you lots of fun and a pleasant week!

Day 7, 8:00 PM

- 5) Send message to inform about end of chat group:

Thank you for participating in this chat group. I hope you profited and had fun.

Tomorrow morning at 8:00 AM, the chat group will be dissolved. Until then, you have time to say goodbye. Thanks again for your participation in the chat group and lots of success for keeping up with eating [more fruits and vegetables / fewer unhealthy snacks]!

Best regards, your Administrator

Day 8, 8:00 AM

- 6) Dissolve the group by removing all participants and then deleting the chat.

2. Graphic Representation of the Statistical Model




Figure S-2. Graphic representation of the statistical model. b = regression coefficient.

3. SAS Code

Example for the outcome “Servings of Fruits and Vegetables” (fnv)

```
PROC MIXED covtest DATA=fnv METHOD=reml cl;  
CLASS id diary_number group int1 int1copy;  
MODEL fnv=int1 time int1*time int1*spline / solution cl DDF=95, 95, 95, 95;  
RANDOM intercept/SUBJECT=group TYPE=un group=int1copy;  
PARMS (0 1.1601 1.5472 1.5251 0.02986 4.9351)/hold=1;  
RANDOM intercept time /SUBJECT=id(group) TYPE=un;  
REPEATED diary_number/SUBJECT=id TYPE=ar(1);  
  
run;
```

Note. id = person identifier; diary_number = diary day (1-10); group = chat group (1-32, 0 = control group); int1 = intervention condition (1 = social support, 0 = control); int1copy = copy of the intervention condition variable; fnv = servings of fruits and vegetables; time = intervention days (centered at 0 = last intervention day); spline = pre-intervention days (centered at last pre-intervention day). The PARMS statement is necessary to suppress the estimation of the “chat group variance” in the control group as the control group does not contain chat groups.

4. Results: Supplemental Tables and Figures

Descriptive Statistics

Table S-4. *Baseline Descriptive Statistics for the Intervention Groups.*

	Control Group + Snack Goal (<i>n</i> =54)	Control Group + F&V Goal (<i>n</i> =49)	Social Support + Snack Goal (<i>n</i> =52)	Social Support + F&V Goal (<i>n</i> =48)
Mean age (SD)	26.2 (7.8)	29.7 (9.0)	28.1 (9.7)	25.9 (7.4)
Mean Body Mass Index (SD)	22.9 (3.2)	24.4 (5.5)	23.7 (4.1)	23.1 (2.6)
Female (%)	45 (83.3)	38 (77.6)	40 (74.1)	31 (66.0)

Note. F&V = Fruit and vegetable.

Additional Analyses: Phase Effect and Covariates**Table S-5.** *Multilevel Model Including the Intervention Phase Parameter for the Three Outcomes: Fruit and Vegetable Consumption, Unhealthy Snack Consumption, and Healthy Eating.***Fruit and Vegetable Goal Subgroup (n=97)**

	Original Model		Model with Intervention Phase		Model with Covariates	
	Estimate	SE	Estimate	SE	Estimate	SE
Number of observations	790		790		650	
Fixed effects (intercept, slopes)	Estimate	SE	Estimate	SE	Estimate	SE
Intercept (mean control condition Day 10)	3.85***	0.43	3.85***	0.29	2.26	1.60
Intervention effect Day 10	1.43*	0.55	1.43**	0.52	1.26†	0.63
Time (slope control condition)	-0.02	0.04	-0.02	0.04	-0.03	0.04
Intervention * time	0.15*	0.07	0.12	0.08	0.17*	0.08
Intervention * pre-intervention time	-0.27	0.24	-0.28	0.25	-0.39	0.28
Intervention * phase	-	-	-0.26	0.45	-	-
Gender (0=male)	-	-	-	-	-0.59	0.48
Age	-	-	-	-	-0.02	0.03
Vegetarian/vegan (0=no)	-	-	-	-	1.33	0.83
Body mass index	-	-	-	-	0.04	0.05
Restrained eating	-	-	-	-	0.68*	0.29
Stress	-	-	-	-	-0.16	0.33
Social desirability	-	-	-	-	0.17	0.15
Active chat participation (0=no)	-	-	-	-	0.21	0.92

Note. Time: Linear trend from pre-intervention day 1 (= -10) to the last day of the intervention (=0), Pre-intervention time = 3 days prior to intervention. Intervention: 1 = social support, 0 = control.

* $p < .05$, ** $p < .01$, *** $p < .001$, † $p < .10$.

Table S-5. (Continued) *Multilevel Model Including the Intervention Phase Parameter for the Three Outcomes: Fruit and Vegetable Consumption, Unhealthy Snack Consumption, and Healthy Eating.***Unhealthy Snack Goal Subgroup (n=106)**

	Original Model		Model with Intervention Phase		Model with Covariates	
	Estimate	SE	Estimate	SE	Estimate	SE
Number of observations	838		838		665	
Fixed effects (intercept, slopes)	Estimate	SE	Estimate	SE	Estimate	SE
Intercept (mean control condition Day 10)	1.61***	0.18	1.71***	0.38	2.12	1.41
Intervention effect Day 10	-0.80**	0.27	-0.84†	0.44	-1.04**	0.32
Time (slope control condition)	<0.01	0.03	0.02	0.03	0.03	0.03
Intervention * time	-0.14**	0.05	-0.12*	0.06	-0.18**	0.05
Intervention * pre-intervention time	0.24	0.15	0.28	0.17	0.32†	0.18
Intervention * phase	-	-	0.23	0.30	-	-
Gender (0=male)	-	-	-	-	0.09	0.34
Age	-	-	-	-	-0.01	0.01
Vegetarian/vegan (0=no)	-	-	-	-	-0.34	0.37
Body mass index	-	-	-	-	0.02	0.04
Restrained eating	-	-	-	-	-0.36†	0.20
Stress	-	-	-	-	0.07	0.21
Social desirability	-	-	-	-	0.01	0.09
Active chat participation (0=no)	-	-	-	-	0.23	0.78

Note. Time: Linear trend from pre-intervention day 1 (= -10) to the last day of the intervention (=0), Pre-intervention time = 3 days prior to intervention. Intervention: 1 = social support, 0 = control.

* $p < .05$, ** $p < .01$, *** $p < .001$, † $p < .10$.

Table S-5. (Continued) *Multilevel Model Including the Intervention Phase Parameter for the Three Outcomes: Fruit and Vegetable Consumption, Unhealthy Snack Consumption, and Healthy Eating.*

Healthy Eating (z standardized, N=203)						
	Original Model		Model with Intervention Phase		Model with Covariates	
	1628		1628		1315	
Number of observations	Estimate	SE	Estimate	SE	Estimate	SE
Fixed effects (intercept, slopes)						
Intercept (mean control condition Day 10)	-0.17*	0.08	-0.18*	0.08	-0.70	0.48
Intervention effect Day 10	0.48***	0.12	0.49***	0.13	0.54**	0.18
Time (slope control condition)	-0.01	0.01	-0.02	0.01	-0.02†	0.01
Intervention * time	0.06***	0.02	0.06**	0.02	0.08***	0.02
Intervention * pre-intervention time	-0.09†	0.05	-0.09	0.06	-0.13*	0.06
Intervention * phase	-	-	-0.06	0.10	-	-
Gender (0=male)	-	-	-	-	-0.16	0.13
Age	-	-	-	-	<0.01	<0.01
Vegetarian/vegan (0=no)	-	-	-	-	0.31†	0.18
Body mass index	-	-	-	-	<0.01	0.01
Restrained eating	-	-	-	-	0.25**	0.08
Stress	-	-	-	-	-0.04	0.08
Social desirability	-	-	-	-	0.02	0.03
Active chat participation (0=no)	-	-	-	-	-0.08	0.25

Note. Time: Linear trend from pre-intervention day 1 (= -10) to the last day of the intervention (=0), Pre-intervention time = 3 days prior to intervention. Intervention: 1 = social support, 0 = control.

* $p < .05$, ** $p < .01$, *** $p < .001$, † $p < .10$.

Missing Analysis

Table S-6. *Multilevel Model Results for the Original Model, and for Different Dropout Replacement Models for the Three Outcomes: Fruit and Vegetable Consumption, Unhealthy Snack Consumption, and Healthy Eating.*

Fruits and Vegetables, (servings per day), F&V Goal Group (n = 97)				
	Original Model	1st Day Value Carried Forward	3rd Day Value Carried Forward	Last Observation Carried Forward
Fixed effects (intercept, slopes)	Estimate (SE)	Estimate (SE)	Estimate (SE)	Estimate (SE)
Intercept (mean control condition Day 10)	3.85 (0.34)***	3.81 (0.27)***	3.81 (0.27)***	3.81 (0.29)***
Intervention effect Day 10	1.43 (0.55)*	1.35 (0.47)**	1.35 (0.47)**	1.41 (0.49)**
Time (slope control condition)	-0.02 (0.04)	-0.02 (0.03)	-0.02 (0.03)	-0.02 (0.03)
Intervention * time	0.15 (0.07)*	0.14 (0.06)*	0.14 (0.06)*	0.14 (0.06)*
Intervention * pre-intervention time	-0.27 (0.24)	-0.27 (0.21)	-0.27 (0.21)	-0.25 (0.21)

Unhealthy Snacks (servings per day), Snack Goal Group (n = 106)				
	Original Model	1st Day Value Carried Forward	3rd Day Value Carried Forward	Last Observation Carried Forward
Fixed effects (intercept, slopes)	Estimate (SE)	Estimate (SE)	Estimate (SE)	Estimate (SE)
Intercept (mean control condition Day 10)	1.61 (0.18)***	1.64 (0.23)***	1.63 (0.22)***	1.62 (0.22)***
Intervention effect Day 10	-0.80 (0.27)**	-0.55 (0.30)†	-0.54 (0.30)†	-0.69 (0.28)*
Time (slope control condition)	0.01 (0.03)	0.01 (0.03)	0.01 (0.03)	0.01 (0.03)
Intervention * time	-0.14 (0.05)**	-0.12 (0.04)**	-0.12 (0.04)**	-0.12 (0.04)**
Intervention * pre-intervention time	0.24 (0.15)	0.26 (0.14)†	0.26 (0.14)†	0.25 (0.14)†

Note. Time: Linear trend from pre-intervention day 1 (= -10) to the last day of the intervention (=0), Pre-intervention time = 3 days prior to intervention. Intervention: 1 = social support, 0 = control. F&V = fruit and vegetable.

* $p < .05$, ** $p < .01$, *** $p < .001$, † $p < .10$.

Healthy Eating (z standardized), Total Sample (N = 203)

	Original Model	1st Day Value Carried Forward	3rd Day Value Carried Forward	Last Observation Carried Forward
Fixed effects (intercept, slopes)	Estimate (SE)	Estimate (SE)	Estimate (SE)	Estimate (SE)
Intercept (mean control condition Day 10)	-0.17 (0.08)*	-0.17 (0.07)*	-0.17 (0.07)*	-0.17 (0.08)*
Intervention effect Day 10	0.48 (0.12)***	0.39 (0.12)**	0.39 (0.12)**	0.45 (0.12)***
Time (slope control condition)	-0.01 (0.01)	-0.01 (0.01)	-0.01 (0.01)	-0.01 (0.01)
Intervention * time	0.06 (0.02)***	0.06 (0.01)***	0.06 (0.01)***	0.06 (0.01)***
Intervention * pre-intervention time	-0.09 (0.05)†	-0.10 (0.05)*	-0.10 (0.05)*	-0.10 (0.05)*

Note. Time: Linear trend from pre-intervention day 1 (= -10) to the last day of the intervention (= 0), Pre-intervention time = 3 days prior to intervention. Intervention: 1 = social support, 0 = control. F&V = fruit and vegetable.

* $p < .05$, ** $p < .01$, *** $p < .001$, † $p < .10$.

Main Results




Figure S-2. Model predicted intervention effects on self-reported healthy eating over time. Bolded lines represent model predicted fixed effects, lighter circles and error bars represent observed means and 2 times the standard errors. F&V: Fruit and vegetable.