

1 **Production potential of greater duckweed *Spirodela polyrhiza* (L. Schleiden) and its**
2 **biochemical composition evaluation**

3

4

5 JaiGopal Sharma^{a*}, William D. Clark^b, Avanish Kumar Shrivastav^a, Ravi Kumar Goswami^c,
6 Douglas R. Tocher^b, Rina Chakrabarti^c

7

8 ^a*Department of Biotechnology, Delhi Technological University, Delhi 110042, India*

9 ^b*Institute of Aquaculture, Faculty of Natural Sciences, University of Stirling, Stirling FK9*

10 *4LA, Scotland, UK.*

11 ^c*Aqua Research Lab, Department of Zoology, University of Delhi, Delhi 110 007, India*

12

13

14

15

16

17

18

19

20 **Running title:** Production of *Spirodela polyrhiza* and its biochemical composition study

21

22 *Correspondence

23 +91-11-27666496, e-mail: sharmajaigopal@yahoo.com

24

25

26 **ABSTRACT**

27 The culture technique of greater duckweed *Spirodela polyrhiza* (L. Schleiden) was
28 standardized in outdoor tanks using three different manures: manure 1 - cattle manure, poultry
29 droppings and mustard oil cake, manure 2 - urea, potash and triple superphosphate and manure
30 3 - cattle manure, urea, potash and triple superphosphate. Significantly ($p<0.05$) higher
31 production was recorded in manure 1 compared to others. Manure 1 was subsequently selected
32 for pond culture. In ponds, the production of duckweed was 2020 ± 150 kg ha⁻¹ month⁻¹ dry
33 weight basis. Protein content was significantly higher ($p<0.05$) in duckweed cultured in manure
34 1. The amino acid profile study showed the presence of essential (37.4%), non-essential
35 (58.2%) and free (4.5%) amino acids. Leucine, isoleucine and valine contributed 51.4% of total
36 essential amino acids. Duckweed contained 7% lipid and α -linolenic acid (36-37%) was the
37 major fatty acid. The study showed the nutritional value of duckweed as an animal feed
38 ingredient.

39 *Keywords:* *Spirodela polyrhiza*, Organic manure, Proximate composition, Amino acids, Fatty
40 acids

41

42

43

44

45

46

47

48

49

50

51 **1. Introduction**

52 The greater duckweed *Spirodela polyrhiza* (L. Schleiden) is a free-floating, fast
53 growing aquatic plant, widely distributed in the still and slow-flowing water bodies globally.
54 Morphologically, this monocotyledon plant is simple and lack specialised structures such as
55 leaves or stems, but consist of flat ovoid leaf-like structures termed fronds with a rootlet for
56 stabilisation. The bright green (upper part) and purple (lower side) colours of the fronds
57 enhance its aesthetic value and make it suitable candidate for aquarium. Recent study shows
58 the whole genome sequencing of *S. polyrhiza*, the most primitive member of Lemnaceae family
59 (Michael et al., 2017; Hoang et al., 2018). It is a useful tool for further investigation with this
60 duckweed. In accordance with other Lemnaceae, the usefulness and potential of *S. polyrhiza*
61 has been recognized in recent days. It has utilisation for various purposes such as waste water
62 remediation as it is able to remove nitrogen (particularly ammonia) with high efficiency (Culley
63 and Epps, 1973; Sutton and Ornes, 1975, 1977), bio-fuel production (Jarvis et al., 1998; Zhao
64 et al., 2012, 2014) and recombinant protein production (Khvatkov et al., 2018). It is also
65 reported as a promising substrate for bio-hydrogen production, and recognised as an ideal plant
66 in bioremediation and carbon cycle research (Kuehdorf et al., 2014; Olah et al., 2015; Tang et
67 al., 2014; Wang et al., 2012, 2015; Xu and Deshusses, 2015; Xu et al., 2015). The copy number
68 of the genes involved in the biosynthesis of two enzymes glutamine synthetase (GS) and
69 glutamate synthase (GOGAT) are amplified in greater duckweed. GS and GOGAT are the
70 major biochemical module for ammonium assimilation (Wang et al., 2014). In recent year,
71 duckweeds are also considered as rich protein source for human consumption (Appenroth et
72 al., 2018; de Beukelaar et al., 2019).

73 *S. polyrhiza* is also gathering interest as a feed material/ingredient for fish, poultry and
74 pigs (Cruz-Velásquez et al., 2014; FAO, 2001; Hasan and Chakrabarti, 2009). Less fibre
75 content of the plant makes it easily digestible. In grass carp *Ctenopharyngodon idella*, a 75%

76 digestibility of *S. polyrhiza* has been observed (Wee, 1991). Similarly, analysis of the
77 proximate composition showed that *S. polyrhiza* are a rich source of protein, although content
78 varies from 23.8 - 40.9% (Hasan and Edwards, 1992; Hillman and Culley, 1978). Amado et al.
79 (1980) reported the amino acid composition of 94 different strains of duckweeds. They
80 suggested that all essential amino acids (except methionine) are present in sufficient amount in
81 all strains of duckweeds. Recently, Appenroth et al. (2017) found around 25% protein level in
82 *S. polyrhiza* cultured in nutrient medium. They also suggested that the levels of critical amino
83 acids in duckweeds are within the recommended range of World Health Organization (WHO)
84 for human. It is also a rich source of pigments, especially carotene and xanthophylls (Leng et
85 al., 1995). Notably the nutritional and biochemical value of such macrophytes is highly variable
86 and depends largely on water quality of the culture system (Boyd, 1971). Therefore, there is an
87 urgent requirement to develop large-scale culture techniques for the production of nutrient-rich
88 duckweeds. There is immense scope for large scale production of duckweeds in tropical climate
89 (Chakrabarti, 2017).

90 In intensive management, supply of water and nutrient are essential for the continuous
91 duckweed production of a predictable and useful biochemical composition (Hasan and
92 Chakrabarti, 2009). Moreover, duckweeds are commonly cultured in wastewater which may
93 contain unwanted components that are unsuitable for consumption by fish, other livestock, and
94 ultimately human consumers. Inorganic and organic manures were successfully applied in
95 Bangladesh for the production of duckweeds (BFRI, 1997; DWRP, 1998). The aim of the
96 present study is to standardise the culture technique for the production of greater duckweed
97 *Spirodela polyrhiza* in small tanks, and then large-scale production in ponds. The proximate,
98 amino acid and fatty acid profiles of cultured *S. polyrhiza* are evaluated to establish its
99 nutritional quality and suitability as an animal feed ingredient.

100 **2. Materials and methods**

101 2.1. Tank culture

102 *S. polyrhiza* were cultured in cemented outdoor tanks (1.2 m x 0.35 m x 0.30 m) using
103 both organic manures and inorganic fertilizers between December 2016 - March 2017. Three
104 different combinations of manures used for the production of duckweeds were as follows.
105 Manure 1: cattle manure, poultry droppings and mustard oil cake (1:1:1) were used at the rate
106 of 1.052 kg m⁻³ (Srivastava et al., 2006). Manure 2: urea, potash and triple superphosphate were
107 used at the rate of 20, 4 and 4 kg ha⁻¹ day⁻¹, respectively (DWRP, 1998). Manure 3: cattle
108 manure, urea, potash and triple superphosphate were used at the rate of 750, 7.5, 1.5 and 1.5
109 kg ha⁻¹day⁻¹, respectively (BFRI, 1997). There were three replicates for each treatment. *S.*
110 *polyrhiza*, grown in the outdoor facility was introduced in the culture tanks (15 g tank⁻¹, fresh
111 weight) after 5 days of manure application. All tanks were re-manured at 10 day intervals for
112 sustainable duckweed production. In manure 1, organic manures were applied at a rate of one
113 fourth dose of the initial dose. In manure 2 and manure 3, the amount of manure was equal to
114 the initial dose. All manures were decomposed (5 days) before application. In each treatment,
115 when the surface was fully covered, harvesting was initiated, except the fifth harvest in manure
116 3. At the time of fifth harvest, growth of duckweeds was poor in this treatment; duckweeds are
117 totally harvested from all treatments. In all harvests (except the final), 50% of the total
118 duckweeds were collected; all duckweeds were collected after 118 days of culture and the
119 production was recorded as kg ha⁻¹ month⁻¹ (dry weight, DW).

120 2.2. Pond culture

121 Three cemented ponds at the Central Institute of Fisheries Education (Indian Council
122 of Agricultural Research), located at Rohtak, Haryana were used for the production of *S.*
123 *polyrhiza* between July - August 2017. Each pond was 200 m² (20 m x 10 m) with water level
124 maintained as 50 cm. Among the three manures used in the tank culture of greater duckweeds,
125 highest production was obtained in manure 1, and so this treatment was selected for pond

126 production. All the organic manures, cattle manure, poultry dropping and mustard oil cake
127 (Srivastava et al., 2006) were decomposed for 5 days initially. *S. polyrhiza* cultures were
128 produced in a clean environment (outdoor tanks of Department of Zoology, University of
129 Delhi); then the plants were introduced in each pond at the rate of 1 kg pond⁻¹ (fresh weight).
130 Initially, these greater duckweeds covered a small area of the water body (Fig. 1). In each pond,
131 after the initial dose, one fourth dose of manure was applied at intervals of 10 days. Greater
132 duckweeds were harvested thrice at 10 days intervals during 30 days of culture period. The
133 harvesting pattern was similar to tank production, i.e. duckweeds were harvested when the
134 whole water surface was covered. In first and second harvest, 50% duckweeds were harvested
135 and plants were totally collected during the third harvest. The production was expressed as kg
136 ha⁻¹ month⁻¹ (DW).

137 2.3. *Water quality*

138 Major water quality parameters were recorded at weekly intervals in both tanks and
139 ponds. A Solar Light lux meter (PMA 2100, USA) was used for the measurement of light
140 intensity in the outdoor systems at fixed time (10.00 a.m.) and it was expressed as an average
141 of replicates of individual treatment. A HACH Multi-meter (HQ 40d, USA) was used for
142 the estimation of temperature, pH, conductivity, dissolved oxygen, ammonia and nitrate
143 levels. Standard methods were followed for the estimation of phosphate and nitrite levels of
144 water (APHA, 2012).

145 2.4. *Relative Growth Rate (RGR)*

146 The RGR of *S. polyrhiza* was estimated with the formula:

$$147 \text{RGR} = \ln (W_t/W_0)/t$$

148 Where, W_t and W_0 were the weights of duckweeds at time t and zero reference time,
149 respectively; t was the time interval in days. RGR was expressed as g g⁻¹ day⁻¹.

150 2.5. *Biochemical assays*

151 The proximate composition of *S. polyrhiza* was assayed following standard methods
152 (AOAC, 2000). Briefly, samples were dried for 24 h at 110 °C in an oven for the estimation of
153 moisture contents. Ash content was determined after incineration at 600 °C for 16 h. Crude
154 protein content was assayed by Kjeldahl distillation and nitrogen content (N x 6.25) was
155 determined using a Tecator Kjeltac Auto 1030 analyser (Foss, Warrington, UK). Crude lipid
156 level was determined gravimetrically using a Tecator Soxtec 2050 (Foss, Warrington, UK)
157 after Soxhlet extraction by Hydrotec 8000 digester (Foss, Warrington, UK). Carbohydrate
158 content of sample was subsequently determined by subtraction of protein, lipid and ash values.

159 The amino acid profile of greater duckweeds was estimated with an L-8900
160 Automatic Amino Acid Analyser (Hitachi Co. Ltd., Tokyo, Japan). The powdered
161 duckweed sample was first hydrolysed using 6 N HCl for 22 h at 110 °C. Then hydrolysed
162 sample was dried in a Nitrogen Evaporator (PCi Analytics, EV PLUS 08, Maharashtra,
163 India). In the sample, 0.02 N HCl was added and the concentration of protein was 0.5 mg
164 mL⁻¹ of sample. The sample was kept in the Auto sampler and sample injection volume was
165 20 µL. As methionine, cysteine and tryptophan are destroyed during hydrolysis of sample
166 with 6 N HCl, specific reagents are used for the estimations of these amino acids. Performic
167 acid and hydrobromic acid (48%) were used for methionine and cysteine. For tryptophan,
168 the sample was hydrolysed with 4 N methanesulfonic acid and 3-(2-aminoethyl) indole. The
169 remaining methodology was identical for all amino acids. The ninhydrin derivative of
170 proline and hydroxyproline was monitored at 440 nm, and other amino acids were monitored
171 at 570 nm. The amino acids (peak areas) were quantified using the supplied Amino Acids
172 Mixture Standard Solutions, Type B and Type AN-2 (Wako Pure Chemical Industries,
173 Limited, Japan). Standard solutions for glutamine and tryptophan (Sigma-Aldrich, USA)
174 were prepared before analysis.

175 Further *S. polyrhiza* samples were dried at 40 °C and ground prior to extraction of
176 total lipid for fatty acid composition analysis. Total lipid was extracted from 1 g sample
177 (DW) by homogenising in chloroform/methanol (2:1, v/v) using an Ultra-Turrax tissue
178 disrupter (Fisher Scientific, Loughborough, UK), and content determined gravimetrically
179 (Folch et al., 1957). Fatty acid methyl esters (FAME) were prepared from total lipid by acid-
180 catalysed transesterification at 50 °C for 16 h (Christie, 2003), and FAME extracted and
181 purified (Tocher and Harvie, 1988). The FAME were separated and quantified by gas-liquid
182 chromatography using a Fisons GC-8160 (Thermo Scientific, Milan, Italy) equipped with a
183 30 m × 0.32 mm (i.d.) × 0.25 µm ZB-wax column (Phenomenex, Cheshire, UK), on-column
184 injector, and a flame ionisation detector. Data were collected and processed using
185 Chromcard software for Windows (version 2.01; Thermoquest Italia S.p.A., Milan, Italy).
186 Individual FAME was identified by comparison to known standards and published data
187 (Tocher and Harvie, 1988).

188 2.6. Statistical analysis

189 Data were presented as mean ± SE unless otherwise stated. One-way analysis of
190 variance, ANOVA, Duncan's multiple range test, DMR (Montgomery, 1984). Student's t-
191 test were used for the statistical analysis with significance accepted at $p < 0.05$ level.

192 3. Results

193 3.1. Culture in tanks

194 3.1.1. Water quality

195 Major water quality parameters were recorded in all treatments before the application
196 of manures. There was no significant ($p > 0.05$) difference in temperature, pH, dissolved
197 oxygen, ammonia, nitrite, nitrate and phosphate levels among treatments at the beginning of
198 the study. A wide range of water temperature 9.4 - 26.7 °C was recorded during the culture of
199 duckweed between December and March and this influenced the productivity (Table 1). The

200 whole culture period was broadly divided into three phases based on the temperature and light
201 intensity in the culture tanks. In phase I (December 2016 - January 2017), water temperature
202 and light intensity were 16.5 °C and 26.0 $\mu\text{mol photons m}^{-2} \text{ s}^{-1}$ at the beginning and then
203 gradually decreased. The lowest temperature and light intensity were recorded in January. In
204 phase II (February - March 2017) and phase III (March, 2017), water temperature and light
205 intensity showed increasing trends. There was no significant ($p > 0.05$) difference in
206 temperature and light intensity among the three different treatments during the culture period.
207 Among these three different treatments, there was variation in pH in different phases.

208 Significantly ($p < 0.05$) higher dissolved oxygen levels were found with manure 2
209 compared to the other two treatments throughout the study period (Fig. 2A). This group was
210 followed by manure 3 and lowest dissolved oxygen level ($<1 \text{ mg L}^{-1}$) was found in manure 1.
211 Ammonia levels were significantly ($p < 0.05$) higher in manure 1 compared to the other two
212 treatments throughout the study period (Fig. 2B). In manure 1, ammonia levels ranged from
213 1.34 - 30.65, 7.52 - 18.57 and 15.25 - 17.85 mg L^{-1} in the first, second and third phases,
214 respectively. In manure 2, ammonia levels ranged from 1.94 - 9.34, 0.03 - 7.71 and 1.44 - 3.33
215 mg L^{-1} in the first, second and third phases, respectively. In manure 3, ammonia level ranged
216 from 0.17 - 10.97, 0.27 - 4.08 and 0.23 - 0.41 mg L^{-1} in the first, second and third phases,
217 respectively. The lowest range of ammonia levels were found in the third phase regardless of
218 manures.

219 Nitrite level was significantly ($p < 0.05$) higher in manure 2 and manure 3 in the first
220 phase compared to manure 1 (Table 1). There was no significant ($p > 0.05$) difference between
221 these two former groups. In the second and third phases, nitrite levels were significantly ($p <$
222 0.05) higher in manure 2 compared to the other two treatments. Nitrate level was significantly
223 ($p < 0.05$) higher in manure 2 compared to the other two treatments throughout the study period.
224 Phosphate level was significantly ($p < 0.05$) lower in manure 2 compared to the other two

225 treatments throughout the study period (Fig. 2C). Conductivity was significantly ($p < 0.05$)
226 higher in manure 1 compared to the other treatments throughout the study period (Fig. 2D). In
227 manure 1, conductivity ranged from 516 - 1196 $\mu\text{S cm}^{-1}$.

228 3.1.2. Production and relative growth rate (RGR)

229 The production of *S. polyrhiza* was affected by water temperature. The relative growth
230 rate of greater duckweeds was slow (0.02 - 0.04 $\text{g g}^{-1} \text{day}^{-1}$) at the beginning of the culture
231 period due to low temperature regardless of treatments. Greater duckweeds were first harvested
232 after 69 days of initial introduction in all three treatments. As water temperature increased, the
233 growth rate also increased and duckweeds were harvested another four times; second and fourth
234 harvests were performed after 10 days of the respective previous harvest and third and fifth
235 harvests were after 12 days of the respective previous harvest. The RGR values ranged from
236 0.021 - 0.158, 0.007 - 0.12 and -0.024 - 0.129 $\text{g g}^{-1} \text{day}^{-1}$ in manures 1, 2 and 3, respectively
237 throughout the study period. In manure 3, poor growth of plant at fifth harvest compared to the
238 previous one resulted into negative RGR value. The average RGR values were 0.08 ± 0.02 ,
239 0.06 ± 0.03 and 0.07 ± 0.03 $\text{g g}^{-1} \text{day}^{-1}$ in manures 1, 2 and 3, respectively. Total production of
240 duckweeds was significantly ($p < 0.05$) higher in manure 1 compared to the other manures (Fig.
241 3). This group was followed by manure 3 and minimum production was found in manure 2.

242 3.2. Culture in ponds

243 3.2.1. Water quality

244 In three different ponds at the Rohtak centre, water temperature and pH ranged from
245 32.4 - 30.5 $^{\circ}\text{C}$ and 7.76 - 8.30, respectively during the study period. Dissolved oxygen level
246 ranged from 1.25 - 4.57 mg L^{-1} on various days of study. Ammonia, nitrite and nitrate levels
247 of ponds ranged from 5.02 - 17.57, 0.003 - 0.12 and 0.23 - 2.44 mg L^{-1} , respectively. Phosphate
248 level ranged 1.15 - 2.0 mg L^{-1} during the study period (Table 2). Conductivity ranged from
249 1032 - 1251 $\mu\text{S cm}^{-1}$ throughout the culture period of greater duckweed.

250 3.2.2. Production and relative growth rate (RGR)

251 *S. polyrhiza* was harvested three times from the ponds at 10 days intervals (Fig. 4A-B).
252 Greater duckweeds were harvested from the ponds and were cleaned thoroughly with tap water
253 to remove organic material, excess water was removed, air dried and then dried at 40 °C in an
254 oven. Dried duckweed was packed in airtight containers for further use. The RGR values were
255 0.48, 0.14 and 0.03 g g⁻¹ day⁻¹ in the first, second and third harvests, respectively. The average
256 RGR value was 0.22 ± 0.13 g g⁻¹ day⁻¹. Total production was 2020 ± 150 kg ha⁻¹ month⁻¹ on
257 dry matter basis, equivalent to 24 tonnes ha⁻¹ yr⁻¹ (Fig. 5).

258 3.3. Biochemical composition

259 There was a difference in the proximate composition of greater duckweed cultured with
260 organic manures (manure 1) and inorganic fertilizers (manure 2) in tanks. Protein content was
261 significantly ($p < 0.05$) higher, and carbohydrate and ash contents were significantly ($p < 0.05$)
262 lower, in duckweed cultured in manure 1 compared to manure 2 (Table 3). The amino acid
263 profile of greater duckweed cultured in organic manures showed the presence of essential
264 (37.4%), non-essential (58.2%) and free amino acids (4.5%). Among essential amino acids,
265 three branched chain amino acids, leucine, isoleucine and valine contributed 51.4%. Glutamic
266 acid and glutamine consisted 28.3% of the total non-essential amino acids in the greater
267 duckweed. The presence of taurine enhanced the nutritional value of greater duckweed (Table
268 4).

269 The fatty acid composition of *S. polyrhiza* was dominated by polyunsaturated fatty
270 acids (PUFA), which accounted for 47-53% of total fatty acids, primarily α -linolenic acid
271 (ALA, 18:3n-3) at around 36-39% (Table 5). Total saturated fatty acids accounted for 32-39%,
272 followed by linoleic acid (LA, 18:2n-6) at 11-14% and monoenes at 9-11%. As with proximate
273 composition, fatty acid profile was affected by manures. *S. polyrhiza* grown in inorganic
274 fertilizers (manure 2) having a higher proportion of ALA, LA and total PUFA, and lower

275 saturated and monounsaturated fatty acids. Due to the slightly higher (although not statistically
276 significant) lipid content of *S. polyrhiza* grown in manure 2, all fatty acids were in higher
277 absolute amounts ($\text{mg}\cdot 100\text{g}^{-1}$ dry mass) in macrophytes grown in inorganic fertilizers. *S.*
278 *polyrhiza* lipid contained no long-chain PUFA such as docosahexaenoic acid (22:6n-3),
279 although there was a trace level of eicosapentaenoic acid (20:5n-3), most likely due to minor
280 microalgal contamination within the macrophyte biomass.

281 **4. Discussion**

282 Water temperature and sunlight are major environmental factors that influence the
283 growth of duckweed compared to the nutrient concentrations in the water (Hasan and
284 Chakrabarti, 2009). In tank culture, *S. polyrhiza* was first harvested after 69 days of culture.
285 The water temperature was generally below 15 °C during this period of culture, and lowest
286 light intensity was also recorded during this period. Water temperature increased above 16 °C
287 at the second phase of culture and only then duckweed grew well and harvested. Higher light
288 intensity was also recorded at the second phase compared to the first one. In a comparative
289 study, growth performance of *S. polyrhiza* was recorded at two temperature ranges of 10 - 12
290 and 26 - 28 °C (Song et al., 2006). It was found that cell growth, the synthesis, and absorption
291 ability of duckweed decreased at low temperature compared to duckweed cultured at higher
292 temperature. There was no change in frond number for 15 days at low temperature range.

293 In the present study, the relative growth rate (RGR) of greater duckweed was low during
294 the first phase of tank culture and then increased regardless of treatments. In manure 3, RGR
295 reduced in fifth harvest of phase three. Among the three manures, significantly ($p < 0.05$)
296 higher production was found with manure 1 compared to the inorganic fertilizers. Therefore,
297 organic manures were applied in pond culture of greater duckweed. In contrast to the tank
298 culture, RGR value was maximum at first harvest in pond culture of greater duckweed and the
299 average RGR value was higher in pond compared to tank production. The production rate of

300 greater duckweed was 0.08 ± 0.02 fronds day⁻¹ in laboratory conditions (Lemon et al., 2001).
301 Higher temperature also resulted in enhanced growth rate in ponds in the present study. In
302 Bangladesh, highest growth of *S. polyrrhiza* was found at 22.2 - 22.5 °C in pond (Khondker et
303 al., 1993), although *S. polyrrhiza* survived at 10 - 12 °C, it could not grow well at a low
304 temperature (Song et al., 2006). The duckweed exposed to oxidative damage at low
305 temperature. Appenroth (2002) suggested that 15 °C temperature (combined with 30 µM
306 phosphate level) was the dominant turion formation inducing factor. In laboratory axenic
307 culture, *S. polyrrhiza* were exposed at 100 µmol m⁻² s⁻¹ white light (Appenroth et al., 2017). In
308 the present study, good growth of *S. polyrrhiza* was found at light intensity between 105 - 151
309 µmol photons m⁻² s⁻¹ in natural outdoor light.

310 In Bangladesh and India, a pH range from 6.5 - 7.5 (Islam and Khondkar, 1991) and
311 6.8 - 8.5 (Gopal and Chamanlal, 1991; Kaul and Bakaya, 1976) was found to be optimum for
312 the production of greater duckweed. In the present study, pH ranged from 6.98 - 7.86 and 7.76
313 - 8.30 in tank and pond culture systems, respectively. There was no direct effect of dissolved
314 oxygen on the production of greater duckweed as highest production was recorded in manure
315 1 with minimum dissolved oxygen level in tank culture. Leng et al. (1995) suggested that
316 maintenance of low dissolved oxygen with 6 - 7 pH should be the strategy for duckweed pond
317 management.

318 It was found that the root length was shorter in *S. polyrrhiza* that grown at low
319 temperature compared to the plants grown at a higher temperature. *S. polyrrhiza* with shorter
320 root length were inefficient in absorbing nitrogen, phosphorus and other nutrients from water
321 (Reddy and DeBusk, 1985). In tank culture, highest ammonia level was recorded in manure 1
322 at first phase and no production was recorded during this period. The ammonia level gradually
323 reduced in the second and third phases and the growth of duckweed enhanced. Even with the
324 same manure system (manure 1), lower levels of ammonia were found in ponds compared to

325 tanks. Absorption of nutrients helped in the higher production of duckweeds in ponds. The
326 fluctuation of pH between 7.4 and 9.0 enhanced the ammonia toxicity in laboratory culture
327 (Caicedo et al., 2000). In tank culture of duckweed, highest RGR was found in the second
328 phase at $15.25 \pm 1.0 \text{ mg L}^{-1}$ ammonia concentration in manure 1. It is also interesting to see
329 that in tank culture, poor growth of duckweeds in manure 3 during fifth harvest might be related
330 to the low ammonia level in the culture tank. Leng et al. (1995) suggested that 7 - 12 mg N L⁻¹
331 was optimum to maintain a protein content of 40% in duckweed. A TKN content of 20 - 30
332 mg L⁻¹ was required for optimum growth (Culley et al., 1981) and maintenance of high protein
333 content. In the present study, the ammonia level in the pond water also helped in the proper
334 growth of the duckweed. Nitrification rate was slower in manure 1 compared to the other two
335 treatments in the tank culture of duckweed. In manure 1, nitrate level was significantly higher
336 in the second phase compared to the other phases. Phosphorus is a major limiting nutrient,
337 although it is required in lesser amount. In the present study, the phosphate levels in manure 1
338 helped in the production of duckweed in both tanks and ponds. The optimum conductivity for
339 maximum production of *S. polyrrhiza* was 650 - 1000 $\mu\text{S cm}^{-1}$ (Gopal and Chamanlal, 1991).
340 *S. polyrrhiza* completely disappeared in May due to reduced conductivity and alkalinity
341 (Khondker et al., 1993). In the tank culture, the growth of greater duckweed was less in the
342 first phase and the conductivity was minimum during this phase regardless of manures applied.
343 Then conductivity increased with higher production of duckweed. In pond culture, the
344 conductivity was always $>1000 \mu\text{S cm}^{-1}$.

345 In ponds, the production of greater duckweed was encouraging, $2020 \pm 150 \text{ kg ha}^{-1}$
346 month⁻¹ ($24 \text{ tonnes ha}^{-1} \text{ yr}^{-1}$) on dry matter basis. Literature showed a wide variation in the
347 production of duckweed, with various climatic conditions and nutrient availability mostly
348 being responsible for this variation. Edwards et al. (1990) reported $\sim 20 \text{ tonnes ha}^{-1} \text{ year}^{-1}$ (DM)
349 production of *S. polyrrhiza* during 1-3 months culture period; the yield decreased ($\sim 9 \text{ tonnes ha}^{-1}$

350 ¹ year⁻¹) when the duration of culture period increased to 6 months. The yield of greater
351 duckweeds in domestic wastewater (Reddy and Debusk, 1985), sewage effluent (Sutton and
352 Ornes, 1975) and nutrient non-limited water (Reddy and DeBusk, 1985) were 17 - 32, 14.6 and
353 11.3 tonnes ha⁻¹ yr⁻¹, respectively. Based on the available data, an average harvest of 10 - 20
354 tonnes duckweed ha⁻¹ year⁻¹ could be expected under optimum environmental conditions
355 (Hasan and Chakrabarti, 2009). In a similar study, *Lemna minor* was produced in ponds using
356 organic manures. The production was lower (702.5 kg ha⁻¹ month⁻¹, DW) compared to *S.*
357 *polyrhiza* (Chakrabarti et al., 2018). The initial amount of duckweed introduced for culture also
358 influenced production. A seeding rate of 60 kg m⁻² for *S. polyrhiza* was recommended (DWRP,
359 1998). In the pond culture, only 1 kg pond⁻¹ (200 m²) *S. polyrhiza* was introduced in the present
360 study.

361 The proximate composition of greater duckweed varied with nutrient availability of the
362 culture system. In the present study, the protein, lipid, ash and carbohydrate contents of greater
363 duckweeds were influenced by the quality of the manures. The protein content of the
364 duckweeds (30.5 ± 0.03 - 35.82 ± 0.14%) was higher in the present study compared to some
365 previous studies. The duckweeds collected from Thailand showed 23.8 ± 0.8% protein content
366 (Hasan and Edwards, 1992), whereas 25.6 ± 0.2% protein content was recorded in plants
367 collected from a pond in Nigeria (Fasakin et al., 1999). In USA, 13.1% crude protein was found
368 in greater duckweed collected from low-nutrient lagoon (Culley et al., 1981), whereas 40.9%
369 crude protein was found in plants grown in a dairy cattle-waste lagoon (Hillman and Culley,
370 1978). In the present study, lipid contents of duckweeds ranged from 7.11 - 7.2%, whereas lipid
371 contents of 2.5 - 6.7% were reported in the earlier studies (Hasan and Chakrabarti, 2009).
372 Appenroth et al. (2017) found around 5% lipid content in duckweed. Similarly, the ash content
373 of the duckweed in the present study (18.51 ± 0.02 - 20.64 ± 0.26%) was comparable with

374 earlier studies, in which ash contents varied from 15.2 ± 0.4 - $18.3 \pm 1.0\%$ in greater duckweeds
375 collected from different geographical areas (Hasan and Edwards, 1992).

376 These data showed that culture of greater duckweed with a specific management
377 strategy helped in the production of valuable animal feed ingredients. *S. polyrhiza* is a new
378 generation sustainable crop (Hoang et al., 2018). Song et al (2006) reported that temperature
379 also influenced the soluble protein, chlorophyll α , chlorophyll β and carotenoid pigment of
380 duckweeds. The present study confirmed the earlier study. The presence of essential amino
381 acids viz. histidine, isoleucine, leucine, lysine, methionine, phenylalanine, threonine, and
382 valine were documented in greater duckweeds (Ismail, 1998). The present study showed that
383 all the essential (including tryptophan) and non-essential amino acids were present in adequate
384 quantity in cultured duckweed. The present study also showed the presence of taurine in the
385 duckweeds. The presence of glutamic acid and glutamine confirmed the role of greater
386 duckweed in reducing nitrogenous materials in the water. Similar amino acids composition was
387 found in *L. minor* (Chakrabarti et al., 2018). The nutritional value of duckweed is comparable
388 with alfalfa, being a rich source of lysine and arginine (Guha, 1997). The composition of
389 essential amino acids in greater duckweed is comparable with soybean (NRC, 1998), the most
390 commonly used ingredient in the diet formulation of fish (Table 6). The amino acid
391 requirements of important cultivable species are documented (NRC, 2011). It is clear from the
392 present study that the amino acid profile of greater duckweed meets the nutritional
393 requirements of the cultivable species. The amino acid profiles of *Landoltia punctata* (= *S.*
394 *oligorhiza*) and different clones of *Wolffia arrhiza* were sufficient to fulfilled the requirements
395 for human recommended by WHO ((Ismail, 1998; Appenroth et al., 2018).

396 In addition, *S. polyrhiza* demonstrated reasonable lipid content with ALA being the
397 major fatty acid component in present study. Inorganic fertilizers resulted in slightly higher
398 lipid content and relative percentage of ALA, which individually did not reach statistical

399 significance, but together had a significant effect, increasing the absolute content of ALA. The
400 PUFA content of *S. polyrhiza* grown in culture media was higher compared to the present study
401 though the total lipid level was higher in the latter (Appenroth et al., 2017). It was interesting
402 that in different species of *Wolffia* fat content was low, varied from 1-5%. PUFA levels were
403 above 60% of total fat. The n-3 PUFA level was higher compared to n-6 PUFA (Appenroth et
404 al., 2018). In the present study, the lipid and PUFA contents were higher in *S. polyrhiza*
405 compared to *Wolffia* spp. In *L. minor*, 60 - 63% of total fatty acid was PUFA; around 41-43%
406 α -linolenic acid and 17-18% linoleic acid (Chakrabarti et al., 2018).

407 **5. Conclusions**

408 The application of organic manures helped in the production of greater duckweed *S.*
409 *polyrhiza* in a sustainable manner. The temperature, light intensity, ammonia, phosphate
410 and conductivity significantly influenced the productivity of the water bodies. Proximate
411 composition, especially amino acid and fatty acid profiles confirmed the suitability of the
412 greater duckweed as a potential ingredient for the development of diets for fish and other
413 livestock.

414 **Acknowledgements**

415 The present investigation was supported by Department of Biotechnology (DBT),
416 Government of India, New Delhi, India (Dy. No. 102/IFD/SAN/4678/2015-2016, dated
417 28.3.2016) and the Biotechnology and Biological Science Research Council (BBSRC)
418 Newton Fund Global Research Partnership Project (BB/N005031/1) “Development of
419 alternative sustainable fish feeds to promote human health using novel non-conventional
420 indigenous ingredients (SNIPH)”. Authors are thankful to the Director, CIFE (ICAR) for
421 providing ponds and other facilities to conduct the study.

422 **References**

423 Amado, R., Muller-Heimeyer, R., Marti, U., 1980. Proteingehalt, aminosäure-
424 zusammensetzung and neutralzucker gehalt von lemnaceen. Veroff. Geobot. Inst.
425 ETH Stiftung Rubel, Zurich. 70, 102–117.

426 AOAC, Association of Official Analytical Chemists, 2000. Official Methods of Analysis.
427 Association of Official Analytical Chemists Inc., Washington, DC.

428 APHA, American Public Health Association, 2012. Standard Methods for the Examination
429 of Water and Waste Water. Twenty second ed. American Public Health
430 Association, American Water Works Association, Water Environment Federation.
431 Washington DC.

432 Appenroth, K.J., 2002. A co-action of temperature and phosphate in inducing turion
433 formation in *Spirodela polyrhiza* (Great duckweed). Plant Cell Environ. 25, 1079–
434 1085. doi: [10.1046/j.1365-3040.2002.00885.x](https://doi.org/10.1046/j.1365-3040.2002.00885.x)

435 Appenroth, K.J., Sree, K.S., Böhm, V., Hammann, S., Vetter, W., Leiterer, M., Jahreis, G.,
436 2017. Nutritional value of duckweeds (Lemnaceae) as human food. Food
437 chem. 217, 266–273. <https://doi.org/10.1016/j.foodchem.2016.08.116>

438 Appenroth, K.J., Sree, K.S., Bog, M., Ecker, J., Seeliger, C., Böhm, V., Lorkowski, S.,
439 Sommer, K., Vetter, W., Tolzin-Banasch, K., Kirmse, R., 2018. Nutritional value
440 of the duckweed species of the genus *Wolffia* (Lemnaceae) as human food. Front.
441 Chem. 6, 483. <https://doi.org/10.3389/fchem.2018.00483>

442 BFRI, 1997. Bangladesh Fisheries Research Institute. Research Progress Report.

443 Boyd, C.E., 1971. The limnological role of aquatic macrophytes and their relationship to
444 reservoir management. Reservoir Fisheries and Limnology Special Publication No.
445 8, American Fisheries Society. pp. 153–166.

446 Caicedo, J.R., Vander Steen, N.P., Arce, O., Gijzen, H.J., 2000. Effect of total ammonia
447 nitrogen concentration and pH on growth rates of duckweed (*Spirodela*
448 *polyrrhiza*). Water Res. 34, 3829–3835.

449 Chakrabarti, R., 2017. Culture of zooplankton and aquatic macrophytes as non-conventional
450 livelihood, in: Dhanze, R., Ninawe, A.S., Dhanze, J.R. (Eds.), Aquaculture for
451 Nutritional and Livelihood Security. Narendra Publishing House, New Delhi, pp.
452 189–203.

453 Chakrabarti, R., Clark, W. D., Sharma, J. G., Goswami, R. K., Shrivastav, A. K., Tocher,
454 D. R., 2018. Mass production of *Lemna minor* and its amino acid and fatty acid
455 profiles. Front. Chem. 6, 479. doi: [10.3389/fchem.2018.00479](https://doi.org/10.3389/fchem.2018.00479)

456 Christie, W.W., 2003. Lipid Analysis, third ed. Oily Press, Bridgewater, UK.

457 Cruz-Velásquez, Y., Kijora, C., Vergara-Hernandez, W., Schulz, C., 2014. On-farm
458 evaluation of Cachama blanca and Nile tilapia fed fermented aquatic plants in a
459 polyculture. Orinoquia 18, 269–277.

460 Culley, D.D., Epps, A.E., 1973. Use of duckweeds for waste treatment and animal feed. J.
461 Water Pollut. Control Fed. 45, 337–347.

462 Culley, D.D., Rejmnkov, E., Event, J., Frye, J.B., 1981. Production, chemical quality and
463 use of duckweeds (Lemnaceae) in aquaculture, waste management and animal
464 feeds. J. World Aquacult. Soc. 12, 27–49. [https://doi.org/10.1111/j.1749-](https://doi.org/10.1111/j.1749-7345.1981.tb00273.x)
465 [7345.1981.tb00273.x](https://doi.org/10.1111/j.1749-7345.1981.tb00273.x)

466 de Beukelaar, M.F., Zeinstra, G.G., Mes, J.J., Fischer, A.R., 2019. Duckweed as human
467 food. The influence of meal context and information on duckweed acceptability of
468 Dutch consumers. Food Qual. Prefer. 71, 76–86.
469 <https://doi.org/10.1016/j.foodqual.2018.06.005>

470 DWRP, 1998. Duckweed Research Project. Duckweed in Bangladesh; Ministry of Fisheries
471 and Livestock, Government of Bangladesh and Royal Netherlands Embassy.
472 Dhaka, Bangladesh, pp. 91.

473 Edwards, P., Pacharaprakiti, C., Yomjind, M., 1990. Direct and indirect use of septage for
474 culture of Nile tilapia *Oreochromis niloticus*. Asian Fisheries Society, pp. 165–
475 168.

476 FAO, Food and Agricultural Organization, 2001. Duckweed, A tiny aquatic plant with
477 enormous potential for agriculture and environment. Food and Agricultural
478 Organization of the United Nations, Geneva.

479 Fasakin, E.A., Balogun, A.M., Fasuru, B.E., 1999. Use of duckweed, *Spirodela polyrrhiza*
480 L. Schleiden, as a protein feedstuff in practical diets for tilapia, *Oreochromis*
481 *niloticus* L. Aquacult. Res. 30, 313–318.

482 Folch, J., Lees, M., Sloane-Stanley, G.H., 1957. A simple method for the isolation and
483 purification of total lipids from animal tissues. J. Biol. Chem. 226, 497–509.

484 Gopal, B., Chamanlal, 1991. Distribution of aquatic macrophytes in polluted water bodies
485 and their bio-indicator value. Verh. Int. Verein. Limnol. 24, 2125–2129.

486 Guha, R., 1997. Duckweeds. Envis Newsletter, Indian Institute of Science, Bangalore, pp.
487 5–9.

488 Hasan, M.S., Edwards, P., 1992. Evaluation of duckweed (*Lemna perpusilla* and *Spirodela*
489 *polyrrhiza*) as feed for Nile tilapia (*Oreochromis niloticus*). Aquaculture 104, 315–
490 326.

491 Hasan, M.R., Chakrabarti, R., 2009. Use of algae and aquatic macrophytes as feed in small-
492 scale aquaculture: a review, FAO fisheries and aquaculture technical paper 531.
493 Food and Agriculture Organization of the United Nations, Geneva.

494 Hillman, W.S., Culley, D.D., 1978. The uses of duckweed. American Scientist 66, 442–451.

495 Hoang, P.N., Michael, T.P., Gilbert, S., Chu, P., Motley, S.T., Appenroth, K. J., Schubert,
496 I., Lam, E., 2018. Generating a high-confidence reference genome map of the
497 greater duckweed by integration of cytogenomic, optical mapping, and oxford
498 nanopore technologies. *Plant J.* 96, 670–684. doi: 10.1111/tpj.14049

499 Ismail, M., 1998. Chemical characterization of protein concentrates of duckweed (Family
500 Laminaceae). *Pertanika JST.* 6, 7–21.

501 Islam, A.K.M.N., Khondkar, M., 1991. Preliminary limnological investigations of some
502 polluted water covered by duckweeds. *Bangladesh J. Bot.* 20, 73–75.

503 Jarvis, M.J., Jenkins, B., Rodgers, G.A., 1998. Southern hemisphere observations of a long
504 - term decrease in F region altitude and thermospheric wind providing possible
505 evidence for global thermospheric cooling. *J. Geophys. Res.* 103, 20775–20787.
506 doi: 10.1029/98JA01629

507 Kaul, V., Bakaya, U., 1976. Noxious, floating, lemnid *Salvinia* aquatic weed complex in
508 Kashmir, in: Varshney, C. K., Rzoska, J. (Eds.), *Aquatic Weeds in South East Asia.*
509 *Proceedings of Regional Seminar on Noxious Aquatic Vegetation, New Delhi, 12-*
510 *17 December 1973.* The Hague: Dr. W. Junk B. V. Publishers, pp. 183–192.

511 Khondker, M., Islam, A.K.M.N., Nahar, N., 1993. Study on the biomass of *Spirodela*
512 *polyrrhiza* and the related limnological factors of some polluted waters, in: Khan,
513 M. S., Aziz Khan, M.A., Hadiuzzaman, S., Aziz, A. (Eds.), *Plants for the*
514 *Environment. Proceedings of the 7th Botanical Conference, 13-14 December,*
515 *1992, Bangladesh Botanical Society: Dhaka, Bangladesh, pp. 37–40.*

516 Khvatkov, P., Firsov, A., Shvedova, A., Shaloiko, L., Kozlov, O., Chernobrovkina, M.,
517 Pushin, A., Tarasenko, I., Chaban, I., Dolgov, S., 2018. Development of *Wolffia*
518 *arrhiza* as a producer for recombinant human granulocyte colony-stimulating
519 factor. *Front. Chem.* 6, 304. <https://doi.org/10.3389/fchem.2018.00304>

520 Kuehdorf, K., Jetschke, G., Ballani, L., Appenroth, K. J., 2014. The clonal dependence of
521 turion formation in the duckweed *Spirodela polyrhiza* - an ecogeographical
522 approach. *Physiol. Plant.* 150, 46–54. doi: 10.1111/ppl.12065.

523 Lemon, G.D., Posluszny, U., Husband, B.C., 2001. Potential and realized rates of vegetative
524 reproduction in *Spirodela polyrhiza*, *Lemna minor*, and *Wolffia borealis*. *Aquat.*
525 *Bot.* 70, 79–87. doi: 10.1016/S0304-3770(00)00131-5

526 Leng, R.A., Stambolie, J.H., Bell, R., 1995. Duckweed - a potential high-protein feed
527 resource for domestic animals and fish, in: Proceedings of the 7th Animal Science
528 Congress of the Asian-Australasian Association of Animal Production Societies
529 (AAAP) Conference. Indonesian Society of Animal Science: Bali. Jakarta, pp.
530 100–117.

531 Michael, T. P., Bryant, D., Gutierrez, R., Borisjuk, N., Chu, P., Zhang, H., Xia, J., Zhou, J.,
532 Peng, H., El Baidouri, M., ten Hallers, B., 2017. Comprehensive definition of
533 genome features in *Spirodela polyrhiza* by high-depth physical mapping and short-
534 read DNA sequencing strategies. *Plant J.* 89, 617–635. doi: 10.1111/tpj.13400

535 Montgomery, D.C., 1984. Design and Analysis of Experiments, John Wiley, New York.

536 NRC, 1998. National Research Council. Nutrient Requirements of Swine, The National
537 Academic Press, Washington.

538 NRC, 2011. National Research Council. Nutrient Requirements of Fish and Shrimps, The
539 National Academic Press, Washington.

540 Olah, V., Hepp, A., Meszaros, I., 2015. Comparative study on the sensitivity of turions and
541 active fronds of giant duckweed (*Spirodela polyrhiza* (L.) Schleiden) to heavy
542 metal treatments. *Chemosphere* 132, 40–46. doi:
543 10.1016/j.chemosphere.2015.01.050.

544 Reddy, K.R., DeBusk, W.F., 1985. Nutrient removal potential of selected aquatic
545 macrophytes. J. Environ. Qual. 14, 459–462. doi:
546 10.2134/jeq1985.00472425001400040001x

547 Song, G., Hou, W., Wang, Q., Wang, J., Jin, X., 2006. Effect of low temperature on
548 eutrophicated water body restoration by *Spirodela polyrhiza*. Bioresour. Technol.
549 97, 1865–1869. <https://doi.org/10.1016/j.biortech.2005.08.012>

550 Srivastava, A., Rathore, R.M., Chakrabarti, R., 2006. Effects of four different doses of
551 organic manures in the production of *Ceriodaphnia cornuta*. Bioresour. Technol.
552 97, 1036–1040. doi:10.1016/j.biortech.2005.04.044

553 Sutton, D.L., Ornes, W.H., 1975. Phosphorous removal from static sewage effluent using
554 duckweed. J. Environ. Qual. 4, 367–370.
555 doi:10.2134/jeq1975.00472425000400030018x

556 Sutton, D.L., Ornes, W.H., 1977. Growth of *Spirodela polyrhiza* in static sewage effluent.
557 Aquat. Bot. 3, 231–237. doi:10.1016/0304-3770(77)90025-0

558 Tang, J., Zhang, F., Cui, W., Ma, J., 2014. Genetic structure of duckweed population of
559 *Spirodela*, *Landoltia* and *Lemna* from Lake Tai, China. Planta 239, 1299–1307.
560 doi: 10.1007/s00425-014-2053-y.

561 Tocher, D.R., Harvie, D.G., 1988. Fatty acid compositions of the major phosphoglycerides
562 from fish neural tissues; (n-3) and (n-6) polyunsaturated fatty acids in rainbow trout
563 (*Salmo gairdneri*) and cod (*Gadus morhua*) brains and retinas. Fish Physiol.
564 Biochem. 5, 229–239. doi: 10.1007/BF01874800

565 Wang, W., Wu, Y., Messing, J., 2012. The mitochondrial genome of an aquatic plant,
566 *Spirodela polyrhiza*. PLoS One. 7, 135–139. doi:10.1371/journal.pone.0046747

567 Wang, W., Haberer, G., Gundlach, H., Gläßer, C., Nussbaumer, T., Luo, M.C., Lomsadze,
568 A., Borodovsky, M., Kerstetter, R.A., Shanklin, J., Byrant, D.W., Mockler, T.C.,

569 Appenroth, K.J., Grimwood, J., Jenkins, J., Chow, J., Choi, C., Adam, C., Cao, X.-
570 H., Fuchs, J., Schubert, I., Rokhsar, D., Schmutz, J., Michael, T.P., Mayer, K.F.X.,
571 Messing, J., 2014. The *Spirodela polyrhiza* genome reveals insights into its
572 neotenus reduction fast growth and aquatic lifestyle. Nat. Commun. 5, 3311.
573 <https://doi.org/10.1038/ncomms4311>

574 Wang, W., Yang, C., Tang, X., Zhu, Q., Pan, K., Cai, D., Hu, Q., Ma, D., 2015. Carbon and
575 energy fixation of great duckweed *Spirodela polyrhiza* growing in swine
576 wastewater. Environ. Sci. Pollut. Res. Int. 22, 15804–15811. doi: 10.1007/s11356-
577 015-4778-y.

578 Wee, K.L., 1991. Use of non-conventional feedstuffs of plant origin as fish feeds - is it
579 practical and economically feasible, in: De Silva, S. S. (Eds.), Fish Nutrition
580 Research in Asia, Special Publication No. 5. Asian Fisheries Society, pp. 13–32.

581 Xu, J., Deshusses, M.A., 2015. Fermentation of swine wastewater-derived duckweed for
582 biohydrogen production. Int. J. Hydrog. Energy 40, 7028–7036.
583 <https://doi.org/10.1016/j.ijhydene.2015.03.166>

584 Xu, X.J., Sun, J.Q., Nie, Y., Wu, X.L., 2015. *Spirodela polyrhiza* stimulates the growth of
585 its endophytes but differentially increases their fenprothrin-degradation
586 capabilities. Chemosphere 125, 33–40. doi: 10.1016/j.chemosphere.2014.12.084

587 Zhao, X., Elliston, A., Collins, S.R.A., Moates, G.K., Coleman, M.J., Waldron, K.W., 2012.
588 Enzymatic saccharification of duckweed (*Lemna minor*) biomass without
589 thermophysical pretreatment. Biomass Bioenergy 47, 354–361.
590 <https://doi.org/10.1016/j.biombioe.2012.09.025>

591 Zhao, X., Moates, G.K., Wellinder, N., Collins, S.R.A., Coleman, M.J., Waldron, K.W.,
592 2014. Chemical characterization and analysis of the cell wall polysaccharides of
593 duckweed (*Lemna minor*). Carbohydr. Polym. 111, 410–418.

594 <https://doi.org/10.1016/j.carbpol.2014.04.079>

595

596

597

598

599

600

601

602

603

604

605

606

607

608

609

610

611

612 **Figure legends**

613 **Fig. 1** Introduction of *S. polyrhiza* (1 kg pond⁻¹) in Rohtak, Haryana.

614 **Fig. 2** Various water quality parameters (in parenthesis). (A) Dissolved oxygen, (B)
615 ammonia, (C) phosphate and (D) conductivity of water found during three different phases
616 of culture of *S. polyrhiza* in tanks. Phase I: December 2016 - January 2017, Phase II:
617 February - March 2017 & Phase III: March 2017. Bars with different superscripts are
618 significantly ($p < 0.05$) different (n = 3).

619 **Fig. 3** Total production of *S. polyrhiza* cultured with three different organic manures and
620 inorganic fertilizers in tanks. Bars with different superscripts are significantly ($p < 0.05$)
621 different ($n = 3$).

622 **Fig. 4** Production of *S. polyrhiza* (A) in ponds & (B) duckweeds after harvest.

623 **Fig. 5** Relative growth rate (RGR) and total production of *S. polyrhiza* in ponds. RGR was
624 measured thrice at 10 days interval. Bars with different superscripts are significantly ($p <$
625 0.05) different ($n = 3$).

626 **Table 1 Environmental parameters measured in tanks during the culture of *S. polyrhiza*.**

627

Parameters	Manure 1		Manure 2		Manure 3	
	Range	Mean \pm SE	Range	Mean \pm SE	Range	Mean \pm SE
Phase I (December 2016 - January 2017)						
Temperature ($^{\circ}$ C)	9.36 – 16.55	14.38 \pm 0.34	9.36 – 16.55	14.38 \pm 0.336	9.36 – 16.55	14.38 \pm 0.34
Light intensity (μ mol photons m^{-2} s^{-1})	14.56 – 49.43	27.29 \pm 2.45	14.56 – 49.43	27.29 \pm 2.45	14.56 – 49.43	27.29 \pm 2.45
pH	7.20 – 7.91	---	7.04 – 7.86	---	6.98 – 7.85	---
Nitrite (mg L^{-1})	0.007 – 0.26	0.116 \pm 0.01	0.13 – 1.01	0.47 \pm 0.05	0.06 – 1.04	0.49 \pm 0.07
Nitrate (mg L^{-1})	1.68 – 18.70	5.77 \pm 1.13	6.58 – 43.66	30.76 \pm 2.60	8.44 – 35.48	24.40 \pm 2.01
Phase II (February - March 2017)						
Temperature ($^{\circ}$ C)	15.70 – 19.33	17.80 \pm 0.43	15.70 – 19.33	17.80 \pm 0.43	15.70 – 19.33	17.80 \pm 0.43
Light intensity (μ mol photons m^{-2} s^{-1})	49.21 – 105.08	89.89 \pm 3.25	49.21 – 105.08	89.89 \pm 3.25	49.21 – 105.08	89.89 \pm 3.25
pH	7.09 – 7.59	---	7.24 – 7.82	---	7.26 – 7.72	---
Nitrite (mg L^{-1})	0.02 – 0.12	0.055 \pm 0.015	0.11 – 0.84	0.44 \pm 0.08	0.006 – 0.12	0.09 \pm 0.02
Nitrate (mg L^{-1})	5.95 – 44.73	25.77 \pm 6.04	15.04 – 46.87	29.38 \pm 4.58	16.15 – 34.94	24.42 \pm 2.61
Phase III (March 2017)						
Temperature ($^{\circ}$ C)	23.26 – 26.70	24.98 \pm 1.72	23.26 – 26.70	24.98 \pm 1.72	23.26 – 26.70	24.98 \pm 1.72
Light intensity (μ mol photons m^{-2} s^{-1})	137.41 – 151.16	143.79 \pm 6.39	137.41 – 151.16	143.79 \pm 6.39	137.41 – 151.16	143.79 \pm 6.39
pH	7.27 – 7.56	-	7.18 – 7.43	-	7.28 – 7.39	-
Nitrite (mg L^{-1})	0.015 – 0.02	0.016 \pm 0.00	0.37 – 0.07	0.52 \pm 0.16	0.082 – 0.12	0.10 \pm 0.02
Nitrate (mg L^{-1})	11.68 – 18.54	15.11 \pm 3.44	33.51 – 36.95	35.23 \pm 1.72	16.51- 34.94	24.41 \pm 2.61

628 **Table 2**

629 **Environmental parameters measured** in *S. polyrhiza* culture ponds during the study period.

Parameter	Range	Mean ± SE
Temperature (°C)	30.5 - 33.0	32.00 ± 1.0
pH	7.76 - 8.30	---
Dissolved oxygen (mg L ⁻¹)	1.25 - 4.57	2.50 ± 0.25
Ammonia (mg L ⁻¹)	5.02 - 17.57	15.25 ± 0.7
Nitrite (mg L ⁻¹)	0.005 - 0.01	0.008 ± 0.002
Nitrate (mg L ⁻¹)	0.05 - 2.05	0.921 ± 0.3
Phosphate (mg L ⁻¹)	1.15 - 2.00	1.52 ± 0.07
Conductivity (µS cm ⁻¹)	1032 - 1251	1150 ± 37.0

630

631

632

633

634

635

636

637

638

639

640

641

642

643

644 **Table 3**645 Proximate composition of *S. polyrhiza* (% of dry weight).

Parameter	Manure 1 (Organic)	Manure 2 (Inorganic)
Protein	35.82 ± 0.14	30.50 ± 0.03*
Lipid	7.11 ± 0.11	7.19± 0.06
Ash	18.51 ± 0.02	20.64 ± 0.26*
Carbohydrate	38.38 ± 0.26	41.68± 0.17*

646 Data are presented as means ± SEM (n=3). *Denotes significant difference ($p < 0.05$)
647 between the two manures as determined by Student's t-test.

648

649

650

651

652

653

654

655

656

657

658

659

660

661

662

663

664

665
666
667
668
669
670
671
672
673
674
675
676
677
678
679
680
681
682
683
684
685
686
687
688
689
690
691
692
693
694
695
696
697
698
699
700
701
702

Table 4

Amino acid (g 100 g⁻¹ of dry weight) profile of *S. polyrhiza* cultured with organic manures in tanks.

Amino acids	Concentration
Essential	
Histidine (His)	0.771 ± 0.053
Isoleucine (Ile)	1.703 ± 0.150
Leucine (Lue)	3.322 ± 0.207
Lysine (Lys)	2.280 ± 0.129
Methionine (Met)	0.694 ± 0.059
Phenylalanine (Phe)	2.159 ± 0.144
Threonine (Thr)	1.502 ± 0.386
Tryptophan (Trp)	0.282 ± 0.018
Valine (Val)	2.383 ± 0.139
Non-essential	
Alanine (Ala)	2.384 ± 0.130
Arginine (Arg)	2.386 ± 0.120
Asparatate (Asp)	4.094 ± 0.212
Cysteine (Cys)	0.369 ± 0.039
Glutamic acid (Glu)	5.103 ± 0.380
Glutamine (GluNH ₂)	1.250 ± 0.300
Glycine (Gly)	2.369 ± 0.110
Proline (Pro)	1.001 ± 0.110
Serine (Ser)	1.904 ± 0.120
Tyrosine (Tyr)	1.558 ± 0.050
Free	
Phosphoserine (p-Ser)	0.060 ± 0.002
Taurine (Tau)	0.023 ± 0.006
Phospho ethanol amine (PEA)	0.072 ± 0.001
α Amino adipic acid (α-AAA)	0.020 ± 0.001
α Amino-n- butaric acid (α-ABA)	0.141 ± 0.014
Cystathionine (Cysthi)	0.115 ± 0.001
β -Alanine (β-Ala)	0.072 ± 0.011
β -Amino isobutyric acid (β-AiBA)	0.354 ± 0.015
Ethanol amine (EOHNH ₂)	0.112 ± 0.004
Ornithine (Orn)	0.027 ± 0.002
1 Methylhistidine (1 Mehis)	0.048 ± 0.003
Hydroxy proline (Hypro)	0.197 ± 0.010
γ- Amino isobutyric acid (γ-AiBA)	0.478 ± 0.024

703 **Table 5**
 704 Fatty acid composition of *S. polyrhiza* as percentage of total fatty acids (Percentage)
 705 or as mg fatty acids per 100 g dry weight (Absolute).
 706

Fatty acid	Manure 1		Manure 2	
	Percentage	Absolute	Percentage	Absolute
14:0	1.01 ± 0.22	16.9 ± 1.86	1.10 ± 0.30	23.65 ± 7.42
15:0	0.60 ± 0.04	10.1 ± 0.46	0.40 ± 0.01*	8.56 ± 0.55
16:0	31.22 ± 2.33	524.1 ± 18.32	25.50 ± 0.40	547.04 ± 33.88
18:0	2.33 ± 0.23	39.1 ± 0.35	2.02 ± 0.13	43.39 ± 4.69
20:0	0.40 ± 0.04	6.6 ± 0.10	0.33 ± 0.01	7.04 ± 0.55
22:0	0.77 ± 0.10	12.9 ± 0.32	0.85 ± 0.03	18.17 ± 0.24*
24:0	3.05 ± 0.15	51.3 ± 3.16	2.28 ± 0.05*	48.85 ± 1.15
Total saturated	39.38 ± 3.12	661.0 ± 20.21	32.48 ± 0.76	696.70 ± 48.49
16:1n-9	4.76 ± 2.23	86.4 ± 27.76	6.75 ± 0.14	144.61 ± 3.60
17:1 n	0.00 ± 0.00	0.0 ± 0.00	0.30 ± 0.02*	6.34 ± 0.22*
18:1n-9	2.09 ± 0.13	35.2 ± 1.68	3.01 ± 0.64	64.93 ± 16.74
18:1n-7	2.24 ± 0.23	37.6 ± 0.25	1.34 ± 0.06*	28.68 ± 2.59*
Total monoenes	9.09 ± 6.37	159.2 ± 124.19	11.39 ± 0.53	244.57 ± 22.72
18:2n-6	11.35 ± 0.76	190.7 ± 8.09	13.49 ± 0.23	289.08 ± 8.33*
20:4n-6	0.00 ± 0.00	0.0 ± 0.00	0.33 ± 0.01*	7.03 ± 0.08*
Total n-6 PUFA	11.35 ± 0.76	190.7 ± 8.09	13.82 ± 0.24	296.11 ± 8.41*
18:3n-3	35.75 ± 2.18	600.6 ± 29.28	38.95 ± 1.08	834.63 ± 15.44*
20:5n-3	0.38 ± 0.12	6.3 ± 1.37	0.60 ± 0.08	12.98 ± 2.30
Total n-3 PUFA	36.13 ± 0.30	606.9 ± 27.91	39.56 ± 1.00	847.61 ± 17.75*
Total DMA	4.04 ± 0.18	68.0 ± 4.35	2.76 ± 0.06*	59.09 ± 1.49
Total PUFA	47.48 ± 3.07	797.6 ± 35.99	53.37 ± 1.241	1143.72 ± 26.16*
Total Fatty acids		1685.8 ± 275.3		2144.1 ± 329.9

707
 708 Data are presented as means ± SEM (n=3). *Denotes significant difference ($p < 0.05$) between the two
 709 manures as determined by Student's t-test. DMA, dimethyl acetals; PUFA, polyunsaturated fatty
 710 acids.

711
 712
 713
 714
 715
 716
 717
 718
 719
 720
 721
 722
 723

724 **Table 6**
 725 The essential amino acid profiles of soybean (*Glycine max*) meal and *S. polyrhiza* and their
 726 requirement for *Cyprinus carpio* and *Oreochromis niloticus* (NRC, 1998, 2011).
 727

Amino acids	<i>Glycine max</i> meal (g 100 g ⁻¹)	<i>Spirodela</i> <i>polyrhiza</i> (g 100 g ⁻¹)	<i>Cyprinus</i> <i>carpio</i> (g 100 g ⁻¹ diet)	<i>Oreochromis</i> <i>niloticus</i> (g 100 g ⁻¹ diet)
Histidine (His)	1.17	0.77	0.5	1.0
Isoleucine (Ile)	1.99	1.70	1.0	1.0
Leucine (Lue)	3.42	3.32	1.4	1.9
Lysine (Lys)	2.83	2.28	2.2	1.6
Methionine (Met)	0.61	0.7	0.7	0.7
Phenylalanine (Phe)	2.18	2.15	1.3	1.1
Threonine (Thr)	1.73	1.50	1.5	1.1
Tryptophan (Trp)	0.61	0.28	0.3	0.3
Valine (Val)	2.06	2.38	1.4	1.5
Arginine (Arg)	3.23	2.38	1.7	1.2
Cysteine (Cys)	-	0.36	-	-
Tyrosine (Tyr)	-	1.55	-	-
Methionine + Cysteine	1.31	1.07	1.0	1.0
Phenylalanine + Tyrosine	-	3.7	2.0	1.6

728
 729

730
731



Fig. 1.

732




Fig. 2 (A)

733




Fig. 2 (B)

734

735

736




Fig. 2 (C)

737
738
739
740




Fig. 2 (D)

741