

1 Survival rates for *Nephrops norvegicus*
2 discarded from Northern European trawl
3 fisheries

4 Clive J Fox¹

5 Amaya Albalat²

6 Daniel Valentinsson³

7 Hans C Nilsson³ Frank

8 Armstrong⁴

9 Peter Randall⁴

10 Thomas Catchpole⁴

11

¹ Scottish Association for Marine Science, Dunstaffnage, Oban, PA37 1QA, UK

² Institute of Aquaculture, Pathfoot Building, University of Stirling, Stirling, FK9 4LA, UK

³ Department of Aquatic Resources, Institute of Marine Research, Swedish University of Agricultural Sciences, Turistgatan 5, SE-453 30 Lysekil, Sweden

⁴ Centre for Fisheries and Aquaculture Science, Pakefield Road, Lowestoft, Suffolk, NR33 OHT, UK

12 **Abstract**

13 When discarded from bottom trawl fisheries, survival of *Nephrops norvegicus* may be
14 sufficiently high that this species can be exempted from the EU Landing Obligation. In three
15 studies, *Nephrops* were sampled from trawlers in northern European waters and the fate of
16 individuals monitored for a minimum of 13 days in onshore tanks. Winter estimates of captive
17 survival (means \pm 95% confidence intervals), including immediate mortality during catch
18 sorting, were $62 \pm 2.8\%$ for the West of Scotland, $57 \pm 1.8\%$ for the Farne Deep (North Sea),
19 and $67 \pm 5.4\%$ for the Skagerrak. The Farne Deep fishery is not active in summer, but captive
20 survival rates in summer in the other two areas were reduced to $47 \pm 3.4\%$ for West of Scotland
21 and $40 \pm 4.8\%$ for the Skagerrak. Linear modelling of the West of Scotland and Skagerrak data
22 suggested that higher survivals in winter were related to colder water or air temperatures
23 although temperatures during captive observation may also have had an impact. Net
24 modifications in the Skagerrak study had an effect on survival, which was higher for *Nephrops*
25 sampled from nets equipped with the more selective Swedish sorting grid compared with Seltra
26 trawls.

27 Keywords: Discards survival; *Nephrops norvegicus*; Trawl fisheries; Landing Obligation

28 **Introduction**

29 One of the main aims of the European Union's reformed Common Fisheries Policy
30 (Regulation (EU) No. 1380/2013) is to reduce unwanted catches through a phased landing
31 obligation for regulated species, the obligation being fully implemented in 2019. Whilst
32 technical measures that allow unwanted animals to escape before being brought onto the vessel
33 are encouraged (Catchpole *et al.*, 2017), such measures rarely eliminate the unwanted
34 components of the catch completely (Broadhurst *et al.*, 2006). The landing obligation thus
35 includes exemptions and flexibility tools including for "species for which scientific evidence

36 demonstrates high survival rates, taking into account the characteristics of the gear, of the
37 fishing practices and of the ecosystem”. Producing robust estimates of post-discard survival
38 has thus become a focus for research because evidence from such studies influences whether
39 exemptions will be granted (Morfin *et al.*, 2017). Allowing continued discarding of organisms
40 with demonstrated high survivability does make conservation sense since a high proportion
41 should survive and contribute to the stock (Rihan *et al.*, 2019).

42 *Nephrops norvegicus* is a small decapod crustacean that has become increasingly important
43 in many European fisheries from Norway to the Bay of Biscay (Ungfors *et al.*, 2013). Because
44 individual *Nephrops* are encased in a strong exoskeleton, and lack gas-filled body cavities, it
45 has been suggested that this species should be a suitable candidate for the high survivability
46 exemption from the landing obligation. Most discard survival estimates have come from
47 captive observation where *Nephrops* are sampled from fishing vessels and held in captivity,
48 recording their survival over time. However, historical survival estimates from trawl fisheries
49 have been rather variable. Published rates include 17–18% (Campos *et al.*, 2015); 19% or 31%
50 depending on area (Charuau *et al.*, 1982); 42% or 75% dependent on area, trawler type and sea
51 conditions (Edwards and Bennett, 1980); 51% (Méhault *et al.*, 2016) and 56–70% (Guéguen
52 and Charuau, 1975). Some of this variation may be due to differences in fishing gears, methods
53 of handling or environmental conditions but the ICES Workshop on Methods for Estimating
54 Discard Survival (WKMEDS) suggested that variable experimental approaches might also be
55 an important factor (ICES, 2014). For example, some studies have monitored survival using
56 cages or containers placed on the seabed (Guéguen and Charuau, 1975; Campos *et al.*, 2015;
57 Méhault *et al.*, 2016), whilst other studies have monitored survival in aquaria. In some studies,
58 monitoring times may have been too short because mortality can be delayed (Wileman, 1999).
59 As one of their outputs, WKMEDS produced methodological guidance with the aim of

60 improving the robustness and reproducibility of results from discard survival experiments
61 (ICES, 2014). For captive observations, recommendations included assessing initial animal
62 condition using standardised criteria, monitoring for a sufficient time and incorporating control
63 subjects to evaluate the effect of holding conditions.

64 The main aim of the present study was to compare the survival of discarded *Nephrops* across
65 three distinct northern European trawl fisheries. Although the research followed the WKMEDS
66 recommendations, there were inevitably some methodological differences as the three studies
67 were conducted by different research groups (Valentinsson and Nilsson, 2015; Armstrong *et*
68 *al.*, 2016; Fox and Albalat, 2018). Links between survival and biological (sex, damage and
69 vitality), environmental (sea and air temperature) and operational factors (haul duration, catch
70 weights and sorting times), were also examined in order to suggest changes in trawling practice
71 that might increase the survival of discarded *Nephrops*.

72 **Methods**

73 *Operational factors*

74 Study 1 was undertaken using the MFV Ocean Trust (PD787), a 24 m stern trawler operating
75 out of Mallaig (Scottish west coast, ICES Divivision VIa, Figure 1). Fishing took place in
76 winter and summer on commercial grounds that were reasonably close to Mallaig to allow
77 experimental animals to be returned to the Scottish Association for Marine Science (SAMS)
78 aquarium, which is approximately 86 miles south by road, in reasonable time. Fishing gear
79 comprised a commercial twin-rig *Nephrops* trawl with both nets fitted for half the hauls with
80 80 mm and half the hauls with 100 mm diamond mesh cod-ends. A 200 mm square-mesh
81 escape panel (SMP) was fitted in the top-sheet of each net in accordance with local regulations,
82 but the nets did not have any further selectivity modifications (Table 1).

83 Study 2 was undertaken using the MFV Luc (SN36), an 18 m single-rig stern trawler
84 operating out of North Shields (English northeast coast, ICES Division IVb). Experimental
85 fishing took place at the southern edge of the Farne Deeps in winter only (Figure 1). The net
86 had an 80 mm diamond mesh cod-end and incorporated a NetGrid selectivity device (Table 1).
87 The NetGrid consists of a four-panel box section with a fish escape hole inserted into a standard
88 two-panel trawl with an inclined netting sheet (Armstrong *et al.*, 2016).

89 Study 3 was undertaken in winter and summer using two commercial, twin-rig stern trawlers,
90 the Canopus (LL377; 12 m) and the Ternö (LL388; 14.9 m) fishing on commercial grounds in
91 the eastern Skagerrak (Swedish southwest coast, ICES Division IIIa, Figure 1). Each vessel
92 deployed a standard Swedish grid trawl (hereafter abbreviated as SweGrid) comprising 35 mm
93 bar-spacing with a 70 mm square-mesh cod-end as described in Valentinsson and Ulmestrand
94 (2008), and a Seltra trawl with 90 mm diamond mesh cod-end and a 270 mm diamond-mesh
95 escape window as described in Krag *et al.* (2016).

96 *Environmental factors*

97 Sea surface temperatures were measured at least once a day using a Sonetek Castaway
98 (Sonetek, San Diego, CA, USA) CTD (Study 1), an Oxyguard Handy Polaris 2 (Study 2) and an
99 SD204 (SAIV A/S, Bergen, Norway) CTD (Study 3). Vertical water column profiles were only
100 recorded in studies 1 and 3. However, thermal and salinity stratification is typically minimal in
101 Feb. – Mar. at the trawl sites in study 2 (Janssen *et al.*, 1999), so surface values for these
102 parameters should have been close to those at the seabed. In all three studies, air temperatures
103 were recorded in the catch sorting area of the fishing vessel for each haul using digital
104 thermometers.

105 *Catch sorting, sampling and biological factors*

106 In all three studies, the trawler crews were asked to follow their normal fishing and catch
107 sorting practices. On all four of the fishing vessels, the catch is dropped into a flat-bottomed
108 metal hopper, from where it is raked via a hatch to a sorting table. Drop height in study 1 was
109 1.5 m, in study 2 it was less than 1 m and in study 3, 0.8–1 m. In study 1, we had to assume
110 that effects on *Nephrops* (levels of physical damage etc.) would be similar in both cod-ends as
111 the catches were not kept separate but dropped sequentially into the hopper, following the
112 normal fishing practice. In study 3, the catch from each net was kept separate by dividing the
113 hopper using wooden boards.

114 In study 1, the catch length profiles were based on measurements of at least 100 *Nephrops*
115 taken unselectively from different parts of the catch. For studies 2 and 3, only those *Nephrops*
116 selected for captive survival observation were measured on-board. For these regions, the typical
117 size ranges of *Nephrops* in the catches and discards were estimated using fisheries observer
118 data collected between 2011–2017 for ICES Division IVb, functional unit 6 and from 2015 for
119 ICES Division IIIa.

120 In all four vessels, the normal practice is that discards are returned continuously to the sea
121 throughout catch sorting via a chute at the end of the sorting table. For each haul in study 1
122 (summer), scientific staff sampled *Nephrops* being discarded from the start of catch sorting
123 until a target of 100 live animals was reached. This was subsequently modified (winter season)
124 to take a target of 100 *Nephrops* from the start, and an additional 50 towards the end of catch
125 sorting. For study 2, around 200 *Nephrops* were sampled randomly throughout catch sorting
126 across the whole size range from each haul. For study 3, observers firstly estimated the amount
127 of *Nephrops* likely to be discarded from the catches and then adjusted the rate of sampling to
128 cover the catch sorting period. The number of dead *Nephrops* encountered during sampling

129 was also recorded and used to estimate immediate mortality for each haul. In all three studies,
130 individual carapace lengths of the sampled *Nephrops* were measured using digital callipers.
131 Sex was recorded during studies 1 and 3, but not during study 2. In all three studies, each animal
132 selected for captive observation was examined for signs of visible damage (Table 2) with care
133 taken to examine both ventral and dorsal surfaces. The vitality of each animal was also assessed
134 (excepting Study 3, summer hauls). *Nephrops* sampled for captive observation were then
135 placed into individual compartments in commercial tube-sets (Figure 2). Once sampling was
136 completed, the tube-set boxes were closed using perforated lids and placed into insulated
137 containers filled with seawater. Water in the on-board holding tanks was renewed periodically
138 to ensure conditions did not deteriorate during transport to the onshore holding facilities.

139 *Transport and onshore holding*

140 For study 1, the tube-set boxes were then transported by road from Mallaig to the SAMS
141 aquarium. Cold blocks were added to the insulated containers and air supplied using a portable
142 compressor during transportation. For study 2, once the trawler had returned to port, the tubeset
143 boxes were placed directly into onshore holding tanks located at the quay. In study 3, boxes
144 were moved directly from the fishing vessels into the Kristineberg Marine Research station
145 aquarium. Oxygen levels, temperature and ammonia were checked in the transport containers
146 on arrival at the onshore holding facilities.

147 *Control animals*

148 In study 1, controls were recovered discard fraction *Nephrops* from the previous trip that
149 showed no visual injuries. Between experiments, the control animals were held in a common
150 tank containing pieces of plastic pipe to act as refuges and fed on finely chopped mussel
151 (*Mytilus edulis*) every second day. Ten control animals were added to each box of test animals
152 when the box was transferred to the onshore aquaria, except for the first hauls in each season

153 when recovered *Nephrops* were not yet available. For study 2, control animals were sourced
154 from a local creel fisher working in a different part of the Farne Deeps. These creel caught
155 controls were transferred to the quayside aquaria and held unfed for the two weeks before the
156 first treatment monitoring. The next opportunity to collect control animals was for the third of
157 three monitoring periods, during which the collection of the control and treatment animals was
158 synchronised. For study 3, control *Nephrops* were caught using creels from an area where
159 trawling is not allowed but with similar habitat, depth and environmental conditions to the
160 experimental trawl locations. The creels had a smaller than usual mesh size (20 mm) in order
161 to catch smaller *Nephrops* of sizes comparable to those normally discarded by the trawlers.
162 Control animals were added to the observation boxes on return to the quay i.e. control animals
163 were not held in captivity prior to the experiments.

164 *Observation tanks*

165 Observation tanks were supplied with running seawater at a sufficient rate for replacement at
166 least every 2 h. Seawater for the SAMS aquarium is drawn from a sub-sand beach filter and
167 incoming water temperatures can become high in summer. Observation tanks in study 1 were
168 therefore housed in a constant temperature room with additional chilling of the incoming water.
169 For study 2, temperatures in the observation tanks followed those of the ambient pumped
170 seawater because the observation tanks were located on the dockside. In study 3, seawater is
171 drawn from a deep supply. The observation tanks were housed in a constant temperature room
172 but additional water chilling was not used. Observation tanks were also aerated in studies 1 and
173 3. In study 1, temperatures in the observation tanks were monitored every 10 minutes using
174 Hobo TidbiT loggers (Onset Computer Corp., Bourne, Massachusetts) and salinity was checked
175 daily using a Castaway CTD (Sontek, San Diego, California). Dissolved oxygen (DO) was
176 monitored daily using an YSI (Yellow Springs, Ohio) Pro20 portable oxygen meter, but only

177 during the winter studies due to equipment availability. Ammonia levels were checked daily
178 using API saltwater test strips (Mars Fishcare, Chalfont, Pennsylvania). In study 2, temperature
179 and dissolved oxygen in the onshore holding tanks were measured daily using a portable meter
180 (OxyGuard Handy Polaris2) but salinity was not monitored. In study 3, temperature, salinity
181 and DO were measured daily using portable meters (WTW Multi 3510) and water samples
182 collected and analysed for ammonia.

183 *Captive observations*

184 *Nephrops* were not fed during captive observation. In study 1, *Nephrops* survival was
185 monitored every two days from 1 – 13 days post-sampling. For study 2, inspections occurred
186 daily up to 15 days, plus an additional evaluation of remaining survivors at 21 days. For study
187 3, *Nephrops* were monitored daily up to 15 days post-sampling. In all cases, the boxes were
188 lifted out of the observation tanks and the individual *Nephrops* checked in air. Exposure to air
189 was usually sufficient to cause live individuals to move but any that showed no movement were
190 gently stimulated with blunt forceps. If they still failed to react to physical stimuli, they were
191 recorded as dead and removed from the box.

192 *Statistical analyses*

193 *Nephrops* sizes are reported as carapace lengths in mm. All statistical analyses were
194 performed using R version 3.5.0 (R Core Team, 2018) with additional packages ‘boot’,
195 ‘ordinal’, ‘survival’ and ‘wrs2’. Statistical test results were considered significant at the $p < 0.05$
196 level.

197 *Statistical analyses —data collected on board the fishing vessels*

198 Exploratory analysis of sea and air temperatures, haul durations, catch weights and catch
199 sorting times by study was conducted using pairs-plots and Kendall’s tau to screen for potential
200 collinearity. *Nephrops* size data were visualised using length frequency histograms.

201 Differences in immediate mortalities within each study were explored using boxplots and tested
202 using a non-parametric two-way median test. Potential relationships between immediate
203 mortality and available covariates (sea surface and air temperatures, haul duration, catch
204 weights, catch sorting times, plus gear modification in study 3) were explored using scatterplots
205 and Kendall's tau. Immediate mortalities were then modelled as the total count of alive versus
206 dead *Nephrops* in each haul using quasi-binomial GLMs that were sequentially simplified by
207 eliminating non-significant factors, starting with the full model (Crawley, 2014). The final
208 model fits were assessed using Pearson residuals.

209 Data for physical damage at the time of sampling were summarised and ranked to identify
210 the most common injuries in each study. The mean rate of occurrence of the top five injury
211 types in each study was computed. Non-symmetrical 95% confidence intervals for these means
212 were estimated by boot-strapping as the percentage of occurrence for some injury types was
213 close to zero. Exploratory analysis of potential relationships between the percentages of injured
214 *Nephrops* and available covariates were conducted as described above for immediate mortality.
215 Analysis of the vitality scores from study 2 showed an unexplained increase in the proportions
216 in the 'Excellent' category comparing hauls on the 3rd and 4th February with later dates. To
217 standardise the vitality scores as much as possible within and across the three studies, the
218 'Excellent' category was combined with the 'Good' category to create E/G, and the 'Poor'
219 category combined with the 'Moribund' category to create P/M. This was based on the
220 argument that the criteria for separating high vitality from low vitality animals was likely to be
221 more consistent than when assigning animals to the finer divisions (Table 2), under challenging
222 field conditions. Because vitality might be related to the presence of physical injuries, chisquare
223 tests were applied to the frequencies of animals with injury presence or absence by E/G or P/M
224 categories. Data on physical injury and vitality were then combined by assigning individual

225 *Nephrops* to one of four categories: Uninjured and E/G; Uninjured and P/M; Injured and E/G;
226 Injured and P/M. Potential relationships between the percentage of *Nephrops* in each category
227 and available environmental and operational covariates (sea surface and air temperatures, haul
228 duration, catch weights, catch sorting time, plus gear for Study 3) were explored using
229 scatterplots and Kendall’s tau and modelled using ordinal regression with a logit link for each
230 study. Non-significant terms based on the Wald F-statistics were sequentially removed from
231 the regression models and the proportional odds assumption of final models tested using the
232 ‘nominal_test’ in the R ‘ordinal’ package. *Statistical analyses — data from the captive*
233 *observations*

234 Survival of control *Nephrops* was evaluated by study and the effect of season for studies 1
235 and 3 tested using Fisher’s exact test. The effect of biological factors (sex, presence of damage
236 and vitality at time of sampling) on the survival of individual *Nephrops* in the captive
237 observations was visualised using Kaplan-Meier survival curves with differences being tested
238 using log-rank tests (Moore, 2016; Kleinbaum and Klein, 2012). Because the assumption of
239 independence between each *Nephrops* within an observation box might be invalid, we firstly
240 estimated mean survivals (plus standard errors and 95% confidence intervals) for each haul
241 from the Kaplan-Meier estimator at the time of the final mortality event. These survival
242 estimates were then used to generate group mean survivals by study, season and gear. To
243 account for the uncertainty in the underlying haul-based mean survival estimates, 95%
244 confidence intervals were computed as twice the standard error incorporating propagation of
245 error following formula [1], assuming each haul-based estimate to be independent within the
246 group.

$$SE = \frac{\sqrt{\sum_i \sigma_i^2}}{n}$$

250 where n is the total number of contributing estimates and σ are the variances of each
251 contributing estimate in the group i .

252 Potential relationships between mean survival in each study and available biological
253 (percentages of *Nephrops* in each damage presence/absence, E/G or P/M group), environmental
254 (sea surface and air temperatures) and operational covariates (haul duration, catch weights,
255 sorting time, plus gear for Study 3) were explored using scatterplots and modelled using
256 multiple linear regressions with sequential removal of non-significant terms (Crawley, 2014).
257 Whilst percentage data, such as survival, infringe the limits for Gaussian error-distributions this
258 only becomes a serious issue for linear modelling if the response variable values lie close to 0
259 or 100. For a range of 30 – 70%, ordinary linear modelling can be reasonably applied (Long,
260 1997). Final model fits were assessed visually using Pearson residual plots.

261 **Results**

262 *Environmental conditions during trawling*

263 Winter air temperatures in studies 1 (West of Scotland) and 2 (North Sea) were between 6.9
264 – 11.5°C, but were colder in study 3 (Skaggerak). In summer, the air temperatures in both
265 regions reached as high as 19°C. There was also a greater seasonal difference in sea surface
266 temperatures comparing the West of Scotland with the Skagerrak. In study 1, there was little
267 thermal stratification, even in summer but this was apparent in study 3 (Table 3). In studies 1
268 and 3, near bottom salinities were around 34 but surface waters in the Skagerrak were fresher
269 with salinities of 24 – 29. Salinity was not recorded in study 2.

270 *Catches and discarding practices*

271 Based on pairs plots (Figures S1–S4), there were no obvious relationships between haul
272 duration and catch weight in any of the three studies but total catch sorting times were
273 significantly related to the *Nephrops* catch weight in study 1 (Figure S1), and to the total catch

274 weight in study 2 (Figure S2). For study 3, there did not seem to be any strong relationships
275 between total sorting times and catch weights (Figure S3 and S4). There was a noticeable
276 difference in the relative weights of *Nephrops* versus non-*Nephrops* in the catches, this being
277 much lower in study 3, where *Nephrops* comprised as little as 20 kg per net haul (Table 1). In
278 study 1, the non-*Nephrops* components of the catches were mainly spotted dogfish
279 (*Scyliorhinus canicula*), rays (Rajidae), ling (*Molva molva*), mackerel (*Scomber scombrus*),
280 various flatfish including dab (*Limanda limanda*) and juvenile gadoids such as cod (*Gadus*
281 *morhua*), hake (*Merluccius merluccius*) and haddock (*Melanogrammus aeglefinus*). In study 2,
282 the non-*Nephrops* components of the catches were mostly small gadoids. In study 3, the
283 majority of the catches were comprised of flatfishes, gadoids and other benthic invertebrates.
284 Details of the individual hauls are given in Table S1.

285 In study 1, the size range of *Nephrops* caught was 15 – 66 mm with a dominant mode at 28
286 mm and the size range of discarded *Nephrops* was 16 – 36 mm (Figure 3a). The majority of
287 discards (96%) in study 1 were larger than the Minimum Conservation Reference Size (MCRS)
288 for this fishing area. In study 2, the size range of *Nephrops* in the catch was 20 – 55 mm with
289 a dominant mode at 28 mm (Figure 3b). This size range also closely matches that recorded over
290 seven years by fisheries observers on English trawlers fishing in the Farne Deeps. Observer
291 data for ICES Division IVb showed that similar sizes of *Nephrops* are typically discarded as in
292 study 1 but, because the MCRS is larger in Division IVb, a smaller percentage (54%) of these
293 discarded *Nephrops* were above the MCRS (Figure 3b). In study 3, fisheries observer data for
294 2015 showed that *Nephrops* in trawl catches from this area ranged from 20 – 69 mm. Discarded
295 *Nephrops* in study 3 ranged from 20 – 58 mm with a minority (8%) being above MCRS (Figure
296 3c). Compared with the other areas this reflects the larger MCRS in ICES Division IIIa at the
297 time (Hornborg *et al.*, 2017). Thus, in all three study areas *Nephrops* were being discarded for

298 reasons other than the animals being below the minimum legal size, this being a particularly
299 prominent feature in studies 1 (ICES Division VIa) and 2 (ICES Division IVb).

300 *Immediate mortality*

301 In study 1 in winter, the mean immediate mortality (\pm 95% lower confidence level (LCL),
302 upper confidence level (UCL)) was 9.7% (7.8, 11.9) and in summer, it was 14.5% (11.9, 19.7).
303 However, because of variability in the immediate mortalities, neither season nor cod-end mesh
304 size were statistically significant (Figure 4; med2way test: Season $p=0.13$, Cod-end $p=0.51$).
305 Plotting immediate mortality by haul against available covariates (Figure S5) suggested that
306 immediate mortality might be related to total catch weight, *Nephrops* catch weight, sorting time
307 and air temperature with a possible effect of sea surface temperature. However, sequential
308 removal of least significant terms in the GLM resulted in retention of sorting time alone (Table
309 4), although this factor was itself correlated with *Nephrops* catch weight (Figure S1). In study
310 2, no immediate mortality was observed. In study 3 in winter, mean immediate mortality (\pm
311 95% LCL, UCL) was 1.6% (0, 3.2) but in summer increased to 14.6% (11.6, 17.8). Median
312 immediate mortality was significantly related to season, but not to gear (Figure 4; med2way
313 test: Season $p<0.001$, Gear modification $p=0.08$). Scatterplots for study 3 (Figure S6) suggested
314 that immediate mortality might be related to sea surface temperature and this term was retained
315 in the final GLM (Table 4). Residual plots for the GLM models for studies 1 and 3 indicated
316 reasonable fits.

317 *Injury and vitality during catch sorting*

318 The percentage of discarded *Nephrops* with at least one visible injury ranged between 23 –
319 67% of the animals examined from each haul. The most common injuries were loss or damage
320 to one or both chelae, puncture and crush wounds to the thorax or abdomen and damaged rostra

321 (Table 5). Damage to one or more legs, the telson or the eye occurred in less than 1% of the
322 *Nephrops* examined. Scatterplots of the percentage of damaged *Nephrops* against available
323 covariates failed to reveal significant relationships, except in study 1 with non-*Nephrops* catch
324 weight and in study 3 with sea surface temperature (Figures S7, S8). For vitality, the percentage
325 in the E/G category in each haul was related to sea surface temperature in studies 1 and 2, and
326 to haul duration in the winter hauls of study 3 (Figures S9, S10). In all three studies, the
327 presence of at least one physical injury tended to reduce the vitality score of individual
328 *Nephrops* (Study 1: Chisq = 107, df=1, p<0.001; Study 2: Chisq = 228, df=1, p<0.001; Study
329 3: Chisq = 13, df=1, p<0.001) justifying combining the presence of at least one physical injury
330 with vitality. However, ordinal regression of *Nephrops* assigned to these combined injury plus
331 vitality categories failed to identify any significant environmental or operational covariates.

332 *Conditions on-board and during transport*

333 In study 1, the time elapsed between sampling and transfer of tube-boxes into the observation
334 tanks varied from 3 – 9 h with the road transport normally taking around 2 h. Oxygen levels on
335 arrival at the aquarium were between 7.8 – 8.8 mg l⁻¹. Ammonia levels were elevated but not
336 above 1 mg l⁻¹. In study 2, *Nephrops* were held in on-board tanks on the fishing vessel for 2.5
337 – 5.5 hours, oxygen saturation remained above 90% and the animals were then transferred
338 directly to the quayside facility. In study 3, time elapsed between sampling aboard and the
339 transfer of the boxes into the observation tanks varied from 2 – 4 h. Oxygen saturation was
340 always above 90%, and ammonia levels never exceeded 0.15 mg l⁻¹.

341 *Conditions in the captive observation tanks*

342 In study 1, the mean water temperature in the observations tanks was 7.6°C in winter
343 fluctuating by less than 1°C (Table 3). In summer, the mean water temperature was 9.4°C but
344 with larger fluctuations when the chillers struggled to cope with high temperatures of the

345 incoming seawater. However, temperatures did not exceed those measured at the trawling sites
346 during summer (Table 3). Salinities in the observation tanks were slightly lower than equivalent
347 bottom salinities at the trawling sites, reflecting the location of the SAMS aquarium seawater
348 intake. Dissolved oxygen was always above 8 mg l⁻¹ and ammonia levels were usually
349 undetectable but peaked at 1 mg l⁻¹ on a single occasion when the water flow to one recovery
350 tank became temporarily reduced. In study 2, water was drawn directly from the quayside and
351 the tanks were not under temperature control. Nevertheless, as this study was only conducted
352 in the winter, temperatures were generally close to the sea surface temperatures measured at
353 the haul locations (Table 3). In study 3, observation tank temperatures only fluctuated by 1°C,
354 averaging 5.5°C in winter and 14.5°C in summer. However, in summer water temperatures were
355 up to 5°C warmer than the bottom temperatures measured at the haul sites. Salinities were close
356 to those measured at the haul sites (Table 3). Oxygen levels remained above 80% saturation
357 throughout all experiments and ammonia levels were barely detectable.

358 *Size and survival of control animals*

359 The size (mean ± stdev.) of control *Nephrops* in study 1 was 25 ± 2.2 versus 24 ± 2.4 mm in
360 the test animals. In study 2, the relative sizes were 40 ± 3.8 mm and 32 ± 6.4 respectively, and
361 in study 3, 38 ± 2.7 mm and 38 ± 4.6 respectively. Survival for controls during the monitoring
362 of captive *Nephrops* was 96% in study 1 (n=170), 94% in study 2 (n=214) and 97% in study 3
363 (n=390). For studies 1 and 3, seasonal differences in control survival were not statistically
364 significant (Fisher's exact tests; p>0.05). This was not tested for study 2, which took place only
365 in winter.

366 *Factors affecting survival of individual Nephrops*

367 Based on Kaplan-Meier curves and log-rank tests, sex was not a significant factor affecting
368 individual survival in either study 1 or 3 (Figure S11, Log-rank tests: study 1, Chisq=1.0, p=0.3;

369 study 3 Chisq=1.9, p=0.2). Sex was not recorded in study 2. The presence of physical injuries
370 affected individual survival with puncture and crush injuries having the greatest negative
371 impacts (Figure S12). Kaplan-Meier curves (Figure 5) showed significant effects on individual
372 survival for the *Nephrops* in the four presence of injury combined with vitality categories
373 (Logrank tests df = 3: Study 1, Chisq = 299, p<0.001; Study 2, Chisq = 610, p<0.001; Study 3,
374 Chisq = 126, p<0.001). In all three studies, survival of undamaged animals in excellent or good
375 vitality was significantly higher than for injured *Nephrops* in a poor or moribund state at the
376 time of sampling.

377 *Survival estimates*

378 Based on the overall survival curves (Figure S13), about 90% of the observed mortalities had
379 occurred by 8 days and further mortalities had largely ceased by 10 days of observation. Final
380 survival estimates by haul including immediate mortality are given in Table S2, illustrated in
381 Figure S14, and presented grouped by study, season and fishing gear in Table 6 and Figure S15.

382 *Final survival, biological, environmental and operational factors*

383 In study 1 (Scottish west coast), final survival estimates were significantly higher in winter
384 than summer (winter $62 \pm 2.8\%$ versus summer $47 \pm 3.4\%$; ANOVA: Survival ~ Season,
385 Season F=13.0, df=1, P=0.002). In this study, *Nephrops* were only sampled from the start of
386 the catch sorting for the summer hauls and this could have resulted in some over-estimation of
387 survival. The approach was subsequently changed so that the entire catch sorting time was
388 sampled for the winter hauls. In study 2 (Farne Deeps, North Sea), mean final survival was 57
389 $\pm 1.8\%$ but only assessed in winter. In study 3 (Skagerrak), final survival was again higher in
390 winter than in summer (winter $67 \pm 5.4\%$ versus summer $40 \pm 4.8\%$). However, gear also had
391 an effect with final mean survival being higher for *Nephrops* caught using trawls fitted with the
392 Swedish grid (ANOVA: Survival ~ Season*Gear modification, Season F=67.5, df=1,

393 $P < 0.001$, Gear modification $F = 9.71$, $df = 1$, $P = 0.01$, Gear modification by Season, $P > 0.05$.
394 Scatterplots and Kendall's tau suggested that the seasonal effect in study 1 might be linked to
395 differences in air temperature but catch sorting time were also significantly correlated with
396 survival (Figure S16). In the simplified multiple linear regression model of survival, sea surface
397 temperature, and not air temperature, along with catch sorting time were retained (Table 7).
398 Seasonal effects were not tested for in study 2 as this was conducted in winter only. The weight
399 of non-*Nephrops* catch was just significantly correlated with final survival, but in a positive
400 manner (Figure S16). Multiple regression simplification failed to identify any significant
401 predictors of survival for study 2. Although neither sea nor air temperature were selected as
402 significant, the temperature range in study 2 was limited since all hauls were conducted in
403 winter. For study 3, although there were apparent effects of sea surface and air temperature on
404 final survival by gear, the correlations were not statistically significant (Table S17). Sea surface
405 temperature was however retained in the simplified multiple linear regression models of
406 survival for study 3 (Table 7). Final survival results across all three studies did appear consistent
407 with an overall temperature effect (Figure 6). However, because sea surface and air
408 temperatures were correlated it is not possible to say with any certainty which factor was having
409 the stronger impact. In relation to biological factors, scatterplots and Kendall's tau failed to
410 identify any patterns of mean survival for each haul with the proportions of *Nephrops* in the
411 injury presence/absence combined with E/G or P/M vitality groups (Figures S18, S19).

412 **Discussion**

413 *Factors affecting immediate Nephrops mortality*

414 Being caught in trawls results in a range of physiological and physical responses in *Nephrops*.
415 Animals will exhibit vigorous tail flipping as they try to escape from the ground gear (Newland
416 and Chapman, 1989) and such activity results in depletion of muscle ATP and increased levels

417 of anaerobic metabolites (Albalat *et al.*, 2009). Exposure of *Nephrops* to low salinity surface
418 waters during net hauling may lead to further physiological stress, but this is only likely to be
419 important in strongly salinity-stratified waters such as the Kattegat and Skagerrak (Harris and
420 Ulmestrand, 2004). Although haloclines were present in our third study in the Skagerrak, the
421 surface salinities were not as low as the salinity of 15 used in the laboratory experiments
422 conducted by Harris and Ulmestrand (2004). Once on board fishing vessels, *Nephrops* are
423 usually held in air during catch sorting resulting in multiple physiological and immunological
424 changes associated with oxygen deprivation (Spicer *et al.*, 1990; Albalat *et al.*, 2009; Lund *et*
425 *al.*, 2009; Campos *et al.*, 2015). These changes are potentially reversible when *Nephrops* are
426 returned to seawater, but the temperature of the aerial exposure appeared to influence
427 immediate mortality in study 3. In study 1, immediate mortality was related to total catch
428 sorting time, possibly because of prolonged aerial exposure of *Nephrops* in the hopper.

429 *Factors affecting final survival*

430 Consistent with previous studies (Symonds and Simpson, 1971; Wileman *et al.*, 1999;
431 Campos *et al.*, 2015; Albalat *et al.*, 2016), we found clear links between the survival of
432 individual *Nephrops* during captive observation and the presence of physical damage plus
433 vitality at the time of sampling. Puncture and crush injuries in particular are known to lead to
434 loss of haemolymph often resulting in eventual circulatory collapse (Wileman *et al.*, 1999).
435 However, despite the clear link between physical damage plus vitality and survival at the
436 individual level, only sea surface temperature consistently emerged as a significant predictor
437 of final mean survival in studies 1 and 3. However, because sea surface and air temperatures
438 were correlated, it was difficult to determine which factor was having more impact. Although
439 several studies have highlighted the negative link between increased air temperatures and
440 *Nephrops* survival (Spicer *et al.*, 1990; Ridgway *et al.*, 2006), being returned to warmer water

441 in summer, either by being discarded at sea or when placed into observation tanks, might also
442 reduce survival. Since metabolic costs are linked to temperature, elevated energetic costs might
443 reduce an animal's capacity for recovery during summer months. In addition, bacterial and
444 fungal growth rates are likely to be higher in summer, perhaps resulting in poorer survival of
445 injured *Nephrops* recovering in warmer water. Broadhurst *et al.* (2006) suggested that simple
446 measures to keep catches cool, such as ensuring hopper covers are closed after the nets have
447 been emptied or installing chillers, might improve discard survival. However, such measures
448 could be less beneficial in summer if lower survival rates are also due to animals recovering in
449 warmer water. This could be tested in further captive observation trials if the water
450 temperatures in the recovery tanks were kept constant between seasons.

451 *Experimental design*

452 The conclusion that final survivals were linked to temperature must be treated with some
453 caution because most of the hauls at the higher temperatures were from the third study. There
454 is thus scope for inter-study effects to have contributed to the overall relationship. This problem
455 could be overcome by randomly allocating hauls across the full range of covariates but this is
456 difficult to achieve in field-based studies where the activity, in this case trawling and its
457 associated environmental conditions, are not under direct experimental control. Furthermore,
458 temperatures in the observation tanks did not always coincide with those measured in the field.
459 In particular, temperatures were colder in the observation tanks for study 1 summer hauls but
460 warmer in study 3 summer captive observations. We are not aware of any discard recovery
461 studies with *Nephrops* where the effects of different water temperatures during recovery have
462 been investigated, but water temperatures in the observation tanks could have had some impact
463 on the results.

464 The considered opinion of WKMEDS (ICES, 2014) is that to date, there are no satisfactory
465 methods for adjusting discard-survival estimates using control data. Therefore, it is currently
466 recommended that the magnitude of the control mortality should be used as a measure of the
467 validity of the observation method, where control mortalities close to zero suggest a more valid
468 method for accurately estimating discard survival. In the present studies, mortality of control
469 animals was less than 5% suggesting that the observational setups were not causing high levels
470 of stress. It must be noted that control animals were added to the observation boxes when they
471 reached the aquaria. Adding control animals to the observation boxes on-board the trawlers
472 was impractical because the control *Nephrops* would have had to be transported back to the
473 haul locations, in some cases on the previous evening, and thus exposed to even more
474 unrealistic stressors. However, any mortality in the test subjects resulting from being placed
475 into insulated containers on board the trawlers and transported to the aquaria could not be
476 identified with the approach used. Sourcing appropriate control animals for discard survival
477 studies is also challenging (ICES, 2014; Campos *et al.*, 2015; Méhault *et al.*, 2016; Morfin *et*
478 *al.*, 2017; Mérillet *et al.*, 2018). Although previous studies have also used recovered (Mérillet
479 *et al.*, 2018) or creel-caught *Nephrops* (Wileman *et. al.*, 1999), both approaches are open to
480 challenge. The use of recovered animals might not represent the full health and robustness
481 range of *Nephrops* caught in the trawls since recovered animals might be those more resilient
482 to such stresses. However, this approach did ensure that control animals are of similar size to
483 those being discarded. For creel-caught controls, their larger size compared with those being
484 discarded may be an issue but this potential problem was minimised in study 3 by using creels
485 with a reduced mesh size.

486 Studies 1 and 2 were based on single vessels whilst study 3 used two vessels. Any
487 extrapolation of results must be made cautiously because of the variety of operations in the

488 wider fishing fleet. Given the logistical challenges and costs of conducting discard survival
489 experiments across multiple fishing vessels, relationships between survival during captive
490 observation and vitality have been used to extrapolate captive observation findings to a larger
491 number of vessels (Morfin *et al.*, 2017). However, this approach relies on the assumption that
492 survival depends only on vitality plus any environmental covariates identified as statistically
493 affecting captive survival. In the present studies, mean survival by haul did not appear to be
494 strongly linked to such factors, suggesting that a substantial part of the variability in survival
495 is being driven by additional, un-measured factors.

496 *Limitations with captive observation survival estimates*

497 Several publications have pointed out that tank-based discard survival experiments are likely
498 to over-estimate true survival by ignoring predation mortality that may occur at the sea surface,
499 in the water column or when discarded animals reach the seabed (Symonds and Simpson, 1971;
500 Raby *et al.*, 2014, Morfin *et al.*, 2017; Mérillet *et al.*, 2018). Although seabirds probably do not
501 take a large proportion of discarded *Nephrops* (Catchpole *et al.*, 2006; Depestele *et al.*, 2016),
502 this predation risk can be minimized by releasing discards below the sea surface using a
503 protective chute. Little work has been undertaken on predation of discarded *Nephrops* during
504 their descent through the water column but Bergmann *et al.* (2002) suggested that discards
505 would reach the seabed in a few minutes. As far as we are aware, there are no estimates of
506 predation rates of live, discarded *Nephrops* once they reach the seabed although the behaviour
507 of small *Nephrops* released at depths of around 100 m has been observed using a remotely
508 operated vehicle (Fox and Albalat, 2018). It was reported that undamaged *Nephrops*, even after
509 aerial exposure for up to 3 h, recovered rapidly and began exploring their environment and
510 entering available burrows within 10 min. However, these observations were only made on a
511 limited number of dives and the *Nephrops* could only be followed for a short time. The

512 conclusions reached might not apply to grounds with higher abundances of predators, to
513 damaged *Nephrops* or to those previously exposed to prolonged elevated air temperatures.

514 Furthermore, if *Nephrops* are discarded over un-suitable habitat, for example whilst steaming
515 back to port, they will have no chance of finding suitable protection in burrows (Evans *et al.*,
516 1994).

517 The longer-term effects of discarding on *Nephrops* are also difficult to assess. Evans *et al.*
518 (1994) demonstrated that animals lacking one chela were less successful in competing for food
519 and shelter compared with un-injured *Nephrops*. In the present studies, this injury was seen in
520 around 20% of the discards and these animals may be at a competitive disadvantage when
521 returned to the sea. Reducing the occurrence of such injuries should improve survival potential
522 but is challenging as levels of physical damage are related to animal condition, gear type, haul
523 duration, seabed condition, size of catches and composition, catch handling and hopper design
524 (Campos *et al.*, 2015; Méhault *et al.*, 2016). Oliver *et al.* (2017) suggested that trawls that are
525 more selective will result in less physical damage to *Nephrops* in the net, thus potentially
526 increasing discard survival. However, across all three studies we were unable to establish a
527 statistical link between the proportions of *Nephrops* with physical damage and final survival
528 by haul, even though such injuries led to reduced survival at the individual level. Within study
529 3 (Skagerrak), there was an effect of gear with captive survival of discarded *Nephrops* from
530 nets equipped with Swedish grids being higher. Swedish grid trawls are considered more
531 selective than Seltra trawls (Madsen and Valentinsson, 2010). Unfortunately, vitality was only
532 recorded on the winter hauls making it difficult to reach firmer conclusions regarding the
533 interplay of gear selectivity, *Nephrops* condition and subsequent survival.

534 *Comparison with other published studies*

535 Levels of immediate mortality recorded in studies 1 and 3 were quite similar to the 15.6%
536 immediate mortality reported by Mérillet *et al.* (2018) when using a discard chute. In study 2,
537 no immediate mortality was observed. This was unexpected and not explained by any obvious
538 differences between the studies, such as tow lengths or catch weights (Table 1).

539 There are a limited number of published *Nephrops* survival studies undertaken at different
540 seasons but Mérillet *et al.* (2018) reported higher survival in summer (57%) compared with
541 spring (42%). This contrasts with findings of reduced final survival at higher temperatures in
542 the present studies, and with other publications reporting a negative link between *Nephrops*
543 survival and temperature (Méhault *et al.*, 2016). Mean summer survival estimates in study 1
544 (47%) and study 3 (40%) were lower than a recent result of 64% reported by Oliver *et al.* (2017)
545 off the west coast of Ireland using Seltra trawls and of 57% reported by Mérillet *et al.* (2018)
546 for the Bay of Biscay using a modified discarding chute. This may reflect genuine differences
547 between the fisheries because the experimental methodology across all these recent studies
548 largely followed the WKMEDS guidelines. *Conclusions and recommendations for future work*

549 Despite some operational differences between the three studies, the final survival estimates
550 were reasonably consistent. In all three winter studies, over half the observed *Nephrops*
551 survived a minimum 13 days captive observation whilst in the two studies conducted in
552 summer, survival was between a third and half. Although what constitutes “high-survivability”
553 is not defined in the Landing Obligation (Regulation (EU) No. 1380/2013), the results
554 presented in the present paper have been reviewed by the Scientific, Technical and Economic
555 Committee for Fisheries (STECF) and accepted by the European Commission as the basis for
556 exemptions in the North Sea and west of Scotland.

557 In the two studies conducted across seasons, final survival of discarded *Nephrops* was
558 significantly higher in winter than summer. Sea surface temperature was identified as affecting
559 both immediate mortality and final survival in study 3, but only final survival in study 1.
560 However, the effect of air temperature was only marginally weaker in the models making it
561 hard to conclude which of these two correlated environmental factors might be driving the
562 seasonal response. Altering fishing practices to keep catches cool during catch sorting may thus
563 improve discard survival, particularly in summer. However, poorer captive observation
564 survival in summer could also be related to animals recovering in warmer water, in which case
565 chilling during catch sorting may have less positive effect. Further studies where water
566 temperatures in the captive observation tanks are manipulated could be undertaken to test this.
567 Our results also confirmed that physical damage to *Nephrops* significantly reduces their
568 survival potential with puncture and crush injuries being most deleterious. Although we were
569 unable to link statistically the overall levels of damage within hauls to resultant mean final
570 survival, better survival was observed from catches made with trawls equipped with a Swedish
571 sorting grid compared to a Seltra trawl. Weights for the non-*Nephrops* component of the
572 catches were lower in hauls made with the Swedish grid. Furthermore in study 1, immediate
573 mortality was lower in lighter hauls where the overall sorting times were also lower. Recording
574 levels of physical damage and vitality of *Nephrops* when gear selectivity studies are conducted
575 in future could provide valuable additional data.

576 Once on board, catch-handling practices that may lead to further damage should be avoided.
577 There is potential that changes in hopper design, such as sloping floors allowing the catch to
578 be pulled onto the sorting tables with the assistance of gravity (Albalat *et al.*, 2016), or seawater
579 hoppers (Broadhurst *et al.*, 2006), might be beneficial in reducing damage and improving
580 discard survival. Although we are not aware of any research into this in northern European

581 *Nephrops* fisheries, the benefits of seawater hoppers for improving discard survival have
582 received attention in Australian prawn fisheries (Ocean Watch Australia, 2004). However, it
583 must be cautioned that placing catches into hoppers filled with low salinity seawater may cause
584 additional stress, such measures may therefore not be effective in improving survival in areas
585 with reduced surface salinity, such as the Skagerrak. Similar considerations would apply during
586 summer months if un-chilled seawater hoppers were filled with warm surface seawater.

587 Given that discard survival studies are expensive to conduct (Morfin *et al.*, 2017), it is
588 recognised that future studies need to be standardised as much as possible (ICES, 2014).
589 Despite efforts at standardisation, some differences were apparent between the three studies
590 reported here. For example, physical water column parameters were not measured in a
591 consistent manner, analysis of the vitality data raised some doubts about the consistency of
592 scoring and temperatures in the observation tanks did not always reflect those in the field. Such
593 problems need to be tackled through further training and inter-calibration between laboratories
594 conducting discard survival studies.

595 **Acknowledgements**

596 The authors would like to extend their sincere thanks to the skippers and crews of the
597 participating fishing vessels. The authors would also like to acknowledge the anonymous
598 reviewers whose comments have helped shape and improve the manuscript. Studies in Division
599 VIa were funded by a grant (FIS015) from Fisheries Innovation Scotland; in Division IVb by
600 UK Defra program ASSIST MF1232 and in Division IIIa by the Swedish Agency for Marine
601 and Water Management (grant id. 1861-2019).

602 603 **References**

604

605 Albalat, A., Collard, A., Bruce, M., Coates, C. J., and Fox, C. J. 2016. Physiological condition,
606 short-term survival, and predator avoidance behavior of discarded Norway lobsters (*Nephrops*
607 *norvegicus*). *Journal of Shellfish Research*, 35: 1053–1065.

608

609 Albalat, A., Gornik, S. G., Atkinson, R. J. A., Coombs, G. H., and Neil, D. M. 2009. Effect of
610 capture method on the physiology and nucleotide breakdown products in the Norway lobster
611 (*Nephrops norvegicus*). *Marine Biology Research*, 5: 441–450.

612

613 Armstrong, F., Randall, P., Ribeiro, A., Jones, P., Firmin, C., Doran, S., and Catchpole, T. L.
614 2016. Assessing the survival of discarded *Nephrops* in the English NE *Nephrops* selective trawl
615 fishery. Project report ASSIST MF1232. Centre for Environment, Fisheries and Aquaculture
616 Science, Lowestoft, Suffolk. 29 pp.

617

618 Bergmann, M., Wiczorek, S. K., Moore, P. G., and Atkinson, R. J. A. 2002. Utilisation of
619 invertebrates discarded from the *Nephrops* fishery by variously selective benthic scavengers in
620 the west of Scotland. *Marine Ecology Progress Series*, 233: 185–198.

621

622 Broadhurst, M. K., Suuronen, P., and Hulme, A. 2006. Estimating collateral mortality from
623 hauled fishing gear. *Fish and Fisheries*, 7, 180–218.

624

625 Campos, A., Fonseca, P., Pilar-Fonseca, T., Leocádio, A. M., and Castro, M. 2015. Survival of
626 trawl-caught Norway lobster (*Nephrops norvegicus* L.) after capture and release - Potential
627 effect of codend mesh type on survival. *Fisheries Research*, 172: 415–422.

628

629 Catchpole, T. L., Frid, C. L. J., and Gray, T. S. 2006. Importance of discards from the English
630 *Nephrops norvegicus* fishery in the North Sea to marine scavengers. Marine Ecology
631 Progress Series, 313: 215-226.

632

633 Catchpole, T. L., Ribeiro-Santos, A., Mangi, S. C., Hedley, C., Gray, T. S. 2017. The challenges
634 of the landing obligation in EU fisheries. Marine Policy 82: 76–86.

635

636 Charuau, A., Morizur, Y., and Rivoalen, J. J. 1982. Survie des rejets de *Nephrops norvegicus*
637 dans le Golfe de Gascogne et en mer Celtique, ICES C.M. 1982/B:13, 6 pp.

638

639 Crawley, M. J. (2014) The R book. 2nd edition, John Wiley & Sons Ltd., Chichester, 1051 pp.

640

641 Depestele, J., Rochet, M.-J., Dorémus, G., Laffargue, P., and Stienen, E. W. M. 2016. Favorites
642 and leftovers on the menu of scavenging seabirds: modelling spatiotemporal variation in
643 discard consumption. Canadian Journal of Fisheries and Aquatic Science, 73: 1446–1459.

644

645 Edwards, E. S., and Bennett, D. B. 1980. Survival of discarded *Nephrops*. ICES CM
646 1980/K:10, 6 pp.

647 Evans, S. M., Hunter, J. E., Elizal, and Wahju, R. I. 1994. Composition and fate of the catch
648 and bycatch in the Farne Deep (North Sea) *Nephrops* fishery. ICES Journal of Marine Science:
649 51: 155–168.

650

651 Fox, C. J., and Albalat, A. 2018. FIS015 - Post-catch survivability of discarded Norway lobsters
652 (*Nephrops norvegicus*): Further investigations within the large-scale fleet operation, Project
653 report for Fisheries Innovation Scotland, 219 pp.

654

655 Guéguen, J., and Charuau, A. 1975. Essai de détermination du taux de survie des langoustines
656 hors taille rejetées lors des opérations de pêche commerciale, ICES CM 1975/K:12, 3 pp.

657

658 Harris, R. R., and Ulmestrand, M. 2004. Discarding Norway lobster (*Nephrops norvegicus* L.)
659 through low salinity layers – mortality and damage seen in simulation experiments. ICES
660 Journal of Marine Science, 61: 127–139.

661

662 Hornborg, S., Jonsson, P., Sköld, M., Ulmestrand, M., Valentinsson, D., Ritzau Eigaard, O.,
663 Feekings, J., Nielsen, J. R., Bastardie, F., and Lövgren, J. 2017. New policies may call for new
664 approaches: the case of the Swedish Norway lobster (*Nephrops norvegicus*) fisheries in the
665 Kattegat and Skagerrak. ICES Journal of Marine Science 74: 134–145.

666

667 ICES. 2014. Report of the ICES Workshop on Methods for Estimating Discard Survival
668 (WKMEDS). CM 2014/ACOM:51, 114 pp.

669 Janssen, F., Schrum, C., Backhaus, J.O. 1999. A climatological data set of temperature and
670 salinity for the Baltic Sea and the North Sea. Deutsche Hydrographische Zeitschrift Supplement
671 9, 245 pp.

672

673 Kleinbaum, D.G., and Klein, M. (2012) Survival Analysis. 3rd edition, Springer, New York,
674 700 pp.

675

676 Krag, L. A., Herrmann, B., Feekings, J., and Karlsen, J. D. 2016. Escape panels in trawls – a
677 consistent management tool? Aquatic Living Resources, 29: 306.

678

679 Long, J.S. (1997). Regression Models for Categorical and Limited Dependent Variables. Sage
680 Publishing. 328 pp.

681

682 Lund, H. S., Wang, T., Chang, E. S., Pedersen, L. F., Taylor, E. W., Pedersen, P. B., and
683 McKenzie, D. J. 2009. Recovery by the Norway lobster *Nephrops norvegicus* (L.) from the
684 physiological stresses of trawling: Influence of season and live-storage position. Journal of
685 Experimental Marine Biology and Ecology, 373: 124–132.

686

687 Madsen, N., and Valentinsson, D. 2010. Use of selective devices in trawls to support recovery
688 of the Kattegat cod: a review of experiments and experience. ICES Journal of Marine Science
689 67(9): 2042-2050.

690

691 Méhault, S., Morandeau, F., and Kopp, D. 2016. Survival of discarded *Nephrops norvegicus*
692 after trawling in the Bay of Biscay. Fisheries Research, 183: 396–400.

693

694 Mérillet, L., Méhault, S., Rimaud, T., Piton, C., Morandeau, F., Morfin, M., and Kopp, D.

695 2018. Survivability of discarded Norway lobster in the bottom trawl fishery of the Bay of
696 Biscay. *Fisheries Research*, 198: 24–30.

697

698 Moore, D. F. 2016. *Applied survival analysis using R*. Springer International Publishing,
699 Switzerland, 226 pp.

700

701 Morfin, M., Kopp, D., Benoît, H. P., Méhault, S., Randall, P., Foster, R., and Catchpole, T.
702 2017. Survival of European plaice discarded from coastal otter trawl fisheries in the English
703 Channel. *Journal of Environmental Management*, 204: 404-412.

704

705 Newland, P. L., and Chapman, C. J. 1989. The swimming and orientation behaviour of the
706 Norway lobster, *Nephrops norvegicus* (L.), in relation to trawling. *Fisheries Research*, 8: 63–
707 80.

708

709 Ocean Watch Australia. 2004. *Hoppers in Australian prawn fisheries - A handbook for fishers*.
710 Ocean Watch Australia Pty Ltd., Pyrmont, New South Wales, 48 pp.

711

712 Oliver, M., McHugh, M., Browne, D., Murphy, S., and Cosgrove, R. 2017. *Nephrops*
713 *survivability in the Irish demersal prawn fishery*, BIM, New Docks, Galway, 14 pp.

714

715 Raby, G. D., Packer, J. R., Danylchuk, A. J., Cooke, S. J. 2014. The understudied and
716 underappreciated role of predation in the mortality of fish released from fishing gears. *Fish and*
717 *Fisheries*, 15: 489-505.

718

719 R Core Team (2018). R: A language and environment for statistical computing. R Foundation
720 for Statistical Computing, Vienna, Austria.

721

722 Ridgway, I. D., Taylor, A. C., Atkinson, R. J. A., Stentiford, G. D., Chang, E. S., Chang, S. A.,
723 and Neil, D. M. 2006. Morbidity and mortality in Norway lobsters, *Nephrops norvegicus*:
724 physiological, immunological and pathological effects of aerial exposure. Journal of
725 Experimental Marine Biology and Ecology, 328: 251–264.

726

727 Rihan, D., Uhlmann, S.S., Ulrich, C., Breen, M., and Catchpole, T., 2019. Requirements for
728 documentation, data collection and scientific evaluations. In: The European Landing
729 Obligation: Reducing Discards in Complex, Multi-Species and Multi-Jurisdictional Fisheries.
730 S.S. Uhlmann, C. Ulrich, S.J. Kennelly (eds) Cham: Springer International Publishing, pp. 49–
731 68.

732

733 Spicer, J. I., Hill, A. D., Taylor, A. C., and Strang, R. H. C. 1990. Effect of aerial exposure on
734 concentrations of selected metabolites in blood of the Norwegian lobster *Nephrops norvegicus*
735 (Crustacea: Nephropidae). Marine Biology, 105: 129–135.

736

737 Symonds, D. J., and Simpson, A. C. 1971. The survival of small *Nephrops* returned to the sea
738 during commercial fishing. ICES Journal du Conseil, 34: 89–97.

739

740 Ungfors, A., Bell, E., Johnson, M. L., Cowing, D., Dobson, N. C., Bublitz, R., and Sandell, J.

741 2013. *Nephrops* Fisheries in European Waters. In *Advances in Marine Biology*, 64: 247–314.

742

743 Valentinsson, D., and Ulmestrand, M., 2008. Species-selective *Nephrops* trawling: Swedish
744 grid experiments. *Fisheries Research*, 90: 109–117.

745

746 Valentinsson, D., and Nilsson, H. C. 2015. Effects of gear and season on discard survivability
747 in three Swedish fisheries for Norway lobster (*Nephrops norvegicus*). Swedish University of
748 Agricultural Sciences. 11 pp.

749 <https://www.slu.se/globalassets/ew/org/inst/aqua/externwebb/radgivning/radgivning->

750 [omfiskemojligheter-och-kvoter/nephrops-discard-survival_2_v2.pdf](https://www.slu.se/globalassets/ew/org/inst/aqua/externwebb/radgivning/radgivning-omfiskemojligheter-och-kvoter/nephrops-discard-survival_2_v2.pdf)

751

752 Wileman, D. A., Sangster, G. I., Breen, M., Ulmestrand, M., Soldal, A. V., and Harris, R. R.

753 1999. Roundfish and *Nephrops* survival after escape from commercial gear. Final report, EC

754 Contract No: FAIR-CT95-0753. 240 pp.

Table 1: Summary of study locations, season (W — winter; S — summer), fishing gears, number of hauls, dates, tow durations and catch weights (mean \pm std dev.) and the biological factors recorded on board the fishing vessels (Y — yes; N — no): CL — Catch length profile; DL — Discard length profile; DS — Discard sex profile; DM — Discard immediate mortality; DP — Discard physical damage; DV — Discard vitality.

Study	ICES Div.	Season	Cod-end mesh (mm)	Gear mods	Number of hauls	Year	Dates	Tow duration (h)	<i>Nephrops</i> catch (kg)	Non- <i>Nephrops</i> catch (kg)	Biological factors recorded					
											CL	DL	DS	DM	DP	DV
1	VIa	W	80	SMP	6	2017	06/03-08/03	3.6 \pm 0.3	189 \pm 55	40 \pm 11	Y	Y	Y	Y	Y	Y
			100	SMP	6	2017	15/02-17/02	3.9 \pm 0.2	183 \pm 81	58 \pm 15	Y	Y	Y	Y	Y	Y
	S	80	SMP	6	2016	19/08-17/09	3.6 \pm 0.3	152 \pm 43	95 \pm 41	Y	Y	Y	Y	Y	Y	
		100	SMP	6	2016	15/07-18/08	3.4 \pm 0.5	309 \pm 120	57 \pm 31	Y	Y	Y	Y	Y	Y	
2	IVb	W	80	NetGrid	12	2016	03/02-11/03	3.2 \pm 0.4	121 \pm 95	29 \pm 12	N	N	N	Y	Y	Y
3	IIIa	W	70	SweGrid	3	2015	05/03-17/03	4.1 \pm 0.8	29 \pm 28	63 \pm 39	N	Y	Y	Y	Y	Y
			90	Seltra	3	2015	05/03-17/03	4.1 \pm 0.8	20 \pm 26	212 \pm 78	N	Y	Y	Y	Y	Y
	S	70	SweGrid	3	2015	31/08-03/09	4.0 \pm 0.1	28 \pm 13	91 \pm 12	N	Y	Y	Y	Y	N	
		90	Seltra	3	2015	31/08-03/09	4.0 \pm 0.1	23 \pm 14	167 \pm 167	N	Y	Y	Y	Y	N	

Table 2: Codes for scoring *Nephrops* semi-quantitative assessments of vitality and damage.

Criterion	Description	
Excellent	Vigorous body movement; all limbs moving and tail extends horizontally, flexed or tail-flips	
Good	All limbs moving but tail hangs limp, no tail-flips	
Poor	Limited or no body movement but movement of maxillipeds	
Moribund	Only slight movement of maxillipeds or limbs in response to gentle prodding	
Dead	0	No response/movement to physical stimuli
No injury	1	Alive with no obvious visible injuries
Chelae	D1	Either claw missing or damaged
	D2	Both claws missing or damaged
Rostrum	DR	Rostrum damaged
Body	PUN	A puncture injury on thorax or tail
Thorax	THC	A crush injury on the thorax
	THP	A puncture injury on the thorax
Tail	TAC	A crush injury on the tail
	TAP	A puncture injury on the tail
Eye	EYE	Damage to one or both eyes
Leg	LEG	One or more walking legs missing or damaged

763

764

Table 3: Field and observation tank environmental conditions as ranges. Season: W — winter; S — summer hauls. CT — constant temperature room.

Study	Season	Field sampling environmental conditions					Captive observation tanks		
		Air temp. (°C)	Sea surface temp. (°C)	Bottom temp. (°C)	Sea surface sal.	Sea bottom sal.	CT set air temp. (°C)	Water temp (°C)	Salinity
1	W	6.9–11.5	7.9–8.4	8.0–8.5	34.5–34.7	34.5–34.7	5	6.7–8.2	30.0–34.0
	S	13.8–19.0	13.3–14.7	12.3–14.3	34.0–34.9	34.0–34.9	5	5.7–13.0	31.0–33.0
2	W	8.0–10.0	6.5–7.6	-	-	-	-	5.2–9.2	-
3	W	2.0–5.9	3.6–4.3	4.9–6.1	24.5–26.4	32.9–34.9	10	5.0–6.0	32.0–33.0
	S	14.9–18.7	16.7–17.9	10.0–10.8	23.9–29.1	33.5–34.2	14	14.0–15.0	33.0–34.0

Table 4: GLM model results for immediate mortality modelled as counts of Nephrops alive versus dead during catch sorting, family=quasi-binomial, link=logit. Final models resulting from sequential removal of insignificant terms are shown for models where residual patterns were acceptable. Note in study 2 no immediate mortality was observed.

	Estimate	SE	t-value	p
Study 1 Intercept	2.67	0.24	11.2	<0.001
Study 1 Sorting time	0.32	0.10	-3.2	0.005
Null deviance	86.3	df = 22		
Residual deviance	59.0	df = 21		
Dispersion	2.9			
Study 3 Intercept	4.83	0.66	7.27	<0.001
Study 3 Sea surface temperature	0.17	0.04	-4.42	0.001
Null deviance	67.1	df = 11		
Residual deviance	15.6	df = 10		
Dispersion	1.5			

766

767

Table 5: Percentages of *Nephrops* showing at least one injury by type during catch sorting as mean (95% LCL– 95% UCL from bootstrap in parentheses). Note that individual *Nephrops* may have had more than one injury type and that abdominal and cephalothorax injuries have been combined.

Injury	Study 1 SMP	Study 2 NetGrid	Study 3 Seltra	Study 3 SweGrid
Damaged – at least one injury	40 (36–43)	32 (30–34)	45 (32–54)	37 (32–41)
One chela missing/damaged	24 (22–26)	21 (19–23)	15 (8–25)	11 (6–15)
Puncture wound	8 (6–10)	2(4–5)	24 (9–41)	19 (9–32)
Crush wound	5 (4–10)	6 (3–7)	4 (1–9)	5 (1–11)
Damaged rostrum	3 (2–4)	4 (3–5)	4 (1–6)	3 (1–4)
Two chelae missing/damaged	4 (3–5)	4 (3–5)	3 (1–7)	2 (1–3)

768


769

Table 6: Final survival estimates from the tank-based observation experiments including immediate mortality. Season as W — winter, S — summer. LCL and UCL are the 95% upper and lower confidence limits for the mean survival estimates averaged by gear, season and study with error propagation from the individual haulbased survival estimates.

Study	Season	Codend (mm)	Gear mod.	N	Final survival estimates (%)			
					Mean	Std. err.	LCL	UCL
1	W	80	SMP	6	60.7	2.8	57.3	64.0
	W	100	SMP	6	64.6	2.9	61.2	68.0
	W	Both	SMP	12	61.7	1.4	59.3	64.0
	S	80	SMP	6	52.0	3.6	48.2	55.8
	S	100	SMP	6	42.1	3.3	38.5	45.8
	S	Both	SMP	12	47.1	1.7	44.4	49.7
	Both	Both	SMP	24	55.3	0.7	53.6	57.0
2	W	80	NetGrid	12	57.2	0.9	55.3	59.2
3	W	70	SweGrid	3	75.2	3.1	69.1	81.4
	W	90	Seltra	3	58.6	4.5	49.6	67.6
	W	Both	Both	6	66.9	2.7	61.5	72.4
	S	70	SweGrid	3	41.7	3.1	35.4	48.0
	S	90	Seltra	3	37.7	3.5	30.7	44.7
	S	Both	Both	6	39.7	2.4	35.0	44.4
	Both	Both	Both	12	53.3	1.8	49.7	56.9

Table 7: Linear model results for mean final survival estimates including immediate mortality by haul against available operational covariates for each study. Final models, resulting from sequential removal of insignificant terms are shown where residual patterns were acceptable.

Study 1 — SMP	Estimate	Std. err.	t-value	p
Intercept	90.07	8.79	10.25	<0.001
Sea surface temperature	-1.90	0.79	-2.42	0.025
Sorting time	-6.52	2.60	-2.52	0.020
Residual standard error	10.52			
Multiple R ²	0.47			
F statistic	9.34	df = 2,21		0.001
Study 2 —_NetGrid	No significant operational covariates			
Study 3 — Seltra	Estimate	Std. err.	t-value	p
Intercept	64.83	1.41	45.98	<0.001
Sea surface temperature	-1.56	0.11	-13.96	<0.001
Residual standard error	1.83			
Multiple R ²	0.98			
F statistic	194.9	df = 1,4		<0.001
Study 3 — SweGrid	Estimate	Std. err.	t-value	p
Intercept	84.91	6.51	13.05	<0.001
Sea surface temperature	-2.47	0.52	-4.79	0.009
Residual standard error	8.46			
Multiple R ²	0.85			
F statistic	23.0	df = 1,4		0.009



774

Figure 1: Locations of experimental hauls.

775

776




Figure 2: Tube-set box used to retain *Nephrops* in individual compartments for captive observation.

777

778


767 (a)



768


769

(c)



770 (b)

771



772


773

774 **Figure 3:** Histograms of *Nephrops* length frequencies in the total catches (grey histograms) and
775 discarded portions of the catches (open histograms) in the three study areas. Vertical dashed
776 lines are the minimum landing size (MCRS) at time studies were completed. Figure 3a: Length
777 frequencies in study 1 (ICES Division VIa). Figure 3b: Length frequencies 2011–2017 in ICES
778 Division IVb. Figure 3c: Length frequencies for 2015 in ICES Division IIIa.

779

780

781 781



782

783

784

785

786

787

788

789

790

Figure 4: Estimates of immediate mortality during catch sorting by study (1–3), season (W — winter, S — summer) and gear (cod-end mesh size and gear modification). Heavy vertical bar indicates median, box is the inter-quartile range, whiskers extend up to 1.5 time the interquartile range, circle is an outlier beyond the whisker range.

782

783

791
792
793
794
795
796
797
798
799
800
801
802
803
804
805
806
807
808
809
810
811





Figure 5: Kaplan-Meier survival curves relating the probability of survival of individual *Nephrops* in the observation tanks against presence of physical damage combined with the vitality categories (E/G — ‘Excellent’ or ‘Good’ versus P/M — ‘Poor’ or ‘Moribund’).

812



813

814 **Figure 6:** Relationship between *Nephrops* final mean survival from each haul and recorded sea surface (left hand panel) and air
815 temperatures (right hand panel) across all three studies. Solid lines linear regressions, dotted lines 95% CIs. Points labelled as study
816 number plus season (W — winter, S — summer). The linear regressions are: Survival = 74.0 - 1.9*Sea surface temperature (F = 26.0, df
817 = 1,46, p = <0.001, r² = 0.36); Survival = 75.3 - 1.8*Air temperature (F = 22.2, df = 1,46, p<0.001, r² = 0.33).
818

Table S1: Details of the individual trawl hauls in the studies.

Haul	Study	Date	Season	Cod-end Mesh	Gear mods	Shoot time	Haul time	Shoot lat	Shoot lon	Haul lat	Haul lon	Shoot depth	Haul depth	Haul speed	Weather
				(mm)				(dec deg)	(dec deg)	(dec deg)	(dec deg)	(m)	(m)	(kts)	
1	1	15/07/2016	S	100	SMP	03:28	07:00	56.799	-5.994	56.813	-6.095	79	73	2.5	Slight chop, overcast
2	1	15/07/2016	S	100	SMP	07:35	10:20	56.795	-6.150	56.785	-6.164	104	90	2.5	Slight swell, rain
3	1	29/07/2016	S	100	SMP	05:15	08:30	56.799	-6.155	56.797	-6.152	93	106	2.6	Calm, dry
4	1	29/07/2016	S	100	SMP	09:25	12:30	56.816	-6.243	56.804	-6.243	93	150	2.5	Calm, dry, sunny
5	1	18/08/2016	S	100	SMP	04:48	08:55	57.116	-6.329	57.124	-6.336	106	88	2.8	Calm, clear, sunny
6	1	18/08/2016	S	100	SMP	09:36	13:25	57.121	-6.334	57.116	-6.323	95	148	2.7	Calm, clear, sunny
7	1	19/08/2016	S	80	SMP	04:33	07:53	56.789	-6.055	56.832	-6.125	60	75	2.5	Cloudy, slight swell, dry
8	1	19/08/2016	S	80	SMP	08:46	12:16	56.887	-6.092	56.903	-6.082	119	73	2.4	Cloudy, swell, dry
9	1	16/09/2016	S	80	SMP	06:10	10:04	57.013	-6.595	56.940	-6.636	88	95	2.6	Clear, sunny, slight breeze
10	1	16/09/2016	S	80	SMP	10:30	14:35	56.940	-6.642	57.017	-6.529	97	90	2.7	Clear, sunny, slight breeze
11	1	17/09/2016	S	80	SMP	05:37	09:05	57.128	-6.310	57.151	-6.303	128	144	2.7	Overcast, slight breeze, slight chop
12	1	17/09/2016	S	80	SMP	10:12	13:28	57.104	-6.232			100		2.5	Overcast, breezy, slight swell
13	1	15/02/2017	W	100	SMP	07:50	12:00	56.963	-6.142	56.953	-6.160	86	110	2.5	Breeze, slight swell, overcast
14	1	15/02/2017	W	100	SMP	12:50	16:45	57.031	-6.090	56.979	-6.020	101	128	2.7	Breeze, slight swell, overcast
15	1	16/02/2017	W	100	SMP	07:20	11:08	57.044	-6.219	57.058	-6.203	104	117	2.8	Calm, overcast, slight precipitation
16	1	16/02/2017	W	100	SMP	11:30	15:35	57.077	-6.208	57.052	-6.041	90	88	2.5	Calm, overcast, slight precipitation
17	1	17/02/2017	W	100	SMP	06:46	10:45	56.935	-6.249	56.929	-6.252	104	121	2.4	Calm, overcast
18	1	17/02/2017	W	100	SMP	11:15	14:49	56.923	-6.250	56.879	-6.241	104	128	2.5	Calm, overcast
19	1	06/03/2017	W	80	SMP	07:45	11:40	56.940	-6.257	56.939	-6.258	115	118	2.4	Breeze, slight swell
20	1	06/03/2017	W	80	SMP	12:15	16:15	56.939	-6.254	56.933	-6.196	126	127	2.7	Breeze, slight swell
21	1	07/03/2017	W	80	SMP	08:20	11:25	56.801	-6.149	56.781	-6.183	100	130	2.5	Strong breeze, swell, cloudy
22	1	07/03/2017	W	80	SMP	12:00	15:20	56.781	-6.171	56.798	-6.149	55	62	2.5	Strong breeze, swell, cloudy
23	1	08/03/2017	W	80	SMP	07:10	10:45	56.893	-6.092	56.900	-6.099	55	51	2.6	Windy, strong swell to rough
24	1	08/03/2017	W	80	SMP	11:15	15:00	56.894	-6.089	56.901	-6.096	49	51	2	Windy, strong swell to rough

Table S1 con/td: Details of the individual trawl hauls in the studies

Haul	Study	Date	Season	Cod-end mesh (mm)	Gear mods	Shoot time	Haul time	Shoot lat (dec deg)	Shoot lon (dec deg)	Haul lat (dec deg)	Haul lon (dec deg)	Shoot depth (m)	Haul depth (m)	Haul speed (kts)	Weather
25	2	03/02/2016	W	80	NetGrid	07:15	11:20	55.083	-1.184	55.221	-1.218	59	73	2.6	Slight/Mod
26	2	03/02/2016	W	80	NetGrid	11:50	14:20	55.221	-1.218	55.083	-1.200	73	66	2.6	Mod
27	2	04/02/2016	W	80	NetGrid	07:25	11:05	55.003	-1.196	54.886	-1.067	64	64	2.6	Slight
28	2	04/02/2016	W	80	NetGrid	11:35	15:00	54.886	-1.067	55.017	-1.134	64	62	2.6	Slight
29	2	18/02/2016	W	80	NetGrid	07:15	11:00	54.933	-1.183	54.800	-1.051	55	55	2.6	Slight
30	2	18/02/2016	W	80	NetGrid	11:30	14:45	54.800	-1.051	54.917	-1.138	55	55	2.6	Slight
31	2	19/02/2016	W	80	NetGrid	06:50	09:50	54.933	-1.167	54.800	-1.133	55	55	2.6	Slight/Mod
32	2	19/02/2016	W	80	NetGrid	10:25	13:00	54.817	-1.133	54.933	-1.167	55	51	2.6	Mod
33	2	10/03/2016	W	80	NetGrid	06:30	09:30	54.967	-1.150	54.888	-1.069	51	55	2.6	Slight
34	2	10/03/2016	W	80	NetGrid	10:00	13:00	54.888	-1.069	54.967	-1.151	55	51	2.6	Slight
35	2	11/03/2016	W	80	NetGrid	06:30	09:45	54.967	-1.150	54.867	-1.068	51	55	2.6	Slight
36	2	11/03/2016	W	80	NetGrid	10:00	13:15	54.872	-1.069	54.967	-1.133	55	51	2.6	Slight
37	3	05/03/2015	W	70	SweGrid	07:50	12:00	58.243	11.229	58.253	11.176	52	59	2.5	W 7 m/s, 1.5 m waves, dry
38	3	05/03/2015	W	90	Seltra	07:50	12:00	58.243	11.229	58.253	11.176	52	59	2.5	W 7 m/s, 1.5 m waves, dry
39	3	14/03/2015	W	70	SweGrid	10:35	14:35	58.374	11.082	58.455	11.128	63	54	2.5	NE 8 m/s, 0.5 m waves, mist
40	3	14/03/2015	W	90	Seltra	10:35	14:35	58.374	11.082	58.455	11.128	63	54	2.5	NE 8 m/s, 0.5 m waves, mist
41	3	17/03/2015	W	70	SweGrid	05:50	09:50	58.382	11.057	58.425	11.014	61	68	2.5	E 8 m/s, 0.5 m waves, overcast
42	3	17/03/2015	W	90	Seltra	05:50	09:50	58.382	11.057	58.425	11.014	61	68	2.5	E 8 m/s, 0.5 m waves, overcast
43	3	31/08/2015	S	70	SweGrid	09:23	13:23	58.257	11.240	58.265	11.254	50	50	2.5	SE 3 m/s, calm, overcast
44	3	31/08/2015	S	90	Seltra	09:23	13:23	58.257	11.240	58.265	11.254	50	50	2.5	SE 3 m/s, calm, overcast
45	3	01/09/2015	S	70	SweGrid	06:00	10:04	58.391	11.120	58.393	11.088	50	58	2.5	E 11 m/s, 1 m waves, overcast
46	3	01/09/2015	S	90	Seltra	06:00	10:04	58.391	11.120	58.393	11.088	50	58	2.5	E 11 m/s, 1 m waves, overcast
47	3	03/09/2015	S	70	SweGrid	06:20	10:20	58.413	11.118	58.409	11.110	52	53	2.5	S 8 m/s, 1 m waves, cloudy
48	3	03/09/2015	S	90	Seltra	06:20	10:20	58.413	11.118	58.409	11.110	52	53	2.5	S 8 m/s, 1 m waves, cloudy

Table S2: Kaplan-Meier based survival at the time of the final death in the captive observations from each haul. N observed is the number of test *Nephrops* in the captive observation, excluding additional control animals; LCL and UCL are the 95% confidence limits of the survival; SMP is square mesh panel. The final survival estimates include the immediate mortality estimated during catch sorting.

Haul	Study	Season	Cod-end (mm)	Gear modification	N observed	Time final death (h)	Survival	SE	LCL	UCL
1	1	S	100	SMP	77	218	57.9	5.1	48.8	68.7
2	1	S	100	SMP	100	261	34.5	4.0	27.6	43.2
3	1	S	100	SMP	100	266	41.9	4.6	33.8	51.8
4	1	S	100	SMP	100	261	27.2	4.2	20.1	36.7
5	1	S	100	SMP	100	266	38.1	4.5	30.3	48.0
6	1	S	100	SMP	100	218	53.2	4.5	45.1	62.8
7	1	S	80	SMP	100	240	50.4	4.6	42.2	60.2
8	1	S	80	SMP	100	293	70.0	4.6	61.6	79.6
9	1	S	80	SMP	100	264	47.1	4.6	38.9	56.9
10	1	S	80	SMP	100	118	65.0	4.8	56.3	75.1
11	1	S	80	SMP	99	267	40.7	4.6	32.6	50.9
12	1	S	80	SMP	100	168	38.7	4.7	30.4	49.2
13	1	W	100	SMP	149	264	63.5	3.9	56.3	71.5
14	1	W	100	SMP	150	215	71.3	3.7	64.5	78.9
15	1	W	100	SMP	150	268	65.4	3.7	58.5	73.2
16	1	W	100	SMP	150	264	72.0	3.7	65.2	79.6
17	1	W	100	SMP	150	266	66.0	3.9	58.8	74.0
18	1	W	100	SMP	150	262	75.3	3.5	68.7	82.6
19	1	W	80	SMP	150	266	41.7	3.8	34.8	49.8
20	1	W	80	SMP	150	262	51.5	3.9	44.4	59.7
21	1	W	80	SMP	147	264	70.1	3.8	63.0	77.9
22	1	W	80	SMP	150	264	65.9	3.7	59.0	73.5
23	1	W	80	SMP	136	269	61.0	4.2	53.4	69.8
24	1	W	80	SMP	149	215	59.1	3.8	52.1	67.2
25	2	W	80	NetGrid	212	324	65.1	3.3	59.0	71.8
26	2	W	80	NetGrid	202	324	62.9	3.4	56.5	69.9
27	2	W	80	NetGrid	212	276	54.7	3.4	48.4	61.8
28	2	W	80	NetGrid	199	324	59.8	3.5	53.4	67.0
29	2	W	80	NetGrid	212	348	70.3	3.1	64.4	76.7
30	2	W	80	NetGrid	202	348	64.9	3.4	58.6	71.8
31	2	W	80	NetGrid	212	300	59.4	3.4	53.2	66.4
32	2	W	80	NetGrid	199	300	51.8	3.5	45.3	59.2
33	2	W	80	NetGrid	212	300	32.5	3.2	26.8	39.5
34	2	W	80	NetGrid	202	300	42.1	3.5	35.8	49.5
35	2	W	80	NetGrid	206	300	54.4	3.5	48.0	61.6
36	2	W	80	NetGrid	205	300	68.8	3.2	62.7	75.4

Table S2 con/td: Kaplan-Meier based survival at the time of the final death in the captive observations from each haul. N observed is the number of test *Nephrops* in the captive observation, excluding additional control animals; LCL and UCL are the 95% confidence limits of the survival; SMP is square mesh panel. The final survival estimates include the immediate mortality estimated during catch sorting.

Haul	Study	Season	Cod- end (mm)	Gear modification	N observed	Time final death (h)	Survival	SE	LCL	UCL
37	3	W	70	SweGrid	81	216	66.7	5.2	57.2	77.8
38	3	W	90	Seltra	26	216	61.5	9.5	45.4	83.4
39	3	W	70	SweGrid	40	48	82.5	6.0	71.5	95.2
40	3	W	90	Seltra	77	336	56.8	5.5	47.0	68.7
41	3	W	70	SweGrid	81	240	76.5	4.7	67.9	86.4
42	3	W	90	Seltra	38	336	57.5	7.8	44.1	75.1
43	3	S	70	SweGrid	67	240	48.7	5.6	38.9	61.0
44	3	S	90	Seltra	34	48	37.8	7.2	26.0	55.0
45	3	S	70	SweGrid	62	192	33.3	5.2	24.5	45.4
46	3	S	90	Seltra	68	336	38.3	5.4	29.0	50.5
47	3	S	70	SweGrid	71	144	43.2	5.5	33.7	55.5
48	3	S	90	Seltra	72	216	37.0	5.4	27.9	49.2




Figure S1: Pairs plot for Study 1 operational co-variables. Panels above the diagonal give Kendall correlation coefficients with font size related to significance and significance ($p < 0.05^*$ $p < 0.01^{**}$ $p < 0.001^{***}$). Symbols are open circles - summer hauls; solid circles — winter hauls. Water — sea surface temperature (°C); Air — hopper air temperature (°C); Haul — haul duration (h); Catch — total catch weight (kg); *Nephrops* — catch weight of *Nephrops* (kg); Non-*Nephrops* — catch weight of organisms other than *Nephrops* (kg); Sort — total sorting time (h).




Figure S2: Pairs plot for Study 2 operational covariables. Panels above the diagonal give Kendall correlation coefficients with font size related to significance and significance ($p < 0.05^*$ $p < 0.01^{**}$ $p < 0.001^{***}$). Symbols are solid circles — winter hauls. Water — sea surface temperature (°C); Air — hopper air temperature (°C); Haul — haul duration (h); Catch — total catch weight (kg); *Nephrops* — catch weight of *Nephrops* (kg); Non-*Nephrops* — catch weight of organisms other than *Nephrops* (kg); Sort — total sorting time (h).

824

825




Figure S3: Pairs plot for Study 3 Seltra trawl hauls operational co-variables. Panels above the diagonal give Kendall correlation coefficients with font size related to significance and significance ($p < 0.05^*$ $p < 0.01^{**}$ $p < 0.001^{***}$). Symbols are solid circles — winter hauls; open circles — summer hauls. Water — sea surface temperature ($^{\circ}\text{C}$); Air — hopper air temperature ($^{\circ}\text{C}$); Haul — haul duration (h); Catch — total catch weight (kg); *Nephrops* — catch weight of *Nephrops* (kg); Non-*Nephrops* — catch weight of organisms other than *Nephrops* (kg); Sort — total sorting time (h).

826

827




Figure S4: Pairs plot for Study 3 SweGrid trawl hauls operational co-variables. Panels above the diagonal give Kendall correlation coefficients with font size related to significance and significance ($p < 0.05^*$ $p < 0.01^{**}$ $p < 0.001^{***}$). Symbols are solid circles — winter hauls; open circles — summer hauls. Water — sea surface temperature ($^{\circ}\text{C}$); Air — hopper air temperature ($^{\circ}\text{C}$); Haul — haul duration (h); Catch — total catch weight (kg); *Nephrops* — catch weight of *Nephrops* (kg); *Non-Nephrops* — catch weight of organisms other than *Nephrops* (kg); Sort — total sorting time (h).




Figure S5: Scatterplots of immediate mortality against available covariates from each haul in Study 1. The Kendall tau correlation and its significance ($p < 0.05^*$, $p < 0.01^{**}$, $p < 0.001^{***}$) are shown in each panel. Symbols are solid circles — winter hauls; open circles — summer hauls.




Figure S6: Scatterplots of immediate mortality against available operational covariates from each haul in Study 3 by gear (left hand column — Seltra trawl hauls; right hand column — Swedish grid trawls). The Kendall tau correlation and its significance ($p < 0.05^*$; $p < 0.01^{**}$; $p < 0.001^{***}$) are shown in each panel. Symbols are solid circles — winter hauls; open circles — summer hauls.




Figure S7: Scatterplots of proportion of *Nephrops* with at least one recorded physical injury during catch sorting for studies 1 (left hand column) and 2 (right hand column) against available operational covariates from each haul. The Kendall tau correlation and its significance ($p < 0.05^*$, $p < 0.01^{**}$, $p < 0.001^{***}$) are shown in each panel. Symbols are solid circles — winter hauls; open circles — summer hauls.




Figure S8: Scatterplots of percentage of *Nephrops* with at least one recorded physical injury during catch sorting against available operational covariates from each haul in Study 3 by gear (left hand column — Seltra trawl hauls; right hand column — Swedish grid trawls). The Kendall tau correlation and its significance ($p < 0.05^*$; $p < 0.01^{**}$; $p < 0.001^{***}$) are shown in each panel. Symbols are solid circles — winter hauls; open circles — summer hauls.




Figure S9: Scatterplots of percentage of *Nephrops* sampled for captive observation in the ‘Excellent’ plus ‘Good’ (E/G) vitality category during catch sorting for studies 1 (left hand column) and 2 (right hand column) against available operational covariates from each haul. The Kendall tau correlation and its significance ($p < 0.05^*$ > $p < 0.01^{**}$ > $p < 0.001^{***}$) are shown in each panel. Symbols are solid circles — winter hauls; open circles — summer hauls.




Figure S10: Scatterplots of percentage of *Nephrops* sampled for captive observation in the ‘Excellent’ plus ‘Good’ (E/G) vitality category during catch sorting for Study 2. The Kendall tau correlation and its significance ($p < 0.05^*$ > $p < 0.01^{**}$ > $p < 0.001^{***}$) are shown in each panel. Vitality was not recorded for the summer tows in study 3. Symbols are open circles — winter hauls with Seltra trawl; solid circles — winter hauls with SweGrid trawl.




Figure S11: Kaplan-Meier survival curves relating the probability of survival of individual *Nephrops* in the observation tanks against sex for studies 1 and 3. Solid lines are the mean curves, dashed lines are the 95% confidence intervals. Note that these plots exclude immediate mortality because sex was not recorded for dead *Nephrops* during catch sorting.




Figure S12: Kaplan-Meier survival curves for effect on survival of the presence of different physical injuries: D1 — loss of one chela; D2 — loss of 2 chelae; DR — damaged rostrum; PUN — puncture; CRU — crush. Note that individual *Nephrops* may have had more than one type of injury. For clarity, confidence intervals are omitted.




Figure S13: Kaplan-Meier survival curves for individual *Nephrops* monitored in captivity by season and study. For clarity, confidence intervals are omitted.




Figure S14: Kaplan-Meier estimates of final mean survival, including immediate mortality, at the time of last observation by haul. Symbols indicate the mean and the whiskers are \pm 95% confidence intervals. S — summer, W — winter. By haul symbols: Open circles — Study 1 SMP 100 mm cod-end summer hauls; Filled circles — Study 1 SMP 80 mm cod-end summer hauls; Open triangles — Study 1 SMP 100 mm cod-end winter hauls; Filled triangles — Study 1 SMP 80 mm cod-end summer hauls; Open inverted triangles — Study 2 NetGrid 80 mm cod-end winter hauls; Open diamonds — Study 3 SweGrid 70 mm cod-end winter hauls; Filled diamonds — Study 3 Seltra 90 mm cod-end summer hauls; Open squares — Study 3 SweGrid 70 mm cod-end summer hauls; Filled squares — Study 3 Seltra 90 mm cod-end summer hauls.




Figure S15: Boxplots of observation tank-based Kaplan-Meier final survival estimates, including immediate mortality, amalgamated by study (left panel), study and season (middle panel), and study, season and gear (right panel). Seasons labelled as W — winter, S — summer; Gears labelled as cod-end mesh size plus gear modifications as described in the text. Heavy vertical bars indicate medians and boxes inter-quartile ranges; left and right whiskers are the lower and upper quartile minus or plus 1.5 times the inter-quartile range respectively.




Figure S16: Scatterplots of final mean survival estimates for studies 1 (left hand column) and 2 (right hand column) against available environmental and operational covariates for studies 1 (left hand column) and 2 (right hand column). The Kendall tau correlation and its significance

($p < 0.05^*$ > $p < 0.01^{**}$ > $p < 0.001^{***}$) are shown in each panel. Symbols are solid circles — winter hauls; open circles — summer hauls.




Figure S17: Scatterplots of final mean survival estimates for study 3 against available environmental and operational covariates for study 3 by gear (left hand column — Seltra trawl hauls; right hand column — Swedish grid trawls). Symbols are solid circles — winter hauls; open circles — summer hauls.




Figure S18: Scatterplots of final mean survival estimates for studies 1 (left hand column) and 2 (right hand column) against biological factor percentage of Nephrops in the four injury combined with vitality groups. The Kendall tau correlation and its significance ($p < 0.05^*$)

$p < 0.01^{**}$ > $p < 0.001^{***}$) are shown in each panel. Symbols are solid circles — winter hauls; open circles — summer hauls.




Figure S19: Scatterplots of final mean survival estimates for study 3 against biological factor proportion of *Nephrops* in the four injury combined with vitality groups by gear. Correlations were not computed due to small sample size. Symbols are solid circles — winter hauls; Note that vitality was not recorded for the summer hauls.