# WILEY

### Research Article

## Glucose Variability and Postprandial Hyperglycaemia After Breakfast in Children and Young People With Type 1 Diabetes

Julie Johnson <sup>(1)</sup>, <sup>1</sup> Victoria L. Franklin, <sup>2</sup> Ashley Shepherd <sup>(1)</sup>, <sup>1</sup> Grace Chau <sup>(1)</sup>, <sup>3</sup> Kate Keen <sup>(1)</sup>, <sup>4</sup> Sophie Lennon, <sup>5</sup> Maria Leveridge <sup>(1)</sup>, <sup>6</sup> Kirsty Maclean <sup>(1)</sup>, <sup>7</sup> Julie Nicol, <sup>8</sup> Vanessa Phillipson <sup>(1)</sup>, <sup>9</sup> Sue Roach <sup>(1)</sup>, <sup>10</sup> Adele Swart, <sup>11</sup> and Stuart D. R. Galloway <sup>(1)</sup>

<sup>1</sup>Health Sciences and Sport, University of Stirling, Stirling, UK

<sup>2</sup>NHS Highland, Paediatrics, Raigmore Hospital, Inverness, UK

<sup>3</sup>Nutrition and Dietetics, NHS Grampian, Royal Aberdeen Children's Hospital, Westburn Road, Aberdeen, UK

<sup>4</sup>Nutrition and Dietetics, Epsom and St Helier University Hospitals NHS Trust, Wrythe Lane, Carshalton, Surrey, UK

<sup>5</sup>Nutrition and Dietetics, Betsi Cadwaladr University Health Board, Ysbyty Gwynedd, Penrhosgarnedd, Bangor, Gwynedd, UK

<sup>6</sup>Nutrition and Dietetics, Royal Devon University Healthcare NHS Foundation Trust, Barrack Road, Exeter, UK

<sup>7</sup>Glasgow Children's Diabetes Service, NHS Glasgow and Greater Clyde, Royal Hospital for Children, 1345 Govan Road, Govan, Glasgow, UK <sup>8</sup>Nutrition and Dietetics, NHS Fife, Kirkcaldy Health Centre, Whyteman's Brae, Kirkcaldy, UK

- <sup>9</sup>Nutrition and Dietetics, Surrey and Sussex Healthcare NHS Trust, East Surrey Hospital, Canada Avenue, Redhill, UK
- <sup>10</sup>Nutrition and Dietetics, University Hospitals of Leicester NHS Trust, Leicester Royal Infirmary, Infirmary Square, Leicester, UK <sup>11</sup>Lewisham and Greenwich NHS Trust, Kaleidoscope CYP Centre, 32 Rushey Green, London, UK

Correspondence should be addressed to Julie Johnson; julie.johnson1@stir.ac.uk

Received 23 October 2024; Accepted 18 March 2025

Academic Editor: Andrea Scaramuzza

Copyright © 2025 Julie Johnson et al. Pediatric Diabetes published by John Wiley & Sons Ltd. This is an open access article under the terms of the Creative Commons Attribution License, which permits use, distribution and reproduction in any medium, provided the original work is properly cited.

**Aims:** This study aimed to describe glucose variability (GV) and explore postprandial glycaemia and the association with food composition following breakfast in children and young people (CYP) with type 1 diabetes (T1D).

**Methods:** This was an observational study of CYP aged 1–17 years using continuous glucose monitoring (CGM). Retrospective CGM data were collected to assess GV along with questionnaires about the breakfast meal and 4-h postprandial period for 7 days. Statistical analysis included Student's *t*-tests and linear mixed models.

**Results:** Ninety-six CYP were recruited, 89 shared their CGM data (mean age  $10.1 \pm 3.8$  years) (44.9% females), of which 74 submitted questionnaires. Diurnal percentage coefficient of variation (%CV) (mean:  $38.1\% \pm 4.3\%$ ) was significantly higher than nocturnal %CV ( $36.4 \pm 5.2$ ) (95% confidence interval [CI] [0.9, 2.5], Cohen d = 0.5, p < 0.001). Continuous subcutaneous insulin infusion (CSII) users had a significantly higher time in range (TIR) (mean:  $61.7\% \pm 11.0\%$ ) than those using multiple daily injections (MDIs) (mean:  $55.5\% \pm 15.0\%$ ) (95% CI [0.2, 12], Cohen's d = 0.5, p = 0.04). Data on 387 breakfast meals were analysed. The preprandial glucose was significantly lower for CSII users (mean:  $7.4 \pm 2.3$  mmol/L) compared with MDI users (mean:  $9.5 \pm 2.9$  mmol/L) (95% CI [1.3,3.3], d = 0.9, p < 0.001). Preprandial glucose was significantly associated with mean postprandial glucose ( $R^2 = 0.27$ , p < 0.001). Compared with breakfast meals containing a protein food (n = 71), ingested meals containing breakfast cereals only (n = 76) resulted in a significantly higher mean postprandial glucose (p = 0.01), peak excursion (p = 0.03), area under the curve (AUC) (p = 0.03) and time above range (TAR) (p < 0.001) and significantly higher glucose excursion at 30, (p < 0.001), 60 (p < 0.001) and 90 min (p = 0.02) compared with breakfast meals containing a protein food.

**Conclusions:** GV is significantly higher in the diurnal period. Managing T1D with CSII and including a protein food in the breakfast meal may reduce postprandial hyperglycaemia after breakfast in CYP with T1D. The trial was registered retrospectively with Clinicaltrials.gov (10/08/2024): NCT06546657.

Keywords: carbohydrate; continuous glucose monitoring; diabetes mellitus; glycaemic control; meal; nutrition; protein

#### 1. Introduction

Postprandial hyperglycaemia is a major cause of glucose variability (GV) in type 1 diabetes (T1D) [1], which has been shown to be more harmful than prolonged glucose elevations [2] and is associated with both macro and microvascular complications [3, 4]. Following its discovery, the challenge of matching insulin action with carbohydrate absorption to avoid both hypo- and hyperglycaemia was acknowledged by Banting et al. [5]. The advent of continuous glucose monitoring (CGM) highlighted the problem of GV and postprandial hyperglycaemia in children and young people (CYP) with T1D [6, 7]. There is some evidence that postprandial hyperglycaemia is more likely to occur after breakfast than other meals [8-10]. This is of concern as CYP are often actively learning during the postbreakfast period, and as hyperglycaemia can negatively affect cognitive ability [11], this may put CYP with T1D at a disadvantage compared with their nonT1D peers.

The dietetic management of postprandial glucose is complex. Whilst the preprandial insulin bolus and the quantity of carbohydrate in a meal are the principal influencers of glycaemic control [12], there are many other factors to consider, including the quality of carbohydrate [9], the glycaemic index (GI) of foods and the glycaemic load (GL) of the meal [13]. Diet quality is considered important as it has been suggested that postprandial hyperglycaemia after breakfast is mainly due to the consumption of highly processed carbohydrates [9]. Despite this, there is a paucity of evidence for the type of food that CYP with T1D consume for breakfast, however, national nutritional surveys report that, within the general population, CYP regularly consume ready-to-eat breakfast cereals that are highly processed [14, 15].

In recent years, most of the dietetic research in paediatric diabetes has focused on the impact of fat and protein on postprandial glucose, which has strengthened the empirical evidence that all three macronutrients need to be considered when calculating the prandial insulin dose [16]. One possible solution to managing postprandial hyperglycaemia at breakfast, which is alluded to in the International Society of Paediatric and Adolescent Diabetes (ISPAD) 2022 Nutritional Management Guidelines, is to add protein to the breakfast meal to lower the glycaemic response [13]. However, there is a lack of empirical evidence of the effectiveness of this strategy at breakfast, and most of what is known arises from clinical practice.

The aim of this study was to report on the following in CYP with T1D: (1) Glycaemic variability in the diurnal and nocturnal periods, (2) the composition of the breakfast meal, (3) the extent of postprandial hyperglycaemia at breakfast and (4) the relationship between postprandial glucose at breakfast and the breakfast meal composition.

#### 2. Methods

This observational multicenter study of CYP with T1D who were recruited from 12 diabetes centres in the UK was conducted over a 1-year period in 2021. The inclusion criteria were age of 1–17 years, a diagnosis of T1D for more than 1 year and using insulin regimens of either multiple daily injection (MDI) or continuous subcutaneous insulin infusion (CSII) via an insulin pump along with Dexcom CGM. Those who had other medical conditions or were taking antihyperglycaemic medication, for example, metformin and/or antidepressants, were not eligible to participate in the study owing to the potential impact on glycaemic control. Potential participants brought their own Dexcom CGM G6 devices to the study, and this was approved by Dexcom Inc. Dexcom CGM G6 was the only CGM used in the study as per the inclusion criteria. No data were collected on the number of CYP using Dexcom CGM in the UK, nor how this was funded for those joining the study. However, the majority of Dexcom CGM was self-funded at the time of the study, as CGM was not part of standard treatment provided by the National Health Service (NHS).

The study was advertised to the CYP with T1D who were attending the diabetes centres located in the NHS Trusts that had joined the study. This was achieved by various methods, including waiting room posters, via NHS social media and during clinic appointments. Those CYP who were interested in joining the study had their eligibility assessed and were then recruited by their local dietitian and invited to complete, for 7-days, a twice daily questionnaire about their breakfast meal and the 4-h postprandial period along with the submission of a photograph of their breakfast meal and any food packaging where applicable. Participant information sheets were provided, and consent forms were obtained from the parents or carers and assent forms from the participants.

Baseline information was collected at recruitment from clinical records. This included: sex, date of birth, HbA1c and insulin regimen. The breakfast questionnaire sought information about the breakfast meal and details of diabetes management before and after the meal. This included the timing and duration of the meal as well as the type and amount of food and fluids consumed, and the specific brands of foods where applicable. Diabetes management information included the amount of insulin administered, and any adjustments required for the treatment of hypo- or hyperglycaemia in the preprandial period. The second daily questionnaire was about the 4-h postprandial period and included questions about insulin and food adjustments that were required for hypo- or hyperglycaemia. This questionnaire also sought information about any food and fluid intake and physical activities undertaken during the 4-h postprandial period.

The questionnaire submissions and meal photographs were reviewed and analysed by a dietitian using Nutritics (Nutrition Analysis Software, 2019) and from the food labels provided via photographs. The analysis included: energy (kcal), fats (g), saturated fats (g), carbohydrate (g), sugars (g), protein (g), fibre and salt (g) and GI and GL. Macronutrient intakes were compared with ISPAD recommendations [13].

The GI values of the carbohydrate foods contained within each breakfast meal were obtained using the International tables of GI [17, 18]. Where the GI was not known, the GI of a similar food was used. The total GI value of each breakfast meal was estimated by summing the means of the carbohydrate foods as described in Dodd et al. [19]. No adjustments were made for the impact on the GI of the fat and protein contained in the meals. The GI values of carbohydrate foods have been shown to correlate with the glucose response to mixed meals [20–22], and the GI is the main determinant of the glycaemic response to mixed meals [22]. The GL was calculated by multiplying the GI of the individual food by its available carbohydrate and dividing by 100 [23].

The breakfast meals were also categorised into two subgroups of breakfast cereal-only meals (cereal only) and those which included a protein food (added protein). The cerealonly subgroup included meals that contained a ready-to-eat cereal with milk and with no other foods included. The added protein subgroup included the meals which contained a food from the protein food group as per the UK Eat Well Guide [24], in addition to cheese from the dairy food group. Insulin dose timing was categorised into three types of lag time:  $\geq 15$ min and < 15 min preprandial and postprandial dosing.

CGM data were collected retrospectively, via Dexcom Clarity, for up to 90 days to assess diurnal (06.00–22.00) and nocturnal (22.00–06.00) periods for analysis of GV and for the 7-day breakfast questionnaire recording period. For the assessment of GV, the CGM metrics included: mean glucose and percentage coefficient of variation (%CV) along with the time in range (TIR) metrics of time below range (TBR), TIR and time above range (TAR) [25].

For the breakfast meal analysis, the primary outcome measurement was the mean postprandial glucose, which was taken from the CGM sensor readings at 5-min intervals over 240 min. The following outcomes, which were taken from the CGM sensor readings over the 240 min period and the questionnaire responses, included (1) preprandial glucose (this was the closest reading taken prior to the breakfast meal commencing); (2) the timing of the insulin dose; (3) mean glucose excursion (defined as the change in glucose levels from baseline); (4) peak glucose excursion; (5) time to peak; (6) the mean area under the curve (AUC) (defined as the sum of the AUC of glucose excursions above baseline glucose over the 240 min period); (7) %CV (defined as the ratio of the standard deviation (SD) to the mean glucose) and (8) TBR, TIR and TAR. Only those breakfast meals which had  $\geq$ 70% of CGM data were included in the analysis as recommended in Schnell et al. [26]. For the analysis of glucose excursions, only those breakfast meals which had a reading at baseline were included in the analysis. This applied to the analysis of the mean glucose excursion, peak excursion, time to peak and AUC. No adjustments were made for other missing CGM data points.

Statistical analysis was performed using Jamovi (The Jamovi project, Sydney, Australia, 2023). (Version 2.2.5 retrieved from https://www.jamovi.org). Figures were produced using Graph-Pad 9.5.0 (GraphPad Software, San Diego, California, USA). The incremental AUC was calculated using the trapezoidal method via GraphPad 9.5.0 (GraphPad Software, San Diego, California, USA). Parametric outcomes are presented as mean and SD, and for nonparametric outcomes, the median and interquartile range (IQR) are presented. GV outcomes were analysed using Student's *t*-tests. For the breakfast postprandial period, paired Student's *t*-tests and simple linear regression were

TABLE 1: Participant demographic and diabetes characteristics.

Characteristic	N = 89
Age (years)	$10.1\pm3.8$
Females $(n)$ $(\%)$	40 (44.9%)
Males ( <i>n</i> ) (%)	49 (55.1%)
Duration of T1D (years)	4.0 (4.5)
Regimen – CSII ( <i>n</i> ) (%)	66 (75%)
Of which - HCL, $n$ (%)	14 (16%)
Regimen – MDI, $n$ (%)	22 (25%)
HbA1c (mmol/mol)	$57.7\pm9.3$

*Note*: Mean  $\pm$  SD, median (IQR). Regimen of one participant was unknown.

performed to compare pre and postprandial glucose measurements. In order to account for the nonindependent data and correlations from repeated measurements from the same participants and the resultant risk of pseudo replication, the remaining outcomes from the postprandial breakfast period were analysed using linear mixed models. Furthermore, linear mixed models were used as these are superior at handling missing data compared with repeated measures analysis of variance (ANOVA). The linear mixed models were performed with restricted maximum likelihood (REML), the Wald method for confidence intervals and the Satterthwaite method for the degrees of freedom. The models included a fixed intercept and slope and a random intercept for the participants. Hypoglycaemic events and insulin correction events for hyperglycaemia were analysed using a generalised linear mixed model with logistic regression. The effect size for the linear mixed models was calculated using the suggested formula as described in Westfall et al. and Brysbaert and Stevens [27, 28].

Where appropriate, post hoc tests were performed using the Bonferroni correction to manage the risk of a type 1 error occurring with multiple statistical tests. p < 0.05 was considered statistically significant.

#### 3. Results

3.1. Participants. Ninety-six participants consented to join the study and provided the baseline information. One was unable to share their CGM data, and six did not respond, resulting in 89 participants who shared their CGM data (Table 1), of which 74 submitted a total of 428 breakfast questionnaires with 368 corresponding postprandial questionnaires. Some meals had missing CGM data during the postprandial period, and this resulted in 387 breakfast meals remaining. Three of the 74 participants submitted a total of four breakfast questionnaires; however, of these, one did not consume any food at breakfast, and the other three meals had no corresponding CGM data, so these were excluded from the analysis, resulting in 71 meal participants.

3.2. GV. Data exported from Dexcom Clarity provided a mean of 79.3 days of CGM sensor readings. Mean percentage of %CV was  $37.8\% \pm 4.2\%$ ; this was significantly higher in the diurnal (mean:  $38.1\% \pm 4.3\%$ ) compared with the nocturnal period (mean:  $36.4\% \pm 5.2\%$ ) with a mean difference of 1.7% (95% CI



FIGURE 1: Comparison of diurnal and nocturnal %CV. Statistically significant difference is indicated by \*\*\*\*p < 0.0001.

[0.9, 2.5], Cohen's d = 0.5, p < 0.001) (Figure 1) and 36% of the participants met the %CV target of ≤36%. Diurnal %CV was significantly lower for those using MDI (mean:  $36.4\% \pm 3.1\%$ ) compared with those using CSII via an insulin pump (mean:  $38.6\% \pm 4.5\%$ ) with a mean difference of 2.3% (95% CI [0.5, 4.0], Cohen's d = 0.6, p = 0.01). The mean percentage TIR was  $60.1\% \pm 12.5\%$  with no significant difference between the diurnal and nocturnal periods for this range of times (p>0.05). CSII users had a significantly higher %TIR (mean:  $61.7\% \pm 11.0\%$ ) compared with those using MDI (mean: 55.6%  $\pm$  15.0%), with a mean difference of 6.1% (95% CI [0.2, 12], Cohen's d = 0.5, p = 0.04). The median %TBR was significantly lower for those using MDI (1.0% [1.8]) compared with CSII (all types) (3.0% [3.0]) (Mann Whitney U = 461, p = 0.01); this was significant for diurnal %TBR (Mann Whitney U = 486, p = 0.02) but not the nocturnal period (p =0.05). For the %TIR target of  $\geq$ 70%, 24.7% met this, whilst 77.5% met the target of <5% for TBR. There was no significant difference in those meeting and not meeting the %TIR target for ages, disease duration, sex or regimen (p>0.05). Comparisons of GV between the CSII users who used either open loop or hybrid closed loop (HCL) systems were not analysed due to the small number of participants using HCL systems.

*3.3. Preprandial Glucose.* Of the breakfast meals which had CGM data and were included in the analysis (n = 387), 54.5% (n = 213) had a preprandial reading within the fasting

glucose target of 4–8 mmol/L; 41.7% were above target, and the remaining 3.8% were below target. The preprandial glucose was significantly lower for the CSII users (mean: 7.4  $\pm$  2.3 mmol/L) compared with MDI users (mean: 9.5  $\pm$  2.9 mmol/L) with a mean difference of 2.2 mmol/L (95% CI [1.3, 3.3], d=0.9, p<0.001).

3.4. Correction Doses and Dose Timing. The odds of needing a correction dose of insulin before breakfast were 3.5 times greater for those using MDI compared with CSII (95% CI Exp [ $\beta$ ] [1.1, 11.2], p = 0.03). The dose time was reported for 380 of the meals which had corresponding CGM data. Of these meals, 45.8% (n = 174) were managed with an insulin dose lag time of  $\geq$ 15 min, 44.2% (n = 168) had a dose lag time of  $\leq$ 15 min and 10% (n = 38) of the meals were managed with a postprandial dose.

3.5. Postprandial Glucose. Mean postprandial glucose was 9.2  $\pm$  2.7 mmol/L and the mean glucose excursion was 1.4  $\pm$  2.6 mmol/L. Two-thirds of the breakfast meals (n=262, 67.7%) resulted in a mean postprandial glucose which was within the TIR target of 3.9–10 mmol/L and 32% of meals (n=124) resulted in a sensor reading >10 mmol/L, of which 5.9% of meals (n=23) were above >13.9 mmol/L. Mean postprandial glucose (mean: 9.2  $\pm$  2.7 mmol/L) was significantly higher than the mean preprandial glucose (mean: 8.0  $\pm$  2.6 mmol/L) with a mean difference 1.2 mmol/L (95% CI [0.9, 1.4], Cohen's d=0.5, p<0.001) and this was observed for meals managed with both CSII and MDI.

There was a significant, positive association between pre and postprandial glucose (Figure 2). Every 1 mmol/L increase in preprandial glucose corresponded with a 0.5 mmol/L increase in mean postprandial glucose (95% CI [0.4, 0.6], d=0.2, p<0.001). This remained significant when adjusting to include covariates and factors of age, disease duration and sex.

Mean postprandial glucose was significantly lower for the meals ingested by the CSII users (mean:  $8.8 \pm 2.5 \text{ mmol/L}$ ) compared with the MDI users (mean:  $10.2 \pm 3.2 \text{ mmol/L}$ ) with a mean difference of 1.5 mmol/L (95% CI [0.4, 2.6], d=0.6, p<0.01). This remained significant after adjusting the model with covariates and factors of age, disease duration and sex, albeit with a higher p value (p=0.03); however, it was not significant when preprandial glucose was also added to the model as a covariate (p=0.58). There was no significant difference in the postprandial glucose measurements across the three types of insulin dose timing.

3.6. The Breakfast Meal. The mean GI of the breakfast meals was  $59\% \pm 10\%$  and the mean GL was  $26 \pm 13$ . The low GI meals accounted for 36% (n = 139) whilst 49.5% (n = 191) were in the medium GI category and the remaining 14.5% of meals (n = 56) were in the high GI category. When categorised by GL, the majority of the breakfast meals, 65.6% (n = 252), were in the high GL category ( $\geq 20$ ), followed by 28.1% of meals in the medium (n = 110) and 6.3% (n = 24) had a low GL category (<10). The proportion of energy from the three macronutrients for all meals was  $52.9\% \pm 14.3\%$  from carbohydrate,  $31.2\% \pm 14.9\%$  from fat and  $14.2\% \pm 5.9\%$  from protein. The mean fibre content was  $3.0 \pm 1.9$  g.



FIGURE 2: Scatterplot with simple linear regression of the relationship between pre and post prandial glucose of 376 breakfast meals consumed by 71 participants for all types of insulin regimen.

The low GI meals were the only breakfast meals to meet the ISPAD recommendations for macronutrient percentage from energy [13], although these recommendations are for all meals, not just breakfast. The protein content of the low GI meals was significantly higher than both the medium GI meals (95% CI [0.8, 4.0], adjusted p = 0.01) and high GI meals (95% CI [2.2, 6.9], p < 0.001).

The most popular main source of carbohydrate was breakfast cereals; this was the main carbohydrate source of 44.6% of meals (n = 191), followed by breads (33.4%, n = 143) and pastries (9.1%, n = 39). Only the breakfast cereals had a mean GI that was in the high GI category. Almost a fifth of meals (18.3%) contained cereal only as the main source of carbohydrate (plus milk). These cereal-only meals contained a significantly higher GI (mean:  $67\% \pm 7\%$ ) than all the other meals (mean:  $57\% \pm 9\%$ ) with a mean difference of 9% (95% CI [6.5, 11.1], *d* = 1.0, *p* < 0.001). They also contained a significantly higher GL (mean:  $28 \pm 15$ ) than all the other breakfast meals (mean:  $25 \pm 13$ ) with a mean difference of 3 (95% CI [0.1, 5], d = 0.2, p = 0.04). These cerealonly meals also contained significantly more carbohydrate per percentage of energy (mean:  $63.3\% \pm 9.0\%$ ) than all the other breakfast meals (50.6%  $\pm$  14.3%), with a mean difference of 14.3% (95% CI [11, 18], d = 1.1, p < 0.001).

A fifth of breakfast meals (19.6%) included a protein food. The protein content was significantly higher (mean:  $20.2 \pm 9.3$  g) than all the other meals (mean:  $10.3 \pm 6.0$  g) with a mean difference of 10.6 g (95% CI [9.1, 12.1], d=1.6, p<0.01). The percentage of energy from carbohydrate was significantly lower for the added protein meals (mean:  $38.8\% \pm 13.4\%$ ) compared with all the other breakfast meals (mean:  $56.4\% \pm 12.3\%$ ) with a mean difference of 19.8% (95% CI [16.8, 22.8], d=1.6, p<0.001). 3.7. The Breakfast Postprandial Period. Ingestion of the high GI breakfast meals resulted in a significantly lower %CV (mean: 20.4%  $\pm$  7.8%) than the medium GI meals (mean: 24.7%  $\pm$  9.9% with a mean difference of 4.4% (95% CI [1.2, 7.5], d=0.4, p=0.01, adjusted p=0.02). There were also fewer hypoglycaemic events following ingestion of the high GI meals (n=9, 16%) than after ingestion of the medium GI meals (n=62, 32.5%) with borderline significance (p=0.06). Ingestion of the high GL meals resulted in a faster time to peak (mean:  $118 \pm 76$  min) than both medium (mean:  $132 \pm 79$  min) and low GL meals (mean:  $148 \pm 87$  min) combined with a mean difference of 22 min (95% CI [0.6, 44], d=0.3, p=0.04). There were no other significant differences in the glucose outcome measurements, nor any difference in the excursion at 30-min intervals between the categories of GI and GL (p>0.05).

The ingestion of the cereal-only meals resulted in a significantly higher mean postprandial glucose over the 240 min (mean:  $9.6 \pm 2.7 \text{ mmol/L}$ ) compared with the added protein meals (mean:  $8.5 \pm 2.0 \text{ mmol/L}$ ) with a mean difference of 1.3 mmol/L (95% CI [0.4, 2.1], d = 0.5, p = 0.01).

The peak excursion was also significantly higher following the ingestion of the cereal-only meals (mean:  $5.2 \pm 3.2$  mmol/L) compared with the added protein meals (mean:  $4.5 \pm$  2.6 mmol/L) with a mean difference of 1.2 mmol/L (95% CI [0.6, 2.3] d=0.4, p=0.03). The time to peak was significantly shorter following the ingestion of the cereal-only meals (mean:  $102 \pm 71$  min) compared with the added protein meals (mean:  $135 \pm 77$  min) with a mean difference of 35 min (95% CI [9, 62], d=0.5, p=0.01). The mean AUC was also significantly higher following the ingestion of the cereal-only meals (mean:  $515 \pm$ 463 [mmol/L] \*min) compared with the added protein meals (mean:  $442 \pm 346$  [mmol/L] \*min) with a mean difference of 167 (mmol/L) \*min (95% CI [20, 313], d=0.4, p=0.03).

TABLE 2: Comparisons of the glucose measurements taken from CGM readings over 240 min after ingestion of cereal-only and added protein breakfast meals.

Glucose measurements	Cereal only	Added protein	Þ
Preprandial glucose (mmol/L)	$8.5\pm2.7$	$7.6\pm2.5$	ns
Mean postprandial glucose (mmol/L)	$9.6\pm2.7$	$8.5\pm2.0$	0.01
Mean excursion (mmol/L)	$1.3\pm2.6$	$1.3\pm2.0$	ns
Peak excursion (mmol/L)	$5.2 \pm 3.2$	$4.5\pm2.6$	0.03
Time to peak (min)	$102\pm71$	$135\pm77$	0.01
AUC ([mmol/L] *min)	$515\pm463$	$442\pm346$	0.03
%CV	$25.3\pm10.4$	$22.3\pm10.8$	ns
Hypo events ( <i>n</i> ) (%)	21 (29.6)	18 (23.7)	ns
Correction events ( <i>n</i> ) (%)	17 (27.4)	16 (22.2)	ns
TIR (min)	$136\pm76$	$167\pm70$	0.01
TAR, high (min)	$59\pm49$	$54\pm58$	ns
TAR, very high (min)	$35\pm51$	$10\pm26$	< 0.001

*Note*: Mean  $\pm$  SD. Linear mixed model fit by REML with fixed intercept and random intercept (subjects). Cereal-only meals n = 71, added protein meals n = 76 (number with primary outcome measure of mean postprandial glucose).

Abbreviation: ns: nonsignificant.

The ingestion of the cereal-only meals resulted in a less TIR (mean:  $136 \pm 76$  min) than the added protein meals (mean:  $167 \pm 70$  min) with a mean difference of 38 min (95% CI [11, 64], d=0.5, p=0.01) and more TAR with a mean difference of 31 min, (95% CI [17, 46], d=0.7, p<0.001) (Table 2).

At 30 min, the glucose excursion was significantly higher after ingestion of the cereal-only meals (mean: 1.8  $\pm$ 2.3 mmol/L) compared with the added protein meals (mean:  $1.0 \pm 2.2$  mmol/L) with a mean difference of 1.5 mmol/L (95% CI [0.7, 2.3], d = 0.6, p < 0.001). This remained significantly higher at 60 min with a mean difference of 2.3 mmol/L (95% CI [1.2, 3.3], d=0.7, p<0.001) and at 90 min with a mean difference of 1.5 mmol/L (95% CI [0.2, 2.7], d = 0.4, p = 0.02). These differences at these times remained significant when the model was adjusted to include the type of regimen. There was no other significant difference in the glucose excursions until 240 min when the glucose excursion was significantly lower after ingestion of the cereal-only meals compared with the added protein meals with a mean difference of 1.4 mmol/L (95% CI [0.1, 2.8], p = 0.04); however this was not significant when the model was adjusted to include type of regimen (p = 0.44) (Figure 3).

3.8. Snacks and Physical Activities. The mean postprandial glucose was significantly lower where the participant had a snack during the postprandial period (mean:  $8.5 \pm 2.4 \text{ mmol/L}$ ) compared with those who did not snack (mean:  $9.5 \pm 2.8 \text{ mmol/L}$ ) with mean difference of 1 mmol/L (95% CI [0.5,1.6], d=0.4, p<0.001). The TIR (min) was also significantly higher for those who snacked during the postprandial period (mean:  $163 \pm 72 \text{ min}$ ) compared with those who did not snack (mean:  $138 \pm 79 \text{ min}$ ) with a mean difference of 26 min (95% CI [9, 43], d=0.3, p=0.01). The activity duration was significantly longer for those who snacked (mean:  $45 \pm 58 \text{ min}$ ) compared to the nonsnack group ( $29 \pm 53 \text{ min}$ ) with a mean difference of 14 min (95% CI [0.5, 28], d=0.3, p=0.04). Activity duration of those who took a snack without insulin was significantly longer (mean:  $62 \pm 69 \text{ min}$ ) than those who

took insulin with their snack  $(31 \pm 42 \text{ min})$ , with a mean difference of 36 min (95% CI [16, 57], d = 0.6, p < 0.001).

#### 4. Discussion

4.1. GV. Although only 36% of the participants met the %CV target of  $\leq$ 36% [25], which is the level of stable glycaemia [29], the mean %CV is lower than others have reported [30–34], suggesting that GV for CYP with T1D is improving. This improvement reflects the use of real time CGM (rtCGM) by all the participants in the present study, as this is known to be superior at improving glycaemic control compared with intermittentscan CGM (isCGM) [33], which was used in some of the other studies reporting on GV [30, 33, 34]. Furthermore, more recent studies have also reported similar levels of %CV to the current finding [35–38].

The higher diurnal %CV compared with the nocturnal period was also observed by both Tansey et al. [7] and Edge et al. [30]. Although in the present study the mean diurnal glucose was lower than in both these previous studies [7, 30], the higher diurnal %CV demonstrates that managing GV in the diurnal period, when food consumption and physical activity occur, remains a challenge. The significantly lower diurnal %CV observed for the MDI users was due to them spending less TBR than the CSII users. This conflicts with findings showing that hypoglycaemia tends to occur less often in those using CSII, however, this is observed for severe hypoglycaemic events [39] and for mild hypoglycaemia, there appears little difference between MDI and CSII users [40]. Other studies have also reported a lower %CV in MDI users when using rtCGM [35, 41]. It is also known that the use of rtCGM, regardless of regimen, reduces the risk of hypoglycaemia [42], which may explain the current findings.

Achieving a TIR of  $\geq$ 70% is more challenging. In this present study, the mean TIR was 60.1%, and only 24.7% of the participants met the  $\geq$ 70% target. Several studies have observed a similar level of %TIR for CYP with T1D [31, 32,



FIGURE 3: Comparison of the glucose excursion from CGM readings over 240 min following the ingestion of "cereal-only" and "added protein" breakfast meals. Number of meals: cereal-only n = 70 and added protein n = 75. Statistically significant difference is indicated by "p < 0.05; \*\*\*p < 0.001.

35, 43], and the higher %TIR for the CSII users compared with MDI users was also observed by Cherubini et al. [41].

The low number of participants meeting the %TIR target demonstrates how difficult this target is to achieve; in a study based at a diabetes camp an improvement in %TIR during camp was observed from 58% to 67% however the authors noted that this involved "great professional effort" and during camp only 30.8% met the %TIR target [44]. However, in this previous study, some of these CYP were using isCGM and none were using HCL systems, which have been shown to significantly improve glycaemic control, including TIR, from both randomised controlled and single arm trials [45–49]. Similarly, in the present study, few participants were using HCL systems, which likely explains the low number of those who met the TIR target. As increasing numbers of CYP in the UK have access to HCL systems, it is expected that this TIR target will be more achievable for CYP with T1D.

4.2. The Preprandial Period. Similar levels of preprandial glucose before breakfast have been reported elsewhere [6, 50, 51]. This present study suggests that CSII users experience more optimal preprandial glucose levels, however, Heptulla et al. [50] did not observe any difference in the preprandial glucose when the participants in their study moved from using MDI to CSII. Despite this, the "Dawn Phenomenon" [52] is the likely cause of the higher preprandial glucose observed in the MDI users as basal insulin, administered via injections, cannot be adjusted to meet the diurnal and nocturnal variability of insulin requirements unlike CSII via an insulin pump and particularly HCL systems [45, 53].

Preprandial glucose is an important factor for postprandial glycaemia. It is known that in adults with noninsulindependent type 2 diabetes, preprandial glucose is strongly positively correlated with postprandial glucose [54], although once glucose levels exceed the renal threshold for reabsorption, there is an inverse association with postprandial glucose [55]. There is also some evidence that preprandial glucose is associated with postprandial hyperglycaemia in adults with T1D [56], as was observed in this study.

The mean insulin dose timing met the ISPAD 2022 guidance of 10–15 min preprandial [57] although national registry data report that up to a third of CYP do not adhere to these recommendations [58, 59] and none of the dose timings resulted in a difference in glucose outcomes which contrasts with the review by Mozillo et al. [60] who concluded that preprandial dose timing was associated with lower peak glucose compared with doses given postprandial.

4.3. *The Postprandial Period.* Reporting on data collected using the first Food and Drug Administration (FDA) approved CGM, Boland et al. [6] highlighted the issue of GV caused by postprandial hyperglycaemia in CYP with T1D, however, they did not observe this to be significantly worse after break-fast than after other meals. In contrast, other studies, also using CGM, observed that postprandial glucose was worse after breakfast compared to other meals [8–10].

The findings of this present study highlight that postprandial hyperglycaemia can occur after breakfast, however, two-thirds of the breakfast meals analysed resulted in a postprandial glucose within the TIR target, therefore, good

glycaemic control after breakfast is possible. The glucose levels were also lower than others have reported [6, 9, 10, 50]. In Boland et al. [6], in their study of 56 CYP with T1D, the majority of whom were using CSII, the peak postprandial glucose after breakfast was  $16.3 \pm 4.7$  mmol/L. This was a similar finding to Heptulla et al. [50] who observed a significant reduction in postprandial glucose after breakfast in their subjects when they were moved from injection therapy to CSII therapy, although this remained above target at  $14.8 \pm$ 0.9 mmol/L (p < 0.003). In Gandrud et al. [9], a study involving children with T1D aged under 7 years, the glucose excursion after breakfast was also above target at a mean of 7.5  $\pm$ 3.9 mmol/L although some of the subjects had higher levels of HbA1c than in the present study, which these same authors observed to be associated with postprandial hyperglycaemia after breakfast.

In the present study, the mean glucose excursion did not exceed recommendations [61]. and as this was also lower than reported elsewhere [6, 9], this suggests that postprandial hyperglycaemia after breakfast may be improving for CYP with T1D. However, this contrasts with the findings of Monzon et al. [10], who reported the mean postprandial glucose after breakfast to be 14 mmol/L, and this was higher after breakfast compared with both lunch and dinner. These findings likely differ because the study involved CYP from the USA, who have been shown to experience higher levels of poor glycaemic control [62]. Furthermore, postprandial hyperglycaemia seems to affect some CYP more than others; in the present study, a third of meals resulted in a mean glucose reading above 10 mmol/L, which is not acceptable, and as those using MDI had a higher preprandial reading, they also experienced higher postprandial glucose.

4.4. The Breakfast Meal. Some of the foods in the present study were like those reported by Mackey et al. [63] for children aged 2–5 years with T1D, although breakfast cereals did not appear to be commonly consumed in Mackey et al., unlike in the present study. The consumption of high GI breakfast cereals is observed in clinical practice. Lower GI cereals are available and account for over a third of those reported in the international tables of GI [18]; however, not all these products are available in the UK.

The percentage of energy from carbohydrate of  $52.9\% \pm 14.3\%$  is like the findings of Fisher et al. [64] (2023), who reported this to be 52.3% for breakfast in their sample of 48 Australian CYP with T1D. In the present study, the percentage of energy from protein was  $14.2\% \pm 5.9\%$ , which was lower than in Fisher et al. [64], who reported these to be a mean of 16.8%, although none of the breakfast meals in Fisher et al. were high in protein. Studies including CYP with T1D from the USA have reported both higher and lower amounts of carbohydrate with Monzon et al. [10] observing an intake of 62% of energy from carbohydrates at breakfast for children aged 2–6 years with T1D, which was higher than the 48.5\% reported in Mackey et al. [63] in a similar age group of CYP with T1D. These studies demonstrate that carbohydrate intake at breakfast in CYP with T1D is variable.

The mean GI of  $59\% \pm 10\%$  is higher than reported by Mackey et al. [63], who observed a GI of 51.6% in breakfast meals consumed by children aged 2–5 years with T1D. As only the low GI breakfast meals met the ISPAD recommendations for macronutrient distribution for all meals [13], this appears to substantiate the recommendation for the GI to be incorporated into carbohydrate counting education. In contrast, the low GL meals only contained 36% of carbohydrate from energy, which would be classed as a low carbohydrate diet if this was followed for all meals [65, 66]. Low intakes of carbohydrate without increases in protein and fat will result in inadequate energy intake and weight loss in CYP [67]. It appears more clinically beneficial to focus on the GI rather than GL for ensuring the ISPAD macronutrient recommendations are met.

4.5. Postprandial Glucose and Food Composition. In the present study, the ingestion of the high GI breakfast meals resulted in a significantly lower %CV compared with medium GI meals and fewer hypoglycaemic events were observed with borderline significance. This infers that the consumption of lower GI meals may increase the risk of hypoglycaemia. There is conflicting evidence in the literature for this. Two studies of adult subjects with T1D did not observe more frequent hypoglycaemic events following consumption of lower GI meals [68, 69] although the latter did report that hypoglycaemic events occurred earlier in the postprandial period following the ingestion of a low GI meal. In their study of CYP with T1D, Nansel et al. [70] observed mild hypoglycaemia and the need for lower doses of insulin when low GI meals were consumed compared to high GI meals. As TBR correlates with %CV [71], hypoglycaemia caused by the consumption of lower GI meals may result in higher levels of GV, and these meals may require insulin adjustment [13].

The largest difference in glycaemic response was observed between the meals which contained only breakfast cereal and those which included a protein food. The significantly higher glucose excursions after the ingestion of the breakfast cerealonly meals are also consistent with other findings [72–74]. Parents also report that breakfast cereals are a problem food, causing high glucose levels [75]. The breakfast cereal-only meals had a significantly higher GI. Foods with a high GI can cause a large rise in the early postprandial period [72, 76, 77] followed by a drop in the later postprandial period, which can cause hypoglycaemia [73]. Although there was no difference in hypoglycaemic events, the breakfast cereal meals did result in a significantly lower glucose excursion in the later postprandial period.

The breakfast cereal meals also had a significantly larger carbohydrate load as per percentage of energy, which may negatively impact longer-term glycaemic control. Mackey et al. [63] observed a trend towards an HbA1c above target for those children who consumed higher amounts of carbohydrate for breakfast as per percentage of energy, however, this was not significant.

In contrast to high GI foods, meals which contain fat or protein can result in a lower glucose excursion in the early postprandial period [78, 79], and this was observed after

ingestion of the "added protein" breakfast meals. CYP with T1D who spend more than 70% of their TIR have been observed to consume more protein than those with lower TIR [80], and in the present study, adding protein to the breakfast meal also resulted in the longest TIR with significantly less time spent above 13.9 mmol/L. In nonT1D, the addition of protein to a high GI carbohydrate meal has been shown to reduce the AUC by almost half [81]. This can be beneficial in lowering the GI of a meal, however, in CYP with T1D, high intakes of protein and fat can cause delayed hyperglycaemia between three and 5-h postprandial [78, 82, 83] with an additive effect in meals which are high in both these macronutrients [82]. This effect can increase the need for more correction doses of insulin during the postprandial periods after ingestion of high-protein meals [84]. However, this was not observed in the present study and following the ingestion of the "added protein" meals, the glucose excursion started to fall by 150 min with no delayed hyperglycaemia and no significant difference in the number of correction events for any type of meal, suggesting that this amount of protein which was a mean of  $20.2 \pm 9.3$  g was not enough to cause delayed hyperglycaemia.

The glucose levels were lower for those who snacked during the postprandial period, and the physical activity of these participants was also of a longer duration. This suggests that much of the snacking that took place was to manage glucose levels during activity, especially for those who did not cover the snack with insulin. Fisher et al. [64], in a diet survey study, found no association between snacks and HbA1c, and furthermore, GV has been shown to be lower for those who snack between meals [10], which is likely due to fewer hypoglycaemia events.

There are several limitations to this study. As an observational study, it lacks randomisation, which can lead to bias and the overestimation of treatment effects when compared with randomised controlled trials [85]. There is also potential for confounding factors as these are not controlled for between groups in observational studies [86]. There were potential confounding factors in the present study arising from the participants' socioeconomic status and food choice, which could have affected glycaemic control. The sample size was also small, and although bias was controlled for by not limiting recruitment to those with good glycaemic control, this may have arisen from only recruiting those using Dexcom CGM. At the time of recruitment, fewer CYP had access to CGM, and those who did have access to CGM were more likely to be from a higher socioeconomic status, as there was a socioeconomic disparity in access to technologies [87]. Furthermore, although only those CYP who had T1D for at least 1 year were recruited to the study, it is not known if any were in partial remission, which would have affected their glycaemic control. The intake of meals was also self-reported, and although the accuracy of this was likely increased via the collection of photographs, the accurate analysis of the breakfast meals did rely on correct submissions. Although this study highlights that high GI ready-to-eat cereals are a commonly consumed breakfast meal, the study did not explore other commonly consumed breakfast foods among different ethnic groups and the impact these have on glycaemic control after breakfast.

#### 5. Conclusions

This study provides some evidence of the impact that breakfast meal composition has on postprandial glucose. There are several facets to this, including the GI, the GL and carbohydrate load and the macronutrient composition of the meal. Many CYP with T1D choose to consume high GI ready-toeat cereals for breakfast. This study provides evidence of the extent of this and the association between these cereals and high levels of postprandial hyperglycaemia, and it supports the use of protein to manage postprandial glycaemia after breakfast in CYP with T1D.

The impact of increasing protein intake at breakfast in T1D requires further investigation in order to establish the amount of protein required to dampen the glucose rise after breakfast and to establish the optimal breakfast meal composition. One such intervention study has now been undertaken with results pending. Given their ability to improve overall glycaemic control, HCL systems may be better able to manage postprandial hyperglycaemia regardless of meal composition and further research is required to establish their effectiveness in managing postprandial glycaemia after breakfast in CYP with T1D.

#### Data Availability Statement

The data to support these findings are available from the corresponding author on request.

#### **Ethics Statement**

The study was approved by the University of Stirling (NHS, Invasive & Clinical Research Committee) and the West of Scotland (5) Research Ethics Committee (20/WS/0123).

#### Disclosure

This paper formed part of a Clinical Doctorate thesis for Julie Johnson titled 'The Breakfast Rise Education and Knowledge Study (The BREAK study).

#### **Conflicts of Interest**

The authors declare no conflicts of interest.

#### **Author Contributions**

Julie Johnson: Conceptualization, data curation, formal analysis, investigation, methodology, project administration, resources, visualization, writing—original draft, writing review and editing. Victoria Franklin: Conceptualization, methodology, supervision, writing—review and editing. Ashley Shepherd: Conceptualization, methodology, project administration, supervision, writing—review and editing. Grace Chau: Investigation, project administration, writing—review and editing. Kate Keen: Investigation, project administration, writing—review and editing. Sophie Lennon: Investigation, project administration, writing—review and editing. Maria Leveridge: Investigation, project administration, writing review and editing. Kirsty Maclean: Investigation, project administration, writing—review and editing. Julie Nicol: Investigation, project administration, writing—review and editing. Vanessa Phillipson: Investigation, project administration, writing—review and editing. Sue Roach: Investigation, project administration, writing—review and editing. Adele Swart: Investigation, project administration, writing—review and editing. Stuart Galloway: Conceptualization, writing review and editing. supervision, validation, writing review and editing.

#### Funding

No funding was received for this research.

#### Acknowledgments

We wish to thank all the children and young people and their families who participated in this study. The abstract of this study was presented at the 50<sup>th</sup> Annual Conference of the ISPAD (2024) [88].

#### References

- D. Rodbard, "Glucose Variability: A Review of Clinical Applications and Research Developments," *Diabetes Technol*ogy & *Therapeutics* 20, no. S2 (2018): S2-5–S2-15.
- [2] M. J. Rewers, K. Pillay, C. De Beaufort, et al., "Assessment and Monitoring of Glycemic Control in Children and Adolescents With Diabetes," *Pediatric Diabetes* 15, no. S20 (2014): 102– 114.
- [3] T. S. Temelkova-Kurktschiev, C. Koehler, E. Henkel, W. Leonhardt, K. Fuecker, and M. Hanefeld, "Postchallenge Plasma Glucose and Glycemic Spikes Are More Strongly Associated With Atherosclerosis Than Fasting Glucose or HbA1c Level," *Diabetes Care* 23, no. 12 (2000): 1830–1834.
- [4] M. Shichiri, H. Kishikawa, Y. Ohkubo, and N. Wake, "Long-Term Results of the Kumamoto Study on Optimal Diabetes Control in Type 2 Diabetic Patients," *Diabetes Care* 23, no. 2 (2000): B21-9.
- [5] F. G. Banting, W. R. Campbell, and A. A. Fletcher, "Further Clinical Experience With Insulin (Pancreatic Extracts) in the Treatment of Diabetes Mellitus," *BMJ* 1, no. 3236 (1923): 8–12.
- [6] E. Boland, T. Monsod, M. Delucia, C. A. Brandt, S. Fernando, and W. V. Tamborlane, "Limitations of Conventional Methods of Self-Monitoring of Blood Glucose: Lessons Learned From 3 Days of Continuous Glucose Sensing in Pediatric Patients With Type 1 Diabetes," *Diabetes Care* 24, no. 11 (2001): 1858– 1862.
- [7] M. Tansey, R. Beck, K. Ruedy, et al., "Persistently High Glucose Levels in Young Children With Type 1 Diabetes," *Pediatric Diabetes* 17, no. 2 (2016): 93–100.
- [8] Diabetes Research in Children Network (DirecNet) Study Group, "Eight-Point Glucose Testing Versus the Continuous Glucose Monitoring System in Evaluation of Glycemic Control in Type 1 Diabetes," *the Journal of Clinical Endocrinology and Metabolism* 90, no. 6 (2005): 3387–3391.
- [9] L. M. Gandrud, D. Xing, C. Kollman, et al., "The Medtronic MiniMed Gold Continuous Glucose Monitoring System: An Effective Means to Discover Hypo- and Hyperglycemia in

Children Under 7 Years of Age," *Diabetes Technology & Therapeutics* 9, no. 4 (2007): 307–316.

- [10] A. D. Monzon, L. B. Smith, S. W. Powers, L. M. Dolan, and S. R. Patton, "The Association Between Glycemic Variability and Macronutrients in Young Children With T1D," *Journal of Pediatric Psychology* 45, no. 7 (2020): 749–758.
- [11] D. D. Martin, E. A. Davis, and T. W. Jones, "Acute Effects of Hyperglycaemia in Children With Type 1 Diabetes Mellitus: The Patient's Perspective," *Journal of Pediatric Endocrinology* and Metabolism 19, no. 7 (2006): 927–936.
- [12] R. Rabasa-Lhoret, J. Garon, H. Langelier, D. Poisson, and J. L. Chiasson, "Effects of Meal Carbohydrate Content on Insulin Requirements in Type 1 Diabetic Patients Treated Intensively With the Basal-Bolus (Ultralente-Regular) Insulin Regimen," *Diabetes Care* 22, no. 5 (1999): 667–673.
- [13] S. F. Annan, L. A. Higgins, E. Jelleryd, et al., "ISPAD Clinical Practice Consensus Guidelines 2022: Nutritional Management in Children and Adolescents With Diabetes," *Pediatric Diabetes* 23, no. 8 (2022): 1297–1321.
- [14] B. McKevith and A. Jarzebowska, "The Role of Breakfast Cereals in the UK Diet: Headline Results From the National Diet and Nutrition Survey (NDNS) Year 1," *Nutrition Bulletin* 35 (2010): 314–319.
- [15] J. A. Grieger, S. Kim, and L. Cobiac, "Where Do Australian Children Get Their Dietary Fibre? A Focus on Breakfast Food Choices," *Nutrition & Dietetics* 70, no. 2 (2013): 132–138.
- [16] K. J. Bell, C. E. Smart, G. M. Steil, et al., "Impact of Fat, Protein, and Glycemic Index on Postprandial Glucose Control in Type 1 Diabetes: Implications for Intensive Diabetes Management in the Continuous Glucose Monitoring Era," *Diabetes Care* 38 (2015): 1008–1015.
- [17] F. S. Atkinson, K. Foster-Powell, and J. C. Brand-Miller, "International Tables of Glycemic Index and Glycemic Load Values," *Diabetes Care* 31, no. 12 (2008): 2281–2283.
- [18] F. S. Atkinson, J. C. Brand-Miller, K. Foster-Powell, A. E. Buyken, and J. Goletzke, "International Tables of Glycemic Index and Glycemic Load Values 2021: A Systematic Review," *The American Journal of Clinical Nutrition* 114, no. 5 (2021): 1625–1632.
- [19] H. Dodd, S. Williams, R. Brown, and B. Venn, "Calculating Meal Glycemic Index by Using Measured and Published Food Values Compared With Directly Measured Meal Glycemic Index," *The American Journal of Clinical Nutrition* 94, no. 4 (2011): 992–996.
- [20] F. R. Bornet, D. Costagliola, S. W. Rizkalla, et al., "Insulinemic and Glycemic Indexes of Six Starch-Rich Foods Taken Alone and in a Mixed Meal by Type 2 Diabetics," *The American Journal of Clinical Nutrition* 45, no. 3 (1987): 588–595.
- [21] I. Chew, J. C. Brand, A. W. Thorburn, and A. S. Truswell, "Application of Glycemic Index to Mixed Meals," *The American Journal of Clinical Nutrition* 47, no. 1 (1988): 53–56.
- [22] T. M. S. Wolever, M. Yang, X. Y. Zeng, F. Atkinson, and J. C. Brand-Miller, "Food Glycemic Index, as Given in Glycemic Index Tables, Is a Significant Determinant of Glycemic Responses Elicited by Composite Breakfast Meals," *The American Journal of Clinical Nutrition* 83, no. 6 (2006): 1306–1312.
- [23] J. Salmerón, A. Ascherio, E. B. Rimm, et al., "Dietary Fiber, Glycemic Load, and Risk of NIDDM in Men," *Diabetes Care* 20, no. 4 (1997): 545–550.
- [24] The Eatwell Guide, 2016, Accessed 16 Feb. 2025 https://www. gov.uk/government/publications/the-eatwell-guide.
- [25] T. Battelino, T. Danne, R. M. Bergenstal, et al., "Clinical Targets for Continuous Glucose Monitoring Data Interpretation: Recommendations From the International Consensus on Time in Range," *Diabetes Care* 42, no. 8 (2019): 1593–1603.

- [26] O. Schnell, K. Barnard, R. Bergenstal, et al., "Role of Continuous Glucose Monitoring in Clinical Trials: Recommendations on Reporting," *Diabetes Technology & Therapeutics* 19, no. 7 (2017): 391–399.
- [27] M. Brysbaert and M. Stevens, "Power Analysis and Effect Size in Mixed Effects Models: A Tutorial," *Journal of Cognition* 1, no. 1 (2018).
- [28] J. Westfall, D. A. Kenny, and C. M. Judd, "Statistical Power and Optimal Design in Experiments in Which Samples of Participants Respond to Samples of Stimuli," *Journal of Experimental Psychology: General* 143, no. 5 (2014): 2020– 2045.
- [29] L. Monnier, C. Colette, A. Wojtusciszyn, et al., "Toward Defining the Threshold Between Low and High Glucose Variability in Diabetes," *Diabetes Care* 40, no. 7 (2017): 832– 838.
- [30] J. Edge, C. Acerini, F. Campbell, et al., "An Alternative Sensor-Based Method for Glucose Monitoring in Children and Young People With Diabetes," *Archives of Disease in Childhood* 102, no. 6 (2017): 543–549.
- [31] J. Petersson, K. Åkesson, F. Sundberg, and S. Särnblad, "Translating Glycated Hemoglobin A1c into Time Spent in Glucose Target Range: A Multicenter Study," *Pediatric Diabetes* 20, no. 3 (2019): 339–344.
- [32] A. Brener, K. Mazor-Aronovitch, M. Rachmiel, et al., "Lessons Learned From the Continuous Glucose Monitoring Metrics in Pediatric Patients With Type 1 Diabetes Under COVID-19 Lockdown," Acta Diabetologica 57, no. 12 (2020): 1511–1517.
- [33] C. Piona, M. Marigliano, E. Mozzillo, et al., "Long-Term Glycemic Control and Glucose Variability Assessed With Continuous Glucose Monitoring in a Pediatric Population With Type 1 Diabetes: Determination of Optimal Sampling Duration," *Pediatric Diabetes* 21, no. 8 (2020): 1485–1492.
- [34] C. Piona, M. Marigliano, V. Mancioppi, et al., "Glycemic Variability and Time in Range Are Associated With the Risk of Overweight and High LDL-Cholesterol in Children and Youths With Type 1 Diabetes," *Hormone Research in Paediatrics* (2023): 1–12.
- [35] F. Lombardo, G. Salzano, B. Bombaci, et al., "Has COVID-19 Lockdown Improved Glycaemic Control in Pediatric Patients With Type 1 Diabetes? An Analysis of Continuous Glucose Monitoring Metrics," *Diabetes Research and Clinical Practice* 178 (2021): 108988.
- [36] M. Marigliano, R. Pertile, E. Mozzillo, et al., "Satisfaction With Continuous Glucose Monitoring Is Positively Correlated With Time in Range in Children With Type 1 Diabetes," *Diabetes Research and Clinical Practice* 204 (2023): 110895.
- [37] S. Foti Randazzese, B. Bombaci, S. Costantino, Y. Giorgianni, F. Lombardo, and G. Salzano, "Discordance Between Glucose Management Indicator and Glycated Hemoglobin in a Pediatric Cohort With Type 1 Diabetes: A Real-World Study," *Children* 11, no. 2 (2024): 210.
- [38] K. K. Kowalczyk-Korcz and A. Szypowska, "Factors Affecting the Prolongation of Glycemic Time in Range Among Children With Type 1 Diabetes Using Continuous Glucose Monitoring Systems: A Case Control Study," *International Journal of Diabetes in Developing Countries* (2024): 1–9.
- [39] B. Karges, A. Schwandt, B. Heidtmann, et al., "Association of Insulin Pump Therapy vs Insulin Injection Therapy With Severe Hypoglycemia, Ketoacidosis, and Glycemic Control Among Children, Adolescents, and Young Adults With Type 1 Diabetes," JAMA 318, no. 14 (2017): 1358–1366.

- [40] R. Hanas and J. Ludvigsson, "Hypoglycemia and Ketoacidosis With Insulin Pump Therapy in Children and Adolescents," *Pediatric Diabetes* 7, no. s4 (2006): 32–38.
- [41] V. Cherubini, R. Bonfanti, A. Casertano, et al., "Time In Range in Children With Type 1 Diabetes Using Treatment Strategies Based on Nonautomated Insulin Delivery Systems in the Real World," *Diabetes Technology & Therapeutics* 22, no. 7 (2020): 509–515.
- [42] T. Battelino, M. Phillip, N. Bratina, R. Nimri, P. Oskarsson, and J. Bolinder, "Effect of Continuous Glucose Monitoring on Hypoglycemia in Type 1 Diabetes," *Diabetes Care* 34, no. 4 (2011): 795–800.
- [43] K. Dovc, S. Lanzinger, R. Cardona-Hernandez, et al., "Association of Achieving Time in Range Clinical Targets With Treatment Modality Among Youths With Type 1 Diabetes," *JAMA Network Open* 6, no. 2 (2023): e230077.
- [44] K. Nagl, I. Bozic, G. Berger, et al., "Time in Range in Children With Type 1 Diabetes Before and During a Diabetes Camp—A Ceiling Effect?" *Children* 9, no. 12 (2022): 1951.
- [45] S. A. Brown, B. P. Kovatchev, D. Raghinaru, et al., "Six-Month Randomized, Multicenter Trial of Closed-Loop Control in Type 1 Diabetes," *New England Journal of Medicine* 381, no. 18 (2019): 1707–1717.
- [46] M. D. Breton, L. G. Kanapka, R. W. Beck, et al., "A Randomized Trial of Closed-Loop Control in Children With Type 1 Diabetes," *New England Journal of Medicine* 383, no. 9 (2020): 836–845.
- [47] R. M. Bergenstal, R. Nimri, R. W. Beck, et al., "A Comparison of Two Hybrid Closed-Loop Systems in Adolescents and Young Adults With Type 1 Diabetes (FLAIR): A Multicentre, Randomised, Crossover Trial," *The Lancet* 397, no. 10270 (2021): 208–219.
- [48] J. Ware, C. K. Boughton, J. M. Allen, et al., "Cambridge Hybrid Closed-Loop Algorithm in Children and Adolescents With Type 1 Diabetes: A Multicentre 6-Month Randomised Controlled Trial," *The Lancet Digital Health* 4, no. 4 (2022): e245–e255.
- [49] T. von dem Berge, K. Remus, S. Biester, et al., "In-Home Use of a Hybrid Closed Loop Achieves Time-in-Range Targets in Preschoolers and School Children: Results From a Randomized, Controlled, Crossover Trial," *Diabetes, Obesity and Metabolism* 24, no. 7 (2022): 1319–1327.
- [50] R. A. Heptulla, H. F. Allen, T. M. Gross, and E. O. Reiter, "Continuous Glucose Monitoring in Children With Type 1 Diabetes: Before and After Insulin Pump Therapy," *Pediatric Diabetes* 5, no. 1 (2004): 10–15.
- [51] E. Bell, S. Binkowski, E. Sanderson, et al., "Substantial Intra-Individual Variability in Post-Prandial Time to Peak in Controlled and Free-Living Conditions in Children With Type 1 Diabetes," *Nutrients* 13, no. 11 (2021): 4154.
- [52] M. I. Schmidt, A. Hadji-Georgopoulos, M. Rendell, S. Margolis, and A. Kowarski, "The Dawn Phenomenon, an Early Morning Glucose Rise: Implications for Diabetic Intraday Blood Glucose Variation," *Diabetes Care* 4, no. 6 (1981): 579–585.
- [53] G. P. Forlenza, L. Ekhlaspour, L. A. DiMeglio, et al., "Glycemic Outcomes of Children 2–6 Years of Age With Type 1 Diabetes During the Pediatric MiniMedTM 670G System Trial," *Pediatric Diabetes* 23, no. 3 (2022): 324–329.
- [54] M. F. Carroll, A. Izard, K. Riboni, M. R. Burge, and D. S. Schade, "Fasting Hyperglycemia Predicts the Magnitude of Postprandial Hyperglycemia," *Diabetes Care* 25, no. 7 (2002): 1247-1248.

- [55] H. Nielsen and G. L. Nielsen, "Preprandial Blood Glucose Values: Influence on Glycemic Response Studies," *The American Journal* of Clinical Nutrition 49, no. 6 (1989): 1243–1246.
- [56] A. R. Majithia, A. B. Wiltschko, H. Zheng, G. A. Walford, and D. M. Nathan, "Rate of Change of Premeal Glucose Measured by Continuous Glucose Monitoring Predicts Postmeal Glycemic Excursions in Patients With Type 1 Diabetes: Implications for Therapy," *Journal of Diabetes Science and Technology* 12, no. 1 (2018): 76–82.
- [57] E. Cengiz, T. Danne, T. Ahmad, et al., "ISPAD Clinical Practice Consensus Guidelines 2022: Insulin Treatment in Children and Adolescents With Diabetes," *Pediatric Diabetes* 23, no. 8 (2022): 1277–1296.
- [58] A. Peters, M. A. Van Name, B. L. Thorsted, J. S. Piltoft, and W. V. Tamborlane, "Postprandial Dosing of Bolus Insulin in Patients With Type 1 Diabetes: A Cross-sectional Study Using Data From the T1D Exchange Registry," *Endocrine Practice* 23, no. 10 (2017): 1201–1209.
- [59] K. A. Datye, C. T. Boyle, J. Simmons, et al., "Timing of Meal Insulin and Its Relation to Adherence to Therapy in Type 1 Diabetes," *Journal of Diabetes Science and Technology* 12, no. 2 (2018): 349–355.
- [60] E. Mozzillo, R. Franceschi, F. Di Candia, et al., "Optimal Prandial Timing of Insulin Bolus in Youths With Type 1 Diabetes: A Systematic Review," *Journal of Personalized Medicine* 12, no. 12 (2022): 2058.
- [61] G. Slama, F. Elgrably, A. Sola, J. Mbemba, and E. Larger, "Postprandial Glycaemia: A Plea for the Frequent Use of Delta Postprandial Glycaemia in the Treatment of Diabetic Patients," *Diabetes & Metabolism* 32, no. 2 (2006): 187–192.
- [62] N. C. Foster, R. W. Beck, K. M. Miller, et al., "State of Type 1 Diabetes Management and Outcomes From the T1D Exchange in 2016–2018," *Diabetes Technology & Therapeutics* 21, no. 2 (2019): 66–72.
- [63] E. R. Mackey, M. Rose, C. Tully, et al., "The Current State of Parent Feeding Behavior, Child Eating Behavior, and Nutrition Intake in Young Children With Type 1 Diabetes," *Pediatric Diabetes* 21, no. 5 (2020): 841–845.
- [64] E. L. Fisher, N. A. Weaver, A. L. Marlow, B. R. King, and C. E. Smart, "Macronutrient Intake in Children and Adolescents With Type 1 Diabetes and Its Association With Glycemic Outcomes," *Pediatric Diabetes* 2023 (2023): 7102890.
- [65] P. Dyson, "Low Carbohydrate Diets and Type 2 Diabetes: What Is the Latest Evidence?" *Diabetes Therapy* 6, no. 4 (2015): 411–424.
- [66] R. Seckold, E. Fisher, M. de Bock, B. R. King, and C. E. Smart, "The Ups and Downs of Low-Carbohydrate Diets in the Management of Type 1 Diabetes: A Review of Clinical Outcomes," *Diabetic Medicine* 36, no. 3 (2019): 326–334.
- [67] M. de Bock, K. Lobley, D. Anderson, et al., "Endocrine and Metabolic Consequences due to Restrictive Carbohydrate Diets in Children With Type 1 Diabetes: An Illustrative Case Series," *Pediatric Diabetes* 19, no. 1 (2018): 129–137.
- [68] L. Lafrance, R. Rabasa-Lhoret, D. Poisson, F. Ducros, and J.-L. Chiasson, "Effects of Different GI Foods and Dietary Fibre Intake on Glycaemic Control in Type 1 Diabetic Patients on Intensive Insulin Therapy," *Diabetic Medicine* 15, no. 11 (1998): 972–978.
- [69] N. H. Mohammed and T. M. S. Wolever, "Effect of Carbohydrate Source on Post-Prandial Blood Glucose in Subjects With Type 1 Diabetes Treated With Insulin Lispro," *Diabetes Research and Clinical Practice* 65, no. 1 (2004): 29–35.

- [70] T. R. Nansel, L. Gellar, and A. McGill, "Effect of Varying Glycemic Index Meals on Blood Glucose Control Assessed With Continuous Glucose Monitoring in Youth With Type 1 Diabetes on Basal-Bolus Insulin Regimens," *Diabetes Care* 31, no. 4 (2008): 695–697.
- [71] A. H. El-Laboudi, I. F. Godsland, D. G. Johnston, and N. S. Oliver, "Measures of Glycemic Variability in Type 1 Diabetes and the Effect of Real-Time Continuous Glucose Monitoring," *Diabetes Technology & Therapeutics* 18, no. 12 (2016): 806–812.
- [72] R. Birnbacher, T. Waldhör, U. Schneider, and E. Schober, "Glycaemic Responses to Commonly Ingested Breakfasts in Children With Insulin-Dependent Diabetes Mellitus," *European Journal of Pediatrics* 154, no. 5 (1995): 353–355.
- [73] K. Dżygało and A. Szypowska, "Impact of Insulins Glulisine and Aspart on Postprandial Glycemia After a High-Glycemic Index Meal in Children With Type 1 Diabetes," *European Journal of Endocrinology* 170, no. 4 (2014): 539–545.
- [74] L. Groele, D. Golicki, M. Błazik, and E. Pańkowska, "Improving the Estimation of Meal-Time Insulin Dose Based on the Glycaemic Load of a Meal in Children With Type 1 Diabetes on Insulin Pump Therapy: A Randomized Study," *The Journal of Diabetes & Metabolism* 5, no. 435 (2014): 2.
- [75] T. A. Smith, A. A. Blowes, B. R. King, P. P. Howley, and C. E. Smart, "Families' Reports of Problematic Foods, Management Strategies and Continuous Glucose Monitoring in Type 1 Diabetes: A Cross-Sectional Study," *Nutrition & Dietetics* 78, no. 4 (2021): 449–457.
- [76] R. L. Ryan, B. R. King, D. G. Anderson, J. R. Attia, C. E. Collins, and C. E. Smart, "Influence of and Optimal Insulin Therapy for a Low-Glycemic Index Meal in Children With Type 1 Diabetes Receiving Intensive Insulin Therapy," *Diabetes Care* 31, no. 8 (2008): 1485–1490.
- [77] M. A. O'Connell, H. R. Gilbertson, S. M. Donath, and F. J. Cameron, "Optimizing Postprandial Glycemia in Pediatric Patients With Type 1 Diabetes Using Insulin Pump Therapy: Impact of Glycemic Index and Prandial Bolus Type," *Diabetes Care* 31, no. 8 (2008): 1491–1495.
- [78] M. A. Paterson, C. E. M. Smart, P. E. Lopez, et al., "Increasing the Protein Quantity in a Meal Results in Dose-Dependent Effects on Postprandial Glucose Levels in Individuals With Type 1 Diabetes Mellitus," *Diabetic Medicine* 34, no. 6 (2017): 851–854.
- [79] M. A. Paterson, B. R. King, C. E. M. Smart, T. Smith, J. Rafferty, and P. E. Lopez, "Impact of Dietary Protein on Postprandial Glycaemic Control and Insulin Requirements in Type 1 Diabetes: A Systematic Review," *Diabetic Medicine* 36, no. 12 (2019): 1585–1599.
- [80] V. Cherubini, M. Marino, M. Marigliano, et al., "Rethinking Carbohydrate Intake and Time in Range in Children and Adolescents With Type 1 Diabetes," *Nutrients* 13, no. 11 (2021): 3869.
- [81] K. A. Hätönen, J. Virtamo, J. G. Eriksson, H. K. Sinkko, J. E. Sundvall, and L. M. Valsta, "Protein and Fat Modify the Glycaemic and Insulinaemic Responses to a Mashed Potato-Based Meal," *British Journal of Nutrition* 106, no. 2 (2011): 248–253.
- [82] C. E. Smart, M. Evans, S. M. O'connell, et al., "Both Dietary Protein and Fat Increase Postprandial Glucose Excursions in Children With Type 1 Diabetes, and the Effect Is Additive," *Diabetes Care* 36, no. 12 (2013): 3897–3902.
- [83] M. A. Paterson, C. E. M. Smart, P. Howley, D. A. Price, D. C. Foskett, and B. R. King, "High-Protein Meals Require

30% Additional Insulin to Prevent Delayed Postprandial Hyperglycaemia," *Diabetic Medicine* 37, no. 7 (2020): 1185– 1191.

- [84] M. van der Hoogt, J. C. van Dyk, R. C. Dolman, and M. Pieters, "Protein and Fat Meal Content Increase Insulin Requirement in Children With Type 1 Diabetes—Role of Duration of Diabetes," *Journal of Clinical & Translational Endocrinology* 10 (2017): 15–21.
- [85] H. Sacks, T. C. Chalmers, and H. Smith Jr., "Randomized Versus Historical Controls for Clinical Trials," *The American Journal of Medicine* 72, no. 2 (1982): 233–240.
- [86] R. J. Ligthelm, V. Borzì, J. Gumprecht, R. Kawamori, Y. Wenying, and P. Valensi, "Importance of Observational Studies in Clinical Practice," *Clinical Therapeutics* 29, no. 6 (2007): 1284–1292.
- [87] A. Addala, M. Auzanneau, K. Miller, et al., "A Decade of Disparities in Diabetes Technology Use and HbA1c in Pediatric Type 1 Diabetes: A Transatlantic Comparison," *Diabetes Care* 44, no. 1 (2021): 133–140.
- [88] J. Johnson, V. Franklin, A. Shepherd, S. D. R. Galloway, and W. Syplie, "Glucose Variability and Glycaemic Responses to Food at Breakfast in Children and Young People With Type 1 Diabetes. Abstract Book 50th Annual Conference ISPAD 2024," *Hormone Research in Paediatrics* 97, no. Suppl. 2 (2024): 1–370.